wwPDB X-ray Structure Validation Summary Report (i)

May 23, 2020 – 07:39 am BST

PDB ID	:	4MMN
Title	:	Structural and biochemical analysis of type II free methionine-R-sulfoxide reductase from Thermoplasma acidophilum
Authors	:	Kim, H.S.; Kwak, G.H.; Lee, K.T.; Jo, C.H.; Hwang, K.Y.; Kim, H.Y.
Deposited on	:	2013-09-09
Resolution	:	2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	78% 17% 5%
1	B	149	% 74% 19% 6%
1	C	149	78% 16% 6%
1	D	149	77% 18% 5%
1	E	149	75% 19% 5%
1	F	149	81% 14% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	149	79% 14% • 6%
1	H	149	78% 15% • 6%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 9089 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative uncharacterized protein Ta0848.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	141	1118	718	178	216	6	0	0	0
			Total	C	N	O	S			
1	B	140	1110	712	177	215	6	0	0	0
			Total	C	N	O	S			
1	C	140	1110	712	177	215	6	0	0	0
			Total	C	N	O	S			
1	D	141	1118	718	178	216	6	0	0	0
			Total	C	N	O	S			
1	E	141	1118	718	178	216	6	0	0	0
			Total	C	N	O	S			
1	F	141	1118	718	178	216	6	0	0	0
			Total	C	N	O	S			
1	G	140	1110	712	177	215	6	0	0	0
			Total	C	N	O	S			
1	H	140	1110	712	177	215	6	0	0	0

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	142	LEU	-	EXPRESSION TAG	UNP Q9HJW5
A	143	GLU	-	EXPRESSION TAG	UNP Q9HJW5
A	144	HIS	-	EXPRESSION TAG	UNP Q9HJW5
A	145	HIS	-	EXPRESSION TAG	UNP Q9HJW5
A	146	HIS	-	EXPRESSION TAG	UNP Q9HJW5
A	147	HIS	-	EXPRESSION TAG	UNP Q9HJW5
A	148	HIS	-	EXPRESSION TAG	UNP Q9HJW5
A	149	HIS	-	EXPRESSION TAG	UNP Q9HJW5
B	142	LEU	-	EXPRESSION TAG	UNP Q9HJW5
B	143	GLU	-	EXPRESSION TAG	UNP Q9HJW5
B	144	HIS	-	EXPRESSION TAG	UNP Q9HJW5
B	145	HIS	-	EXPRESSION TAG	UNP Q9HJW5
B	146	HIS	-	EXPRESSION TAG	UNP Q9HJW5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	147	HIS	-	EXPRESSION TAG	UNP Q9HJW5
B	148	HIS	-	EXPRESSION TAG	UNP Q9HJW5
B	149	HIS	-	EXPRESSION TAG	UNP Q9HJW5
C	142	LEU	-	EXPRESSION TAG	UNP Q9HJW5
C	143	GLU	-	EXPRESSION TAG	UNP Q9HJW5
C	144	HIS	-	EXPRESSION TAG	UNP Q9HJW5
C	145	HIS	-	EXPRESSION TAG	UNP Q9HJW5
C	146	HIS	-	EXPRESSION TAG	UNP Q9HJW5
C	147	HIS	-	EXPRESSION TAG	UNP Q9HJW5
C	148	HIS	-	EXPRESSION TAG	UNP Q9HJW5
C	149	HIS	-	EXPRESSION TAG	UNP Q9HJW5
D	142	LEU	-	EXPRESSION TAG	UNP Q9HJW5
D	143	GLU	-	EXPRESSION TAG	UNP Q9HJW5
D	144	HIS	-	EXPRESSION TAG	UNP Q9HJW5
D	145	HIS	-	EXPRESSION TAG	UNP Q9HJW5
D	146	HIS	-	EXPRESSION TAG	UNP Q9HJW5
D	147	HIS	-	EXPRESSION TAG	UNP Q9HJW5
D	148	HIS	-	EXPRESSION TAG	UNP Q9HJW5
D	149	HIS	-	EXPRESSION TAG	UNP Q9HJW5
E	142	LEU	-	EXPRESSION TAG	UNP Q9HJW5
E	143	GLU	-	EXPRESSION TAG	UNP Q9HJW5
E	144	HIS	-	EXPRESSION TAG	UNP Q9HJW5
E	145	HIS	-	EXPRESSION TAG	UNP Q9HJW5
E	146	HIS	-	EXPRESSION TAG	UNP Q9HJW5
E	147	HIS	-	EXPRESSION TAG	UNP Q9HJW5
E	148	HIS	-	EXPRESSION TAG	UNP Q9HJW5
E	149	HIS	-	EXPRESSION TAG	UNP Q9HJW5
F	142	LEU	-	EXPRESSION TAG	UNP Q9HJW5
F	143	GLU	-	EXPRESSION TAG	UNP Q9HJW5
F	144	HIS	-	EXPRESSION TAG	UNP Q9HJW5
F	145	HIS	-	EXPRESSION TAG	UNP Q9HJW5
F	146	HIS	-	EXPRESSION TAG	UNP Q9HJW5
F	147	HIS	-	EXPRESSION TAG	UNP Q9HJW5
F	148	HIS	-	EXPRESSION TAG	UNP Q9HJW5
F	149	HIS	-	EXPRESSION TAG	UNP Q9HJW5
G	142	LEU	-	EXPRESSION TAG	UNP Q9HJW5
G	143	GLU	-	EXPRESSION TAG	UNP Q9HJW5
G	144	HIS	-	EXPRESSION TAG	UNP Q9HJW5
G	145	HIS	-	EXPRESSION TAG	UNP Q9HJW5
G	146	HIS	-	EXPRESSION TAG	UNP Q9HJW5
G	147	HIS	-	EXPRESSION TAG	UNP Q9HJW5
G	148	HIS	-	EXPRESSION TAG	UNP Q9HJW5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	149	HIS	-	EXPRESSION TAG	UNP Q9HJW5
H	142	LEU	-	EXPRESSION TAG	UNP Q9HJW5
H	143	GLU	-	EXPRESSION TAG	UNP Q9HJW5
H	144	HIS	-	EXPRESSION TAG	UNP Q9HJW5
H	145	HIS	-	EXPRESSION TAG	UNP Q9HJW5
H	146	HIS	-	EXPRESSION TAG	UNP Q9HJW5
H	147	HIS	-	EXPRESSION TAG	UNP Q9HJW5
H	148	HIS	-	EXPRESSION TAG	UNP Q9HJW5
H	149	HIS	-	EXPRESSION TAG	UNP Q9HJW5

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	16	16	16	0	0
			Total	O		
2	B	18	18	18	0	0
			Total	O		
2	C	29	29	29	0	0
			Total	O		
2	D	20	20	20	0	0
			Total	O		
2	E	22	22	22	0	0
			Total	O		
2	F	17	17	17	0	0
			Total	O		
2	G	26	26	26	0	0
			Total	O		
2	H	29	29	29	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative uncharacterized protein Ta0848

Chain A:

- Molecule 1: Putative uncharacterized protein Ta0848

Chain B:



- Molecule 1: Putative uncharacterized protein Ta0848

Chain C:



- Molecule 1: Putative uncharacterized protein Ta0848

Chain D:



- Molecule 1: Putative uncharacterized protein Ta0848

Chain E:



- Molecule 1: Putative uncharacterized protein Ta0848

Chain F:



- Molecule 1: Putative uncharacterized protein Ta0848

Chain G:



- Molecule 1: Putative uncharacterized protein Ta0848

Chain H:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α, β, γ	158.90Å 158.90Å 194.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.09 – 2.60 28.09 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.1 (28.09-2.60) 96.1 (28.09-2.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$< I/\sigma(I) >$ [1]	1.42 (at 2.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.196 , 0.240 0.196 , 0.238	Depositor DCC
R_{free} test set	1993 reflections (3.68%)	wwPDB-VP
Wilson B-factor (Å^2)	44.6	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.28 , 23.5	EDS
L-test for twinning[2]	$< \|L\| > = 0.46$, $< L^2 > = 0.28$	Xtriage
Estimated twinning fraction	0.054 for -2/3*h-1/3*k+2/3*l,-1/3*h-2/3*k-2/3*l,2/3*h-2/3*k+1/3*l 0.052 for -h,1/3*h-1/3*k+2/3*l,2/3*h+4/3*k+1/3*l 0.052 for -1/3*h+1/3*k-2/3*l,-k,-4/3*h-2/3*k+1/3*l 0.479 for -h,2/3*h+1/3*k-2/3*l,-2/3*h-4/3*k-1/3*l 0.479 for 1/3*h+2/3*k+2/3*l,-k,4/3*h+2/3*k-1/3*l 0.479 for -1/3*h-2/3*k-2/3*l,-2/3*h-1/3*k+2/3*l,-2/3*h+2/3*k-1/3*l 0.054 for h,-h-k,-l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	9089	wwPDB-VP
Average B, all atoms (Å^2)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.45	0/1141	0.59	0/1540
1	B	0.49	0/1133	0.58	0/1529
1	C	0.49	0/1133	0.59	0/1529
1	D	0.46	0/1141	0.61	0/1540
1	E	0.47	0/1141	0.60	0/1540
1	F	0.46	0/1141	0.59	0/1540
1	G	0.49	0/1133	0.56	0/1529
1	H	0.49	0/1133	0.60	1/1529 (0.1%)
All	All	0.48	0/9096	0.59	1/12276 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	H	82	LEU	CA-CB-CG	5.90	128.87	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1118	0	1105	18	0
1	B	1110	0	1094	17	0
1	C	1110	0	1094	15	0
1	D	1118	0	1105	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1118	0	1105	20	0
1	F	1118	0	1105	15	0
1	G	1110	0	1094	15	0
1	H	1110	0	1094	15	0
2	A	16	0	0	1	0
2	B	18	0	0	2	0
2	C	29	0	0	3	0
2	D	20	0	0	1	0
2	E	22	0	0	3	0
2	F	17	0	0	1	0
2	G	26	0	0	2	0
2	H	29	0	0	1	0
All	All	9089	0	8796	134	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 134 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:88:THR:HG21	1:H:109:ASP:HB3	1.63	0.78
1:B:88:THR:HG21	1:B:109:ASP:HB3	1.65	0.78
1:F:60:CYS:SG	2:F:211:HOH:O	2.40	0.78
1:G:88:THR:HG21	1:G:109:ASP:HB3	1.62	0.78
1:C:88:THR:HG21	1:C:109:ASP:HB3	1.66	0.76

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/149 (93%)	134 (96%)	5 (4%)	0	100	100
1	B	138/149 (93%)	135 (98%)	3 (2%)	0	100	100
1	C	138/149 (93%)	136 (99%)	2 (1%)	0	100	100
1	D	139/149 (93%)	137 (99%)	2 (1%)	0	100	100
1	E	139/149 (93%)	135 (97%)	4 (3%)	0	100	100
1	F	139/149 (93%)	134 (96%)	5 (4%)	0	100	100
1	G	138/149 (93%)	137 (99%)	1 (1%)	0	100	100
1	H	138/149 (93%)	135 (98%)	3 (2%)	0	100	100
All	All	1108/1192 (93%)	1083 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/131 (94%)	123 (100%)	0	100	100
1	B	122/131 (93%)	121 (99%)	1 (1%)	81	92
1	C	122/131 (93%)	120 (98%)	2 (2%)	62	82
1	D	123/131 (94%)	122 (99%)	1 (1%)	81	92
1	E	123/131 (94%)	123 (100%)	0	100	100
1	F	123/131 (94%)	123 (100%)	0	100	100
1	G	122/131 (93%)	120 (98%)	2 (2%)	62	82
1	H	122/131 (93%)	120 (98%)	2 (2%)	62	82
All	All	980/1048 (94%)	972 (99%)	8 (1%)	81	92

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	142	LEU
1	H	57	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	69	ILE
1	C	95	PRO
1	G	55	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	10	GLN
1	D	54	ASN
1	E	54	ASN
1	F	10	GLN
1	F	54	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	141/149 (94%)	-0.25	0	100	100	52, 63, 84, 102	0
1	B	140/149 (93%)	-0.32	1 (0%)	87	86	53, 65, 81, 89	0
1	C	140/149 (93%)	-0.31	0	100	100	52, 65, 81, 91	0
1	D	141/149 (94%)	-0.30	0	100	100	52, 64, 83, 103	0
1	E	141/149 (94%)	-0.32	0	100	100	52, 63, 82, 102	0
1	F	141/149 (94%)	-0.33	0	100	100	52, 64, 83, 99	0
1	G	140/149 (93%)	-0.32	0	100	100	51, 65, 82, 93	0
1	H	140/149 (93%)	-0.31	0	100	100	52, 64, 82, 93	0
All	All	1124/1192 (94%)	-0.31	1 (0%)	95	95	51, 64, 83, 103	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	141	LYS	2.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.