



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 12, 2025 – 06:06 PM EDT

PDB ID : 9MUT / pdb_00009mut
Title : Reduced state of a turn-on thiol-disulfide redox biosensor with a fluorescence-lifetime readout
Authors : Rosen, P.; Yellen, G.; Lim, D.C.
Deposited on : 2025-01-14
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	
1	B	251	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	302	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7301 atoms, of which 3421 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	226	Total	C	H	N	O	S	0	4	0
			3501	1162	1686	303	342	8			
1	B	226	Total	C	H	N	O	S	0	3	0
			3491	1154	1683	304	344	6			

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	initiating methionine	UNP P42212
A	-12	LYS	-	expression tag	UNP P42212
A	-11	HIS	-	expression tag	UNP P42212
A	-10	HIS	-	expression tag	UNP P42212
A	-9	HIS	-	expression tag	UNP P42212
A	-8	HIS	-	expression tag	UNP P42212
A	-7	HIS	-	expression tag	UNP P42212
A	-6	HIS	-	expression tag	UNP P42212
A	-5	HIS	-	expression tag	UNP P42212
A	-4	GLY	-	expression tag	UNP P42212
A	-3	GLY	-	expression tag	UNP P42212
A	-2	ALA	-	expression tag	UNP P42212
A	-1	SER	-	expression tag	UNP P42212
A	1	VAL	-	insertion	UNP P42212
A	48	SER	CYS	conflict	UNP P42212
A	64	LEU	PHE	conflict	UNP P42212
A	?	-	SER	deletion	UNP P42212
A	66	SWG	TYR	chromophore	UNP P42212
A	?	-	GLY	deletion	UNP P42212
A	72	ALA	SER	conflict	UNP P42212
A	80	ARG	GLN	conflict	UNP P42212
A	145	GLY	-	insertion	UNP P42212
A	146	TRP	TYR	conflict	UNP P42212
A	147	CYS	ASN	conflict	UNP P42212
A	148	GLY	SER	conflict	UNP P42212

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	149	ASP	HIS	conflict	UNP P42212
A	154	THR	MET	conflict	UNP P42212
A	164	ALA	VAL	conflict	UNP P42212
A	176	GLY	SER	conflict	UNP P42212
A	205	CYS	GLN	conflict	UNP P42212
A	207	LYS	ALA	conflict	UNP P42212
A	221	PHE	LEU	conflict	UNP P42212
A	232	LEU	HIS	conflict	UNP P42212
B	-13	MET	-	initiating methionine	UNP P42212
B	-12	LYS	-	expression tag	UNP P42212
B	-11	HIS	-	expression tag	UNP P42212
B	-10	HIS	-	expression tag	UNP P42212
B	-9	HIS	-	expression tag	UNP P42212
B	-8	HIS	-	expression tag	UNP P42212
B	-7	HIS	-	expression tag	UNP P42212
B	-6	HIS	-	expression tag	UNP P42212
B	-5	HIS	-	expression tag	UNP P42212
B	-4	GLY	-	expression tag	UNP P42212
B	-3	GLY	-	expression tag	UNP P42212
B	-2	ALA	-	expression tag	UNP P42212
B	-1	SER	-	expression tag	UNP P42212
B	1	VAL	-	insertion	UNP P42212
B	48	SER	CYS	conflict	UNP P42212
B	64	LEU	PHE	conflict	UNP P42212
B	?	-	SER	deletion	UNP P42212
B	66	SWG	TYR	chromophore	UNP P42212
B	?	-	GLY	deletion	UNP P42212
B	72	ALA	SER	conflict	UNP P42212
B	80	ARG	GLN	conflict	UNP P42212
B	145	GLY	-	insertion	UNP P42212
B	146	TRP	TYR	conflict	UNP P42212
B	147	CYS	ASN	conflict	UNP P42212
B	148	GLY	SER	conflict	UNP P42212
B	149	ASP	HIS	conflict	UNP P42212
B	154	THR	MET	conflict	UNP P42212
B	164	ALA	VAL	conflict	UNP P42212
B	176	GLY	SER	conflict	UNP P42212
B	205	CYS	GLN	conflict	UNP P42212
B	207	LYS	ALA	conflict	UNP P42212
B	221	PHE	LEU	conflict	UNP P42212
B	232	LEU	HIS	conflict	UNP P42212

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			7	2	3	2		
3	A	1	Total	C	H	O	0	0
			7	2	3	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	0	0
			7	2	3	2		
3	B	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 4 is FORMIC ACID (CCD ID: FMT) (formula: CH_2O_2).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			5	1	2	2		
4	B	1	Total	C	H	O	0	0
			5	1	2	2		
4	B	1	Total	C	H	O	0	0
			5	1	2	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $\text{C}_3\text{H}_8\text{O}_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 6 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	H	O	0	0
			24	6	14	4		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	120	Total 120	O 120	0	0
7	B	88	Total 88	O 88	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein



- Molecule 1: Green fluorescent protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	49.95Å 140.84Å 129.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.68 – 2.20 26.68 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.2 (26.68-2.20) 98.4 (26.68-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.199 , 0.245 0.201 , 0.247	Depositor DCC
R_{free} test set	21655 reflections (8.55%)	wwPDB-VP
Wilson B-factor (Å ²)	33.1	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 40.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7301	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.47 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.6214e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FMT, PGE, SWG, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/1847	0.34	0/2499
1	B	0.22	0/1832	0.31	0/2477
All	All	0.23	0/3679	0.32	0/4976

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1815	1686	1748	12	0
1	B	1808	1683	1742	9	0
2	A	8	12	12	0	0
3	A	8	6	6	2	0
3	B	8	6	6	0	0
4	A	3	2	1	0	0
4	B	6	4	2	0	0
5	B	6	8	8	0	0
6	B	10	14	14	1	0
7	A	120	0	0	3	0
7	B	88	0	0	0	0
All	All	3880	3421	3539	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 21 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:GLU:OE2	3:A:302:ACT:H1	1.83	0.78
1:A:66:SWG:HE3	1:A:66:SWG:N2	2.10	0.66
3:A:302:ACT:H3	7:A:450:HOH:O	2.00	0.62
1:A:52:LYS:HD2	1:A:217:ASP:OD2	2.00	0.61
1:B:66:SWG:N2	1:B:66:SWG:HE3	2.16	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/251 (90%)	223 (99%)	3 (1%)	0	100	100
1	B	224/251 (89%)	219 (98%)	5 (2%)	0	100	100
All	All	450/502 (90%)	442 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/214 (91%)	194 (100%)	0	100	100
1	B	193/214 (90%)	193 (100%)	0	100	100
All	All	387/428 (90%)	387 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	ASN
1	A	121	ASN
1	B	121	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SWG	A	66	1	22,25,26	2.93	9 (40%)	27,35,37	2.20	9 (33%)
1	SWG	B	66	1	22,25,26	2.86	10 (45%)	27,35,37	2.55	10 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SWG	A	66	1	-	5/8/29/30	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SWG	B	66	1	-	5/8/29/30	0/3/3/3

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	SWG	CA2-C2	-8.46	1.39	1.48
1	B	66	SWG	CA2-C2	-8.26	1.39	1.48
1	A	66	SWG	OG1-CB1	-5.47	1.19	1.42
1	B	66	SWG	OG1-CB1	-5.45	1.19	1.42
1	A	66	SWG	CG-CB2	3.73	1.54	1.47

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	66	SWG	O2-C2-CA2	-5.56	127.47	131.02
1	B	66	SWG	CZ3-CE3-CD2	-5.32	113.73	120.91
1	B	66	SWG	CE3-CD2-CE2	4.99	124.79	118.17
1	A	66	SWG	CZ3-CE3-CD2	-4.66	114.61	120.91
1	B	66	SWG	CH2-CZ2-CE2	-4.62	113.77	120.09

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	66	SWG	C2-CA2-CB2-CG
1	A	66	SWG	N1-CA1-CB1-OG1
1	B	66	SWG	C2-CA2-CB2-CG
1	B	66	SWG	N1-CA1-CB1-OG1
1	A	66	SWG	N2-CA2-CB2-CG

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	66	SWG	2	0
1	B	66	SWG	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	B	301	-	5,5,5	0.58	0	5,5,5	1.00	0
3	ACT	B	302	-	3,3,3	1.15	0	3,3,3	1.34	0
2	EDO	A	301	-	3,3,3	0.55	0	2,2,2	0.30	0
3	ACT	B	304	-	3,3,3	1.09	0	3,3,3	1.43	0
4	FMT	B	305	-	2,2,2	0.58	0	1,1,1	0.41	0
4	FMT	B	303	-	2,2,2	0.71	0	1,1,1	0.45	0
4	FMT	A	305	-	2,2,2	0.73	0	1,1,1	0.45	0
3	ACT	A	303	-	3,3,3	1.00	0	3,3,3	1.40	0
6	PGE	B	306	-	9,9,9	0.38	0	8,8,8	0.35	0
3	ACT	A	302	-	3,3,3	0.72	0	3,3,3	1.23	0
2	EDO	A	304	-	3,3,3	0.59	0	2,2,2	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	301	-	-	0/4/4/4	-
6	PGE	B	306	-	-	5/7/7/7	-
2	EDO	A	304	-	-	0/1/1/1	-
2	EDO	A	301	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	306	PGE	O2-C3-C4-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	B	306	PGE	O3-C5-C6-O4
6	B	306	PGE	C1-C2-O2-C3
6	B	306	PGE	C3-C4-O3-C5
6	B	306	PGE	C6-C5-O3-C4

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	306	PGE	1	0
3	A	302	ACT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	225/251 (89%)	0.06	4 (1%) 67 64	22, 36, 57, 94	4 (1%)
1	B	225/251 (89%)	0.23	6 (2%) 56 53	26, 41, 67, 94	3 (1%)
All	All	450/502 (89%)	0.14	10 (2%) 62 58	22, 38, 63, 94	7 (1%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	146	TRP	4.2
1	A	3	LYS	3.6
1	B	174	ASP	3.4
1	B	147	CYS	3.3
1	A	174	ASP	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SWG	B	66	23/24	0.89	0.10	22,28,39,44	0
1	SWG	A	66	23/24	0.94	0.07	22,28,32,34	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	PGE	B	306	10/10	0.69	0.19	52,69,91,93	0
2	EDO	A	304	4/4	0.70	0.17	32,39,45,50	0
4	FMT	B	305	3/3	0.71	0.21	40,46,49,55	0
3	ACT	B	304	4/4	0.74	0.17	38,40,55,55	0
5	GOL	B	301	6/6	0.77	0.15	32,39,45,46	0
3	ACT	A	303	4/4	0.77	0.13	47,47,56,56	0
2	EDO	A	301	4/4	0.80	0.12	35,44,49,53	0
4	FMT	A	305	3/3	0.81	0.11	57,60,68,75	0
3	ACT	A	302	4/4	0.84	0.12	28,30,34,35	0
4	FMT	B	303	3/3	0.91	0.10	28,30,36,38	0
3	ACT	B	302	4/4	0.93	0.09	29,33,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.