



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 2, 2021 – 05:55 AM EDT

PDB ID : 1N3R  
Title : Biosynthesis of pteridins. Reaction mechanism of GTP cyclohydrolase I  
Authors : Rebelo, J.; Auerbach, G.; Bader, G.; Bracher, A.; Nar, H.; Hoesl, C.; Schramek, N.; Kaiser, J.; Bacher, A.; Huber, R.; Fischer, M.  
Deposited on : 2002-10-29  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	221	61% 36% ..
1	B	221	64% 33% .
1	C	221	67% 30% .
1	D	221	64% 33% ..
1	E	221	70% 29% .
1	F	221	58% 40% .
1	G	221	62% 35% .
1	H	221	61% 38% .

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain	
1	I	221	65%	33%
1	J	221	68%	30%
1	K	221	65%	33%
1	L	221	71%	29%
1	M	221	71%	25%
1	N	221	69%	28%
1	O	221	66%	32%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GTP	I	1418	-	-	X	-

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 26471 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GTP cyclohydrolase I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	221	1728	1085	307	327	9	0	0	0
1	B	221	1728	1085	307	327	9	0	0	0
1	C	221	1728	1085	307	327	9	0	0	0
1	D	221	1728	1085	307	327	9	0	0	0
1	E	220	1720	1081	305	325	9	0	0	0
1	F	221	1728	1085	307	327	9	0	0	0
1	G	221	1728	1085	307	327	9	0	0	0
1	H	221	1728	1085	307	327	9	0	0	0
1	I	221	1728	1085	307	327	9	0	0	0
1	J	221	1728	1085	307	327	9	0	0	0
1	K	221	1728	1085	307	327	9	0	0	0
1	L	221	1728	1085	307	327	9	21	0	0
1	M	221	1728	1085	307	327	9	21	0	0
1	N	221	1728	1085	307	327	9	0	0	0
1	O	221	1728	1085	307	327	9	0	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	112	SER	HIS	engineered mutation	UNP P0A6T5
B	112	SER	HIS	engineered mutation	UNP P0A6T5
C	112	SER	HIS	engineered mutation	UNP P0A6T5
D	112	SER	HIS	engineered mutation	UNP P0A6T5
E	112	SER	HIS	engineered mutation	UNP P0A6T5
F	112	SER	HIS	engineered mutation	UNP P0A6T5
G	112	SER	HIS	engineered mutation	UNP P0A6T5
H	112	SER	HIS	engineered mutation	UNP P0A6T5
I	112	SER	HIS	engineered mutation	UNP P0A6T5
J	112	SER	HIS	engineered mutation	UNP P0A6T5
K	112	SER	HIS	engineered mutation	UNP P0A6T5
L	112	SER	HIS	engineered mutation	UNP P0A6T5
M	112	SER	HIS	engineered mutation	UNP P0A6T5
N	112	SER	HIS	engineered mutation	UNP P0A6T5
O	112	SER	HIS	engineered mutation	UNP P0A6T5

- Molecule 2 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	B	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
2	D	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	E	1	Total 32	C 10	N 5	O 14	P 3	0	0
2	F	1	Total 32	C 10	N 5	O 14	P 3	0	0
2	G	1	Total 32	C 10	N 5	O 14	P 3	0	0
2	H	1	Total 32	C 10	N 5	O 14	P 3	0	0
2	I	1	Total 32	C 10	N 5	O 14	P 3	0	0
2	J	1	Total 32	C 10	N 5	O 14	P 3	0	0
2	K	1	Total 32	C 10	N 5	O 14	P 3	0	0
2	L	1	Total 32	C 10	N 5	O 14	P 3	0	0
2	M	1	Total 32	C 10	N 5	O 14	P 3	0	0
2	N	1	Total 32	C 10	N 5	O 14	P 3	0	0
2	O	1	Total 32	C 10	N 5	O 14	P 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total 2	O 2	0	0
3	B	4	Total 4	O 4	0	0
3	C	4	Total 4	O 4	0	0
3	D	7	Total 7	O 7	0	0
3	E	2	Total 2	O 2	0	0
3	F	6	Total 6	O 6	0	0
3	G	5	Total 5	O 5	0	0
3	H	6	Total 6	O 6	0	0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
3	I	4	Total O 4 4	0	0
3	J	2	Total O 2 2	0	0
3	K	7	Total O 7 7	0	0
3	L	5	Total O 5 5	0	0
3	M	13	Total O 13 13	0	0
3	N	9	Total O 9 9	0	0
3	O	3	Total O 3 3	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: GTP cyclohydrolase I



- Molecule 1: GTP cyclohydrolase I



- Molecule 1: GTP cyclohydrolase I





- Molecule 1: GTP cyclohydrolase I

Chain D: 64% 33% ..




- Molecule 1: GTP cyclohydrolase I

Chain E: 70% 29% .



- Molecule 1: GTP cyclohydrolase I

Chain F: 58% 40% .



- Molecule 1: GTP cyclohydrolase I

Chain G: 62% 35% .






- Molecule 1: GTP cyclohydrolase I



- Molecule 1: GTP cyclohydrolase I




- Molecule 1: GTP cyclohydrolase I




- Molecule 1: GTP cyclohydrolase I






- Molecule 1: GTP cyclhydrolase I

Chain L: 71% 29%




- Molecule 1: GTP cyclhydrolase I

Chain M: 71% 25%




- Molecule 1: GTP cyclhydrolase I

Chain N: 69% 28%



- Molecule 1: GTP cyclhydrolase I

Chain O: 66% 32%



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	224.47Å 313.26Å 130.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-2.80)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.218 , 0.272	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	26471	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1755	0.70	0/2377
1	B	0.38	0/1755	0.66	0/2377
1	C	0.38	0/1755	0.68	0/2377
1	D	0.40	0/1755	0.71	3/2377 (0.1%)
1	E	0.35	0/1747	0.63	0/2366
1	F	0.39	0/1755	0.67	0/2377
1	G	0.36	0/1755	0.66	0/2377
1	H	0.35	0/1755	0.65	0/2377
1	I	0.39	0/1755	0.67	1/2377 (0.0%)
1	J	0.36	0/1755	0.64	0/2377
1	K	0.39	0/1755	0.67	1/2377 (0.0%)
1	L	0.45	0/1755	0.70	1/2377 (0.0%)
1	M	0.48	0/1755	0.75	0/2377
1	N	0.42	0/1755	0.70	1/2377 (0.0%)
1	O	0.37	0/1755	0.67	1/2377 (0.0%)
All	All	0.39	0/26317	0.68	8/35644 (0.0%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	I	150	VAL	N-CA-C	-6.08	94.58	111.00
1	D	26	PRO	N-CA-C	5.65	126.80	112.10
1	D	25	PRO	C-N-CD	-5.54	108.42	120.60
1	D	150	VAL	N-CA-C	-5.54	96.05	111.00
1	K	148	PRO	N-CA-C	-5.46	97.89	112.10

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1728	0	1766	71	0
1	B	1728	0	1766	64	0
1	C	1728	0	1766	53	0
1	D	1728	0	1766	58	0
1	E	1720	0	1760	39	0
1	F	1728	0	1766	71	0
1	G	1728	0	1766	68	0
1	H	1728	0	1766	60	0
1	I	1728	0	1766	61	0
1	J	1728	0	1766	57	0
1	K	1728	0	1766	56	0
1	L	1728	0	1766	47	0
1	M	1728	0	1766	51	0
1	N	1728	0	1766	53	0
1	O	1728	0	1766	53	0
2	A	32	0	11	5	0
2	B	32	0	11	5	0
2	C	32	0	11	4	0
2	D	32	0	11	5	0
2	E	32	0	11	2	0
2	F	32	0	11	3	0
2	G	32	0	11	5	0
2	H	32	0	11	3	0
2	I	32	0	11	9	0
2	J	32	0	11	3	0
2	K	32	0	11	3	0
2	L	32	0	11	3	0
2	M	32	0	11	2	0
2	N	32	0	11	2	0
2	O	32	0	11	3	0
3	A	2	0	0	0	0
3	B	4	0	0	0	0
3	C	4	0	0	0	0
3	D	7	0	0	3	0
3	E	2	0	0	0	0
3	F	6	0	0	1	0
3	G	5	0	0	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	6	0	0	0	0
3	I	4	0	0	1	0
3	J	2	0	0	0	0
3	K	7	0	0	0	0
3	L	5	0	0	0	0
3	M	13	0	0	3	0
3	N	9	0	0	2	0
3	O	3	0	0	0	0
All	All	26471	0	26649	796	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 796 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:185:ARG:HG2	1:O:185:ARG:HH11	0.97	1.12
1:M:207:GLN:HE22	1:N:208:ASN:HB2	1.17	1.09
1:N:168:THR:HG22	1:N:170:ASN:H	1.27	0.95
1:C:168:THR:HG22	1:C:170:ASN:H	1.31	0.95
1:A:24:ARG:HG2	1:A:25:PRO:HD2	1.47	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	219/221 (99%)	196 (90%)	16 (7%)	7 (3%)	4 13
1	B	219/221 (99%)	196 (90%)	19 (9%)	4 (2%)	8 28
1	C	219/221 (99%)	197 (90%)	18 (8%)	4 (2%)	8 28

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	219/221 (99%)	197 (90%)	17 (8%)	5 (2%)	6	21
1	E	218/221 (99%)	189 (87%)	24 (11%)	5 (2%)	6	21
1	F	219/221 (99%)	191 (87%)	25 (11%)	3 (1%)	11	34
1	G	219/221 (99%)	199 (91%)	17 (8%)	3 (1%)	11	34
1	H	219/221 (99%)	196 (90%)	19 (9%)	4 (2%)	8	28
1	I	219/221 (99%)	193 (88%)	22 (10%)	4 (2%)	8	28
1	J	219/221 (99%)	195 (89%)	21 (10%)	3 (1%)	11	34
1	K	219/221 (99%)	200 (91%)	17 (8%)	2 (1%)	17	46
1	L	219/221 (99%)	199 (91%)	18 (8%)	2 (1%)	17	46
1	M	219/221 (99%)	197 (90%)	19 (9%)	3 (1%)	11	34
1	N	219/221 (99%)	197 (90%)	16 (7%)	6 (3%)	5	17
1	O	219/221 (99%)	200 (91%)	16 (7%)	3 (1%)	11	34
All	All	3284/3315 (99%)	2942 (90%)	284 (9%)	58 (2%)	8	28

5 of 58 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	218	ARG
1	A	220	HIS
1	C	3	LEU
1	D	26	PRO
1	B	27	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/194 (100%)	180 (93%)	14 (7%)	14	38
1	B	194/194 (100%)	190 (98%)	4 (2%)	53	84
1	C	194/194 (100%)	189 (97%)	5 (3%)	46	79
1	D	194/194 (100%)	187 (96%)	7 (4%)	35	69

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	193/194 (100%)	187 (97%)	6 (3%)	40	74
1	F	194/194 (100%)	187 (96%)	7 (4%)	35	69
1	G	194/194 (100%)	189 (97%)	5 (3%)	46	79
1	H	194/194 (100%)	192 (99%)	2 (1%)	76	93
1	I	194/194 (100%)	187 (96%)	7 (4%)	35	69
1	J	194/194 (100%)	187 (96%)	7 (4%)	35	69
1	K	194/194 (100%)	186 (96%)	8 (4%)	30	64
1	L	194/194 (100%)	189 (97%)	5 (3%)	46	79
1	M	194/194 (100%)	185 (95%)	9 (5%)	27	60
1	N	194/194 (100%)	189 (97%)	5 (3%)	46	79
1	O	194/194 (100%)	190 (98%)	4 (2%)	53	84
All	All	2909/2910 (100%)	2814 (97%)	95 (3%)	38	72

5 of 95 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	17	ARG
1	K	218	ARG
1	J	93	MET
1	K	30	MET
1	L	185	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	113	HIS
1	L	113	HIS
1	J	163	GLN
1	K	138	ASN
1	M	113	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GTP	M	1422	-	26,34,34	4.25	13 (50%)	33,54,54	3.21	11 (33%)
2	GTP	G	1416	-	26,34,34	4.16	12 (46%)	33,54,54	3.18	12 (36%)
2	GTP	E	1414	-	26,34,34	4.06	12 (46%)	33,54,54	3.17	12 (36%)
2	GTP	O	1424	-	26,34,34	4.13	12 (46%)	33,54,54	3.19	12 (36%)
2	GTP	K	1425	-	26,34,34	4.01	10 (38%)	33,54,54	3.19	12 (36%)
2	GTP	B	1411	-	26,34,34	4.04	11 (42%)	33,54,54	3.21	12 (36%)
2	GTP	F	1420	-	26,34,34	4.15	11 (42%)	33,54,54	3.16	12 (36%)
2	GTP	C	1412	-	26,34,34	4.19	13 (50%)	33,54,54	3.25	11 (33%)
2	GTP	I	1418	-	26,34,34	4.02	11 (42%)	33,54,54	3.17	12 (36%)
2	GTP	D	1413	-	26,34,34	4.21	12 (46%)	33,54,54	3.07	12 (36%)
2	GTP	H	1417	-	26,34,34	4.04	12 (46%)	33,54,54	3.20	12 (36%)
2	GTP	J	1419	-	26,34,34	4.08	12 (46%)	33,54,54	3.23	11 (33%)
2	GTP	A	1415	-	26,34,34	4.12	12 (46%)	33,54,54	3.17	12 (36%)
2	GTP	N	1423	-	26,34,34	4.17	13 (50%)	33,54,54	3.15	11 (33%)
2	GTP	L	1421	-	26,34,34	4.13	11 (42%)	33,54,54	3.16	12 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GTP	M	1422	-	-	5/18/38/38	0/3/3/3
2	GTP	G	1416	-	-	4/18/38/38	0/3/3/3
2	GTP	E	1414	-	-	5/18/38/38	0/3/3/3
2	GTP	O	1424	-	-	5/18/38/38	0/3/3/3
2	GTP	K	1425	-	-	4/18/38/38	0/3/3/3
2	GTP	B	1411	-	-	5/18/38/38	0/3/3/3
2	GTP	F	1420	-	-	6/18/38/38	0/3/3/3
2	GTP	C	1412	-	-	5/18/38/38	0/3/3/3
2	GTP	I	1418	-	-	5/18/38/38	0/3/3/3
2	GTP	D	1413	-	-	4/18/38/38	0/3/3/3
2	GTP	H	1417	-	-	4/18/38/38	0/3/3/3
2	GTP	J	1419	-	-	4/18/38/38	0/3/3/3
2	GTP	A	1415	-	-	4/18/38/38	0/3/3/3
2	GTP	N	1423	-	-	5/18/38/38	0/3/3/3
2	GTP	L	1421	-	-	5/18/38/38	0/3/3/3

The worst 5 of 177 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	1421	GTP	C6-N1	13.07	1.55	1.33
2	H	1417	GTP	C6-N1	12.81	1.55	1.33
2	O	1424	GTP	C6-N1	12.74	1.55	1.33
2	A	1415	GTP	C6-N1	12.64	1.55	1.33
2	N	1423	GTP	C6-N1	12.62	1.55	1.33

The worst 5 of 176 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1417	GTP	C5-C6-N1	-9.19	110.86	123.43
2	K	1425	GTP	C5-C6-N1	-9.17	110.88	123.43
2	F	1420	GTP	C5-C6-N1	-9.17	110.89	123.43
2	C	1412	GTP	C5-C6-N1	-9.11	110.97	123.43
2	L	1421	GTP	C5-C6-N1	-8.99	111.14	123.43

There are no chirality outliers.

5 of 70 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1415	GTP	C3'-C4'-C5'-O5'
2	B	1411	GTP	O4'-C4'-C5'-O5'
2	B	1411	GTP	C3'-C4'-C5'-O5'
2	C	1412	GTP	O4'-C4'-C5'-O5'
2	C	1412	GTP	C3'-C4'-C5'-O5'


There are no ring outliers.


15 monomers are involved in 57 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	M	1422	GTP	2	0
2	G	1416	GTP	5	0
2	E	1414	GTP	2	0
2	O	1424	GTP	3	0
2	K	1425	GTP	3	0
2	B	1411	GTP	5	0
2	F	1420	GTP	3	0
2	C	1412	GTP	4	0
2	I	1418	GTP	9	0
2	D	1413	GTP	5	0
2	H	1417	GTP	3	0
2	J	1419	GTP	3	0
2	A	1415	GTP	5	0
2	N	1423	GTP	2	0
2	L	1421	GTP	3	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





















## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.