



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 06:19 PM EDT

PDB ID : 5NCS  
Title : Structure of the native serpin-type proteinase inhibitor, miropin.  
Authors : Goulas, T.; Ksiazek, M.; Garcia-Ferrer, I.; Mizgalska, D.; Potempa, J.; Gomis-Ruth, X.  
Deposited on : 2017-03-06  
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	<div> <div>2%</div> <div>85%</div> <div>14%</div> <div>.</div> </div>
1	B	375	<div> <div>4%</div> <div>87%</div> <div>12%</div> <div>.</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5902 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serpin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	374	Total	C	N	O	S	0	1	0
			2947	1878	487	562	20			
1	B	374	Total	C	N	O	S	0	0	0
			2939	1873	484	562	20			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP G8UQY8
A	-4	PRO	-	expression tag	UNP G8UQY8
A	-3	LEU	-	expression tag	UNP G8UQY8
A	-2	GLY	-	expression tag	UNP G8UQY8
A	-1	SER	-	expression tag	UNP G8UQY8
A	174	GLN	ARG	conflict	UNP G8UQY8
B	-5	GLY	-	expression tag	UNP G8UQY8
B	-4	PRO	-	expression tag	UNP G8UQY8
B	-3	LEU	-	expression tag	UNP G8UQY8
B	-2	GLY	-	expression tag	UNP G8UQY8
B	-1	SER	-	expression tag	UNP G8UQY8
B	174	GLN	ARG	conflict	UNP G8UQY8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	12	Total	O	0	0
			12	12		
2	B	4	Total	O	0	0
			4	4		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Serpin



#### • Molecule 1: Serpin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.49Å 78.49Å 351.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.97 – 3.00 46.97 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.97-3.00) 100.0 (46.97-3.00)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.37 (at 3.01Å)	Xtriage
Refinement program	REFMAC, PHENIX, BUSTER-TNT, BUSTER 2.10.2	Depositor
R, $R_{free}$	0.176 , 0.216 0.178 , 0.207	Depositor DCC
$R_{free}$ test set	735 reflections (3.18%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	68.7	Xtriage
Anisotropy	0.465	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 79.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5902	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/3011	0.71	0/4072
1	B	0.46	0/3000	0.70	0/4058
All	All	0.47	0/6011	0.70	0/8130

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2947	0	2950	30	0
1	B	2939	0	2937	24	0
2	A	12	0	0	0	0
2	B	4	0	0	0	0
All	All	5902	0	5887	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:41:ILE:HG23	1:B:396:VAL:HG22	1.68	0.75

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:VAL:HG13	1:A:143:ILE:HD11	1.69	0.75
1:B:138:VAL:HG13	1:B:143:ILE:HD11	1.77	0.67
1:A:56:ARG:HH11	1:A:56:ARG:HG3	1.61	0.64
1:A:-3:LEU:HD21	1:A:40:LYS:HD3	1.82	0.62
1:B:56:ARG:HH11	1:B:56:ARG:HG3	1.66	0.60
1:A:394:THR:OG1	1:A:396:VAL:HG23	2.03	0.58
1:B:305:GLU:HG2	1:B:348:PHE:HD1	1.69	0.57
1:A:183:ILE:O	1:A:307:LYS:HE2	2.04	0.56
1:A:394:THR:O	1:B:372:PRO:HG2	2.06	0.56
1:B:394:THR:OG1	1:B:396:VAL:HG23	2.06	0.55
1:B:183:ILE:O	1:B:307:LYS:HE3	2.08	0.54
1:A:140:GLU:HA	1:A:143:ILE:HD12	1.89	0.54
1:A:370:SER:OG	1:B:-1:SER:HA	2.08	0.54
1:B:131:TRP:CD1	1:B:196:ILE:HD12	2.43	0.53
1:A:131:TRP:CD1	1:A:196:ILE:HD12	2.44	0.53
1:A:305:GLU:HG2	1:A:348:PHE:HD1	1.74	0.53
1:B:244:GLU:H	1:B:244:GLU:CD	2.12	0.53
1:A:74:ALA:O	1:A:77:MET:HB2	2.10	0.52
1:A:-1:SER:HB3	1:B:371:SER:H	1.74	0.52
1:A:41:ILE:HG23	1:A:396:VAL:HG22	1.92	0.51
1:A:132:TYR:CZ	1:A:156:ALA:HB2	2.46	0.51
1:B:81:GLY:HA3	1:B:333:ILE:HG13	1.93	0.51
1:B:74:ALA:O	1:B:77:MET:HB2	2.11	0.50
1:A:52:LEU:HD22	1:A:400:ILE:HG12	1.94	0.50
1:A:126:MET:HG2	1:A:199:VAL:HG13	1.94	0.50
1:B:52:LEU:HD22	1:B:400:ILE:HG12	1.94	0.50
1:A:246:CYS:SG	1:A:274:VAL:HG22	2.53	0.49
1:B:132:TYR:CZ	1:B:156:ALA:HB2	2.48	0.49
1:A:246:CYS:HB2	1:A:248:TYR:CE2	2.49	0.48
1:B:126:MET:HG2	1:B:199:VAL:HG13	1.95	0.48
1:A:291:ARG:NH1	1:B:291:ARG:HB3	2.28	0.48
1:A:56:ARG:HG3	1:A:56:ARG:NH1	2.29	0.47
1:A:295:VAL:HG22	1:A:377:ILE:HB	1.97	0.47
1:B:175:LYS:HD2	1:B:175:LYS:N	2.30	0.46
1:A:138:VAL:HG13	1:A:143:ILE:CD1	2.42	0.46
1:A:252:ASP:OD1	1:A:392:LYS:HE3	2.16	0.45
1:B:252:ASP:OD1	1:B:392:LYS:HE3	2.17	0.45
1:B:246:CYS:SG	1:B:274:VAL:HG22	2.57	0.44
1:B:86:THR:HG23	1:B:323:PRO:HG3	1.98	0.44
1:A:40:LYS:HG2	1:A:107:TYR:CZ	2.52	0.44
1:A:369:THR:HG23	1:A:370:SER:N	2.33	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:TYR:OH	1:A:138:VAL:HG12	2.19	0.42
1:A:161:SER:HA	1:A:162:PRO:HD3	1.96	0.42
1:B:81:GLY:HA2	1:B:332:GLY:HA3	2.02	0.42
1:B:56:ARG:HG3	1:B:56:ARG:NH1	2.33	0.41
1:A:65:PHE:CE2	1:A:349:VAL:HB	2.55	0.41
1:B:65:PHE:CE2	1:B:349:VAL:HB	2.55	0.41
1:A:208:GLN:HG2	1:A:357:GLU:O	2.21	0.41
1:A:127:ALA:HB3	1:A:198:ALA:HB3	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	373/375 (100%)	359 (96%)	14 (4%)	0	100	100
1	B	372/375 (99%)	355 (95%)	16 (4%)	1 (0%)	41	76
All	All	745/750 (99%)	714 (96%)	30 (4%)	1 (0%)	51	85

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	-3	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/323 (100%)	307 (95%)	17 (5%)	23	59
1	B	323/323 (100%)	309 (96%)	14 (4%)	29	66
All	All	647/646 (100%)	616 (95%)	31 (5%)	25	62

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-3	LEU
1	A	42	GLU
1	A	56	ARG
1	A	72	SER
1	A	113	GLU
1	A	136	GLU
1	A	165	LEU
1	A	174	GLN
1	A	182	LYS
1	A	225	THR
1	A	243	ASP
1	A	244	GLU
1	A	307	LYS
1	A	329	ASP
1	A	369	THR
1	A	373	SER
1	A	383	LYS
1	B	39	GLU
1	B	40	LYS
1	B	56	ARG
1	B	72	SER
1	B	113	GLU
1	B	137	LEU
1	B	175	LYS
1	B	225	THR
1	B	244	GLU
1	B	278	LEU
1	B	307	LYS
1	B	329	ASP
1	B	364	VAL
1	B	383	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	174	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	374/375 (99%)	-0.24	7 (1%) 66 37	47, 62, 87, 151	0
1	B	374/375 (99%)	0.03	15 (4%) 38 15	56, 87, 135, 161	0
All	All	748/750 (99%)	-0.10	22 (2%) 51 23	47, 72, 130, 161	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	374	THR	4.2
1	B	337	ALA	4.2
1	B	378	ASN	3.9
1	B	335	ASP	3.9
1	B	326	GLU	3.3
1	B	375	THR	3.2
1	B	-3	LEU	3.2
1	A	373	SER	3.0
1	B	327	THR	3.0
1	B	267	GLU	2.9
1	B	137	LEU	2.8
1	A	374	THR	2.8
1	B	223	ASP	2.8
1	A	136	GLU	2.6
1	B	144	LEU	2.5
1	B	372	PRO	2.4
1	A	135	GLY	2.4
1	A	162	PRO	2.4
1	B	373	SER	2.4
1	A	372	PRO	2.3
1	B	139	LYS	2.1
1	A	85	VAL	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.