



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 28, 2025 – 02:08 PM EDT

PDB ID : 9O46 / pdb\_00009o46  
Title : Crystal structure of the L411W mutant of pregnane X receptor ligand binding domain in complex with SJPYT-331  
Authors : Huber, A.D.; Garcia-Maldonado, E.; Miller, D.J.; Chen, T.  
Deposited on : 2025-04-08  
Resolution : 3.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.46


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2927 (3.04-3.00)
Clashscore	180529	3300 (3.04-3.00)
Ramachandran outliers	177936	3188 (3.04-3.00)
Sidechain outliers	177891	3191 (3.04-3.00)
RSRZ outliers	164620	2939 (3.04-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	355	<div> <div style="display: flex; justify-content: space-between;"> <span>%</span> <span>77%</span> <span>5%</span> <span>19%</span> </div> </div>
1	B	355	<div> <div style="display: flex; justify-content: space-between;"> <span>76%</span> <span>6%</span> <span>17%</span> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4634 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pregnane X receptor ligand binding domain tethered to steroid receptor coactivator-1 peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	288	Total	C	N	O	S	0	0	0
			2254	1447	389	401	17			
1	B	293	Total	C	N	O	S	0	0	0
			2303	1486	387	413	17			

There are 44 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	119	MET	-	initiating methionine	UNP O75469
A	120	LYS	-	expression tag	UNP O75469
A	121	LYS	-	expression tag	UNP O75469
A	122	GLY	-	expression tag	UNP O75469
A	123	HIS	-	expression tag	UNP O75469
A	124	HIS	-	expression tag	UNP O75469
A	125	HIS	-	expression tag	UNP O75469
A	126	HIS	-	expression tag	UNP O75469
A	127	HIS	-	expression tag	UNP O75469
A	128	HIS	-	expression tag	UNP O75469
A	129	GLY	-	expression tag	UNP O75469
A	411	TRP	LEU	engineered mutation	UNP O75469
A	431D	SER	-	linker	UNP O75469
A	431E	GLY	-	linker	UNP O75469
A	431F	GLY	-	linker	UNP O75469
A	431G	SER	-	linker	UNP O75469
A	431H	GLY	-	linker	UNP O75469
A	431I	GLY	-	linker	UNP O75469
A	431J	SER	-	linker	UNP O75469
A	431K	SER	-	linker	UNP O75469
A	431L	HIS	-	linker	UNP O75469
A	431M	SER	-	linker	UNP O75469
B	119	MET	-	initiating methionine	UNP O75469
B	120	LYS	-	expression tag	UNP O75469

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	121	LYS	-	expression tag	UNP O75469
B	122	GLY	-	expression tag	UNP O75469
B	123	HIS	-	expression tag	UNP O75469
B	124	HIS	-	expression tag	UNP O75469
B	125	HIS	-	expression tag	UNP O75469
B	126	HIS	-	expression tag	UNP O75469
B	127	HIS	-	expression tag	UNP O75469
B	128	HIS	-	expression tag	UNP O75469
B	129	GLY	-	expression tag	UNP O75469
B	411	TRP	LEU	engineered mutation	UNP O75469
B	430E	SER	-	linker	UNP O75469
B	430F	GLY	-	linker	UNP O75469
B	430G	GLY	-	linker	UNP O75469
B	430H	SER	-	linker	UNP O75469
B	430I	GLY	-	linker	UNP O75469
B	430J	GLY	-	linker	UNP O75469
B	430K	SER	-	linker	UNP O75469
B	430L	SER	-	linker	UNP O75469
B	430M	HIS	-	linker	UNP O75469
B	430N	SER	-	linker	UNP O75469

- Molecule 2 is methyl 3-{[(1P)-1-(2,5-dimethoxyphenyl)-5-methyl-1H-1,2,3-triazole-4-carbonyl]amino}-4-{[(3S)-hexan-3-yl]oxy}benzoate (CCD ID: WU6) (formula: C<sub>26</sub>H<sub>32</sub>N<sub>4</sub>O<sub>6</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			36	26	4	6		
2	B	1	Total	C	N	O	0	0
			36	26	4	6		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	O	0	0
			2	2		
3	B	3	Total	O	0	0
			3	3		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.69Å 87.04Å 103.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.34 – 3.02 42.34 – 3.02	Depositor EDS
% Data completeness (in resolution range)	99.5 (42.34-3.02) 99.5 (42.34-3.02)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.20	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 3.01Å)	Xtriage
Refinement program	PHENIX 1.21.1_5286	Depositor
R, $R_{free}$	0.197 , 0.232 0.198 , 0.234	Depositor DCC
$R_{free}$ test set	758 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	64.5	Xtriage
Anisotropy	0.300	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 40.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4634	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.69% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: WU6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.22	0/2304	0.40	0/3113
1	B	0.21	0/2356	0.36	0/3187
All	All	0.21	0/4660	0.38	0/6300

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2254	0	2177	7	0
1	B	2303	0	2224	12	0
2	A	36	0	0	0	0
2	B	36	0	0	0	0
3	A	2	0	0	0	0
3	B	3	0	0	0	0
All	All	4634	0	4401	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:273:ILE:HG12	1:B:452:LEU:HD22	1.86	0.57
1:A:349:PHE:O	1:A:366:GLN:HB2	2.08	0.54
1:B:409:GLN:O	1:B:413:ARG:HG3	2.08	0.53
1:B:254:ILE:HD12	1:B:283:LEU:HB3	1.89	0.53
1:B:278:GLY:HA3	1:B:353:ARG:HD2	1.91	0.53
1:B:206:LEU:HD23	1:B:236:ILE:HB	1.91	0.51
1:B:410:ARG:O	1:B:414:ILE:HG13	2.12	0.50
1:A:346:ILE:HD13	1:A:394:MET:HE1	1.95	0.49
1:A:211:VAL:HG12	1:A:308:LEU:HA	1.96	0.47
1:B:339:GLU:OE2	1:B:381:ARG:HD2	2.15	0.47
1:A:258:ALA:HB1	1:A:264:PHE:CD2	2.50	0.46
1:A:377:ILE:HG23	1:A:381:ARG:HD2	1.99	0.45
1:B:279:ALA:HB1	1:B:283:LEU:HD13	1.98	0.44
1:B:454:ARG:O	1:B:458:GLU:HG3	2.17	0.44
1:A:221:SER:HB2	1:B:225:TYR:CE1	2.53	0.44
1:A:373:LEU:HD21	1:A:393:ILE:HG21	1.98	0.44
1:B:173:ARG:NH1	1:B:241:PRO:HB3	2.35	0.41
1:B:330:LEU:HD12	1:B:333:LEU:HD12	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/355 (79%)	276 (98%)	6 (2%)	0	100	100
1	B	287/355 (81%)	277 (96%)	10 (4%)	0	100	100
All	All	569/710 (80%)	553 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/313 (74%)	229 (99%)	3 (1%)	65	84
1	B	239/313 (76%)	239 (100%)	0	100	100
All	All	471/626 (75%)	468 (99%)	3 (1%)	84	92

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	213	LEU
1	A	309	GLU
1	A	418	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	168	HIS
1	A	409	GLN
1	B	289	ASN
1	B	344	GLN
1	B	380	ASN
1	B	457	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	WU6	B	501	-	36,38,38	1.33	4 (11%)	49,52,52	2.38	9 (18%)
2	WU6	A	501	-	36,38,38	1.30	3 (8%)	49,52,52	1.18	5 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	WU6	B	501	-	-	12/28/31/31	0/3/3/3
2	WU6	A	501	-	-	4/28/31/31	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	WU6	C19-N18	3.29	1.45	1.35
2	A	501	WU6	C19-N18	3.28	1.45	1.35
2	B	501	WU6	O14-C12	3.15	1.40	1.33
2	A	501	WU6	O14-C12	3.14	1.40	1.33
2	B	501	WU6	O33-C32	2.77	1.41	1.37
2	B	501	WU6	O14-C15	-2.15	1.40	1.45
2	A	501	WU6	O14-C15	-2.10	1.40	1.45

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	WU6	O33-C32-C25	9.62	128.51	118.45
2	B	501	WU6	C32-C25-N24	6.87	126.34	117.99
2	B	501	WU6	O33-C32-C31	-6.03	114.14	124.30
2	B	501	WU6	C34-O33-C32	5.52	125.60	117.51
2	B	501	WU6	O14-C12-C11	4.91	119.66	112.31
2	A	501	WU6	O14-C12-C11	4.59	119.18	112.31
2	A	501	WU6	C32-C25-N24	2.95	121.58	117.99
2	B	501	WU6	C36-C35-C21	-2.77	126.23	129.67
2	A	501	WU6	C36-C35-C21	-2.57	126.48	129.67
2	B	501	WU6	C16-C17-C08	2.46	121.84	118.95
2	B	501	WU6	C29-O28-C27	-2.37	112.42	117.50
2	B	501	WU6	O14-C12-O13	-2.24	119.09	123.46
2	A	501	WU6	C16-C17-C08	2.19	121.52	118.95
2	A	501	WU6	O20-C19-C21	2.07	124.06	120.75

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	501	WU6	C10-C11-C12-O14
2	B	501	WU6	C16-C11-C12-O14
2	B	501	WU6	O13-C12-O14-C15
2	A	501	WU6	C11-C12-O14-C15
2	B	501	WU6	C11-C12-O14-C15
2	B	501	WU6	C10-C11-C12-O13
2	B	501	WU6	C25-C32-O33-C34
2	B	501	WU6	C16-C11-C12-O13
2	B	501	WU6	C31-C32-O33-C34
2	A	501	WU6	O13-C12-O14-C15
2	B	501	WU6	C30-C27-O28-C29
2	B	501	WU6	C26-C27-O28-C29
2	A	501	WU6	C01-C02-C03-C04
2	B	501	WU6	C01-C02-C03-C04
2	A	501	WU6	O20-C19-C21-C35
2	B	501	WU6	O20-C19-C21-C35


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	288/355 (81%)	-0.22	3 (1%) 79 61	31, 57, 103, 130	0
1	B	293/355 (82%)	-0.22	1 (0%) 90 81	32, 58, 105, 125	0
All	All	581/710 (81%)	-0.22	4 (0%) 84 68	31, 57, 103, 130	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	199	TRP	2.8
1	A	311	THR	2.5
1	A	229	ALA	2.3
1	B	228	PRO	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	WU6	B	501	36/36	0.92	0.12	49,63,73,78	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	WU6	A	501	36/36	0.94	0.10	38,52,66,77	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





**Electron density around WU6 A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.