Full wwPDB X-ray Structure Validation Report (i)

Oct 19, 2023 – 03:21 AM EDT

PDB ID	:	2O50
Title	:	The crystal structure of Toxoplasma gondii Enoyl acyl carrier protein reductase
Authors	:	Muench, S.P.; Prigge, S.T.; McLeod, R.; Rafferty, J.B.; Rice, D.W.
Deposited on	:	2006-12-05
Resolution	:	2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	2% 63% 26% • 8%
1	B	315	3% 67% 23% • 8%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 4299 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enoyl-acyl carrier reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	290	2140	1352	369	415	4	0	0	0
1	B	290	2148	1364	366	414	4	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	168	MSE	MET	modified residue	UNP Q6UCJ9
A	193	MSE	MET	modified residue	UNP Q6UCJ9
A	295	MSE	MET	modified residue	UNP Q6UCJ9
A	304	MSE	MET	modified residue	UNP Q6UCJ9
B	168	MSE	MET	modified residue	UNP Q6UCJ9
B	193	MSE	MET	modified residue	UNP Q6UCJ9
B	295	MSE	MET	modified residue	UNP Q6UCJ9
B	304	MSE	MET	modified residue	UNP Q6UCJ9

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	5	5	5	0	0
2	B	6	6	6	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Enoyl-acyl carrier reductase

Chain A:



- Molecule 1: Enoyl-acyl carrier reductase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α, β, γ	76.00Å 76.00Å 187.47Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.90 29.52 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-2.90) 99.7 (29.52-2.90)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.79 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.209 , 0.277 0.215 , 0.290	Depositor DCC
R_{free} test set	730 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å^2)	43.0	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.31 , 34.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.032 for -h,-k,l	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	4299	wwPDB-VP
Average B, all atoms (Å^2)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.60	1/2183 (0.0%)	0.73	3/2964 (0.1%)
1	B	0.57	0/2192	0.67	0/2978
All	All	0.59	1/4375 (0.0%)	0.70	3/5942 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	86	PRO	C-N	-9.18	1.12	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	86	PRO	O-C-N	6.76	133.52	122.70
1	A	84	ASP	CB-CG-OD2	5.20	122.98	118.30
1	A	241	LYS	CA-C-N	-5.06	106.06	117.20

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	241	LYS	Mainchain
1	A	4	PRO	Peptide
1	A	64	LYS	Peptide
1	A	66	PRO	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2140	0	2067	72	0
1	B	2148	0	2086	60	0
2	A	5	0	0	1	0
2	B	6	0	0	0	0
All	All	4299	0	4153	129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (129) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:244:ILE:HG23	1:B:245:ASP:N	1.70	1.06
1:A:85:LYS:HB3	1:A:87:GLU:OE1	1.57	1.02
1:B:244:ILE:CG2	1:B:245:ASP:N	2.24	1.01
1:B:262:HIS:HD2	1:B:264:ASP:H	1.12	0.96
1:B:6:ASP:OD1	1:B:8:ARG:HD3	1.70	0.91
1:B:244:ILE:CG2	1:B:245:ASP:H	1.83	0.90
1:A:165:GLY:HA2	1:A:168:MSE:HE3	1.52	0.89
1:B:184:ARG:NH2	1:B:293:HIS:O	2.10	0.83
1:A:242:SER:HB2	1:A:244:ILE:HG13	1.59	0.82
1:A:262:HIS:HD2	1:A:264:ASP:H	1.29	0.81
1:A:161:LEU:CD1	1:A:207:LEU:HD13	2.11	0.79
1:B:262:HIS:CD2	1:B:264:ASP:H	1.99	0.79
1:A:161:LEU:HD12	1:A:207:LEU:HD13	1.63	0.79
1:A:5:ILE:HG21	1:A:277:LEU:HG	1.65	0.79
1:B:244:ILE:HG22	1:B:245:ASP:H	1.48	0.78
1:A:53:SER:HB2	1:A:58:ARG:HD3	1.66	0.77
1:A:86:PRO:HD2	1:A:87:GLU:OE1	1.85	0.76
1:B:63:ARG:NH1	1:B:70:LEU:HB3	2.04	0.72
1:A:301:SER:HB3	1:A:304:MSE:HG2	1.72	0.71
1:A:85:LYS:HB3	1:A:86:PRO:HD2	1.71	0.71
1:B:3:PHE:N	1:B:4:PRO:HD2	2.09	0.68
1:A:262:HIS:CD2	1:A:264:ASP:H	2.12	0.67
1:A:85:LYS:CB	1:A:87:GLU:OE1	2.39	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:257:LEU:HD11	1:B:218:ARG:HG2	1.77	0.65
1:A:106:THR:OG1	1:A:109:GLU:HG3	1.98	0.64
1:A:161:LEU:HD23	1:A:168:MSE:HE1	1.80	0.64
1:B:63:ARG:NH1	1:B:71:ILE:O	2.31	0.63
1:B:161:LEU:HD13	1:B:168:MSE:HE1	1.79	0.63
1:A:43:TRP:CE2	1:A:45:PRO:HG3	2.36	0.60
1:A:181:ALA:HA	1:A:184:ARG:O	2.01	0.59
1:A:165:GLY:CA	1:A:168:MSE:HE3	2.31	0.58
1:B:242:SER:OG	1:B:244:ILE:HG22	2.02	0.58
1:B:244:ILE:O	1:B:247:ALA:N	2.36	0.58
1:A:43:TRP:CE2	1:A:45:PRO:CG	2.88	0.57
1:B:44:PRO:HB2	1:B:45:PRO:HD3	1.88	0.56
1:B:183:GLU:OE2	1:B:287:TYR:OH	2.22	0.56
1:B:17:VAL:HG21	1:B:40:LEU:HD22	1.88	0.56
1:A:43:TRP:CD2	1:A:45:PRO:HG2	2.42	0.55
1:A:123:ILE:HG23	1:A:274:LEU:HD22	1.87	0.55
1:A:4:PRO:O	1:A:5:ILE:HG23	2.07	0.55
1:A:51:GLN:HG2	1:A:55:GLN:OE1	2.07	0.54
1:B:7:LEU:HD12	1:B:34:ALA:HB3	1.90	0.54
1:A:64:LYS:C	1:A:65:LEU:O	2.46	0.53
1:B:44:PRO:HB2	1:B:45:PRO:CD	2.39	0.53
1:A:184:ARG:NH2	1:A:293:HIS:O	2.41	0.53
1:A:241:LYS:N	1:A:246:TYR:HH	2.07	0.53
1:B:106:THR:O	1:B:110:VAL:HG23	2.09	0.53
1:A:183:GLU:OE2	1:A:287:TYR:OH	2.27	0.53
1:A:54:LEU:HD23	1:A:59:LEU:HD12	1.90	0.53
1:A:53:SER:O	1:A:58:ARG:HB2	2.09	0.52
1:A:78:PRO:HB2	1:A:105:TYR:HB3	1.90	0.52
1:A:256:PRO:HB3	1:B:209:TRP:HA	1.90	0.52
1:A:104:GLY:HA2	1:A:109:GLU:OE1	2.09	0.52
1:A:63:ARG:O	1:A:70:LEU:HA	2.10	0.51
1:A:154:ALA:O	1:A:155:TYR:C	2.48	0.51
1:A:115:LYS:HD3	1:A:167:ILE:CG2	2.39	0.51
1:A:54:LEU:HD23	1:A:59:LEU:CD1	2.40	0.51
1:B:181:ALA:HA	1:B:184:ARG:O	2.10	0.51
1:A:29:LYS:HA	1:A:71:ILE:HD12	1.92	0.51
1:B:164:PHE:O	1:B:168:MSE:HG3	2.09	0.51
1:B:47:LEU:HD23	1:B:102:VAL:HG11	1.93	0.51
1:A:79:LEU:HG	1:A:110:VAL:HG21	1.94	0.50
1:A:64:LYS:O	1:A:65:LEU:O	2.30	0.50
1:B:301:SER:HB3	1:B:304:MSE:HG2	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:PRO:CD	1:A:87:GLU:OE1	2.59	0.50
1:B:7:LEU:HD23	1:B:10:GLN:NE2	2.27	0.49
1:A:214:LYS:HD3	1:A:215:TYR:CZ	2.47	0.49
1:A:106:THR:HG1	1:A:109:GLU:HG3	1.77	0.49
1:B:23:TYR:CE1	1:B:266:VAL:HG21	2.48	0.49
1:B:242:SER:O	1:B:243:PHE:C	2.51	0.48
1:B:164:PHE:CB	1:B:168:MSE:HE2	2.43	0.48
1:B:244:ILE:O	1:B:245:ASP:C	2.52	0.48
1:A:7:LEU:O	1:A:10:GLN:HB2	2.14	0.48
1:B:63:ARG:NH1	1:B:70:LEU:CB	2.75	0.48
1:B:63:ARG:O	1:B:71:ILE:HG13	2.13	0.48
1:A:28:ALA:HB1	1:A:38:VAL:HG11	1.96	0.47
1:B:141:THR:OG1	1:B:191:GLY:HA2	2.15	0.47
1:A:40:LEU:O	1:A:77:TYR:N	2.43	0.46
1:B:12:ALA:HB2	1:B:123:ILE:HB	1.96	0.46
1:A:92:ASP:O	1:A:96:ASN:HB2	2.15	0.46
1:A:222:ILE:HG23	1:A:288:VAL:HG23	1.98	0.46
1:B:180:LEU:HD12	1:B:180:LEU:HA	1.68	0.46
1:A:43:TRP:CZ2	1:A:45:PRO:HG3	2.51	0.46
1:A:153:SER:HB3	2:A:317:HOH:O	2.15	0.46
1:B:110:VAL:O	1:B:114:VAL:HG23	2.16	0.46
1:A:175:VAL:HA	1:A:220:ASN:O	2.16	0.46
1:A:18:ALA:HB1	1:A:46:VAL:HG21	1.99	0.45
1:A:99:TYR:O	1:A:102:VAL:HG22	2.17	0.45
1:B:44:PRO:HD3	1:B:79:LEU:O	2.16	0.45
1:A:14:VAL:HA	1:A:125:VAL:HB	1.98	0.44
1:B:165:GLY:N	1:B:168:MSE:HE3	2.33	0.44
1:A:209:TRP:CH2	1:B:299:VAL:HG11	2.52	0.44
1:A:55:GLN:HE21	1:A:55:GLN:HB2	1.58	0.44
1:A:95:ASP:O	1:A:96:ASN:C	2.55	0.44
1:A:222:ILE:HG23	1:A:288:VAL:CG2	2.48	0.44
1:B:123:ILE:HG23	1:B:274:LEU:HD22	2.00	0.43
1:B:250:TYR:OH	1:B:298:ALA:HB2	2.18	0.43
1:A:241:LYS:HB3	1:A:245:ASP:OD1	2.18	0.43
1:A:155:TYR:O	1:A:158:VAL:N	2.51	0.43
1:B:218:ARG:NH2	1:B:273:LEU:O	2.42	0.43
1:A:106:THR:O	1:A:107:ILE:C	2.57	0.43
1:A:182:ALA:HB2	1:A:197:LYS:HB2	2.01	0.43
1:A:244:ILE:HG13	1:A:245:ASP:H	1.83	0.43
1:B:3:PHE:N	1:B:4:PRO:CD	2.80	0.43
1:B:47:LEU:O	1:B:51:GLN:HG3	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:85:LYS:O	1:B:88:ASP:HB2	2.18	0.43
1:B:23:TYR:CE1	1:B:227:LEU:HD21	2.54	0.42
1:B:37:ARG:HH12	1:B:118:LEU:HB3	1.83	0.42
1:B:54:LEU:HA	1:B:54:LEU:HD23	1.87	0.42
1:A:94:LYS:HB2	1:A:94:LYS:HE3	1.83	0.42
1:A:115:LYS:HD3	1:A:167:ILE:HG22	2.01	0.42
1:B:278:ALA:HB1	1:B:281:VAL:HB	2.02	0.42
1:A:90:PRO:HG2	1:A:93:ILE:HG13	2.02	0.42
1:B:124:LEU:O	1:B:174:ALA:HA	2.20	0.42
1:B:5:ILE:O	1:B:5:ILE:HG22	2.20	0.41
1:B:78:PRO:HB2	1:B:105:TYR:HB3	2.02	0.41
1:B:63:ARG:HB2	1:B:70:LEU:HD22	2.01	0.41
1:B:138:LEU:HD12	1:B:138:LEU:HA	1.88	0.41
1:A:242:SER:CB	1:A:244:ILE:HG13	2.42	0.41
1:B:81:ALA:HB1	1:B:156:SER:HB2	2.02	0.41
1:B:193:MSE:O	1:B:197:LYS:HG2	2.21	0.41
1:A:43:TRP:CD1	1:A:45:PRO:HD2	2.56	0.41
1:A:44:PRO:N	1:A:45:PRO:HD2	2.35	0.41
1:B:164:PHE:HB3	1:B:168:MSE:HE2	2.03	0.41
1:A:53:SER:CB	1:A:58:ARG:HD3	2.42	0.41
1:B:164:PHE:O	1:B:165:GLY:C	2.59	0.41
1:A:56:SER:OG	1:A:58:ARG:HG3	2.21	0.40
1:B:92:ASP:O	1:B:96:ASN:N	2.54	0.40
1:A:107:ILE:HD13	1:A:159:SER:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	286/315 (91%)	255 (89%)	28 (10%)	3 (1%)	15 45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	286/315 (91%)	255 (89%)	27 (9%)	4 (1%)	11	36
All	All	572/630 (91%)	510 (89%)	55 (10%)	7 (1%)	13	40

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	65	LEU
1	A	66	PRO
1	A	107	ILE
1	B	243	PHE
1	B	66	PRO
1	B	297	GLN
1	B	45	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/238 (91%)	207 (96%)	9 (4%)	30	63
1	B	218/238 (92%)	210 (96%)	8 (4%)	34	68
All	All	434/476 (91%)	417 (96%)	17 (4%)	32	66

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	HIS
1	A	37	ARG
1	A	55	GLN
1	A	153	SER
1	A	195	SER
1	A	242	SER
1	A	244	ILE
1	A	259	ARG
1	A	300	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	3	PHE
1	B	63	ARG
1	B	67	ASP
1	B	85	LYS
1	B	128	LEU
1	B	161	LEU
1	B	178	SER
1	B	302	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	262	HIS
1	A	297	GLN
1	B	10	GLN
1	B	262	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	86:PRO	C	87:GLU	N	1.12

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	286/315 (90%)	-0.11	5 (1%)	70	69	20, 38, 46, 53	0
1	B	286/315 (90%)	-0.14	8 (2%)	53	49	20, 38, 46, 54	0
All	All	572/630 (90%)	-0.12	13 (2%)	60	58	20, 38, 46, 54	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	66	PRO	3.4
1	B	242	SER	2.7
1	A	241	LYS	2.5
1	A	67	ASP	2.4
1	B	61	GLU	2.4
1	A	97	LYS	2.4
1	B	67	ASP	2.4
1	A	242	SER	2.3
1	B	95	ASP	2.3
1	B	68	GLY	2.3
1	A	66	PRO	2.2
1	B	305	PRO	2.0
1	B	59	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.