



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 19, 2023 – 06:49 AM EDT

PDB ID : 2OSL  
Title : Crystal structure of Rituximab Fab in complex with an epitope peptide  
Authors : Du, J.; Zhong, C.; Ding, J.  
Deposited on : 2007-02-06  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	213	 7% 67% 28% 5%
1	L	213	 72% 23% 5%
2	A	224	 7% 67% 29% 5%
2	H	224	 3% 72% 22% 6%
3	P	25	 64% 16% 20%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	Q	25	 60% 28% 12%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called light chain of the Rituximab Fab fragment, light chain of the Rituximab Fab fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	213	Total	C	N	O	S	0	0	0
			1623	1016	273	328	6			
1	B	213	Total	C	N	O	S	0	0	0
			1623	1016	273	328	6			

- Molecule 2 is a protein called heavy chain of the Rituximab Fab fragment, heavy chain of the Rituximab Fab fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	224	Total	C	N	O	S	0	0	0
			1669	1055	274	332	8			
2	A	221	Total	C	N	O	S	0	0	0
			1647	1043	270	327	7			

- Molecule 3 is a protein called B-lymphocyte antigen CD20.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	P	20	Total	C	N	O	S	0	0	0
			153	93	24	34	2			
3	Q	22	Total	C	N	O	S	0	0	0
			173	106	27	38	2			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	L	51	Total	O	0
			51	51	
4	H	61	Total	O	0
			61	61	
4	B	47	Total	O	0
			47	47	

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	45	Total 45	O 45	0	0
4	P	3	Total 3	O 3	0	0
4	Q	5	Total 5	O 5	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: light chain of the Rituximab Fab fragment,light chain of the Rituximab Fab fragment



- Molecule 1: light chain of the Rituximab Fab fragment,light chain of the Rituximab Fab fragment



- Molecule 2: heavy chain of the Rituximab Fab fragment,heavy chain of the Rituximab Fab fragment



- Molecule 2: heavy chain of the Rituximab Fab fragment,heavy chain of the Rituximab Fab fragment





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.42Å 98.81Å 107.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.97 – 2.60 43.97 – 2.59	Depositor EDS
% Data completeness (in resolution range)	90.6 (43.97-2.60) 90.3 (43.97-2.59)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 2.58Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.239 , 0.296 0.238 , 0.254	Depositor DCC
$R_{free}$ test set	1518 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.9	Xtrriage
Anisotropy	0.309	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 34.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.017 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	7100	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.41	0/1662	0.69	1/2260 (0.0%)
1	L	0.45	0/1662	0.72	0/2260
2	A	0.41	0/1691	0.70	1/2307 (0.0%)
2	H	0.45	0/1713	0.77	1/2334 (0.0%)
3	P	0.50	0/157	0.67	0/214
3	Q	0.43	0/178	0.59	0/243
All	All	0.43	0/7063	0.72	3/9618 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	113	SER	N-CA-C	-5.55	96.02	111.00
2	H	104	GLY	N-CA-C	-5.45	99.49	113.10
2	A	103	GLY	N-CA-C	-5.12	100.31	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1623	0	1575	44	0
1	L	1623	0	1574	37	0
2	A	1647	0	1602	44	0
2	H	1669	0	1624	34	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	P	153	0	134	1	0
3	Q	173	0	149	6	0
4	A	45	0	0	8	0
4	B	47	0	0	6	0
4	H	61	0	0	4	0
4	L	51	0	0	9	0
4	P	3	0	0	0	0
4	Q	5	0	0	0	0
All	All	7100	0	6658	159	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 159 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:38:LYS:HE3	4:A:254:HOH:O	1.64	0.97
1:L:20:THR:HG22	1:L:73:THR:HB	1.52	0.89
2:A:100:THR:HG23	2:A:101:TYR:H	1.35	0.89
2:H:35:HIS:HD2	2:H:47:TRP:HE1	1.17	0.88
1:L:80:GLU:HG2	4:L:261:HOH:O	1.74	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	211/213 (99%)	199 (94%)	11 (5%)	1 (0%)	29 52
1	L	211/213 (99%)	201 (95%)	7 (3%)	3 (1%)	11 22
2	A	219/224 (98%)	202 (92%)	14 (6%)	3 (1%)	11 22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	222/224 (99%)	205 (92%)	13 (6%)	4 (2%)	8	16
3	P	18/25 (72%)	16 (89%)	1 (6%)	1 (6%)	2	2
3	Q	20/25 (80%)	17 (85%)	3 (15%)	0	100	100
All	All	901/924 (98%)	840 (93%)	49 (5%)	12 (1%)	12	24

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	103	GLY
1	B	129	ALA
2	A	62	GLN
1	L	83	ALA
2	H	137	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	185/185 (100%)	160 (86%)	25 (14%)	4	6
1	L	185/185 (100%)	165 (89%)	20 (11%)	6	12
2	A	183/186 (98%)	156 (85%)	27 (15%)	3	5
2	H	186/186 (100%)	159 (86%)	27 (14%)	3	5
3	P	19/24 (79%)	17 (90%)	2 (10%)	7	13
3	Q	21/24 (88%)	20 (95%)	1 (5%)	25	49
All	All	779/790 (99%)	677 (87%)	102 (13%)	4	7

5 of 102 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	91	THR
1	B	209	ASN
3	P	168	GLU
1	B	104	GLU

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	134	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	197	HIS
2	A	207	ASN
3	Q	181	GLN
2	A	205	ASN
2	H	207	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	213/213 (100%)	-0.13	3 (1%) 75 71	24, 44, 78, 121	0
1	L	213/213 (100%)	-0.16	0 100 100	24, 40, 64, 90	0
2	A	221/224 (98%)	0.37	16 (7%) 15 11	30, 57, 97, 156	0
2	H	224/224 (100%)	-0.08	6 (2%) 54 48	23, 41, 73, 127	0
3	P	20/25 (80%)	0.05	0 100 100	36, 53, 83, 100	0
3	Q	22/25 (88%)	-0.21	0 100 100	34, 49, 62, 66	0
All	All	913/924 (98%)	-0.00	25 (2%) 54 48	23, 45, 82, 156	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	140	SER	8.7
2	A	102	TYR	7.1
2	A	141	GLY	5.9
2	H	103	GLY	4.7
2	A	104	GLY	4.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.