# Full wwPDB X-ray Structure Validation Report (i)

Dec 16, 2023 – 11:15 AM EST

| | | |
|---|---|---|
| PDB ID | : | 4P3G |
| Title | : | Structure of the SRP68-RBD from Chaetomium thermophilum |
| Authors | : | Grotwinkel, J.T.; Wild, K.; Sinning, I. |
| Deposited on | : | 2014-03-07 |
| Resolution | : | 2.70 Å(reported) |

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

| | | |
|---|---|---|
| MolProbity | : | 4.02b-467 |
| Mogul | : | 1.8.5 (274361), CSD as541be (2020) |
| Xtriage (Phenix) | : | 1.13 |
| EDS | : | 2.36 |
| Percentile statistics | : | 20191225.v01 (using entries in the PDB archive December 25th 2019) |
| Refmac | : | 5.8.0158 |
| CCP4 | : | 7.0.044 (Gargrove) |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.36 |

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|---|---|---|
| $R_{free}$ | 130704 | 2808 (2.70-2.70) |
| Clashscore | 141614 | 3122 (2.70-2.70) |
| Ramachandran outliers | 138981 | 3069 (2.70-2.70) |
| Sidechain outliers | 138945 | 3069 (2.70-2.70) |
| RSRZ outliers | 127900 | 2737 (2.70-2.70) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|---|---|---|---|
| 1 | A | 224 | 7%, 78%, 20%, • |
| 1 | B | 224 | 12%, 76%, 14%, 9% |
| 1 | C | 224 | 2%, 71%, 11%, 18% |
| 1 | D | 224 | 10%, 71%, 15%, •, 13% |

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|---|---|---|---|---|---|---|---|
| 2 | PO4 | B | 301 | - | - | X | - |
| 2 | PO4 | C | 301 | - | - | X | - |
| 2 | PO4 | D | 301 | - | - | X | - |

# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 6247 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Signal recognition particle subunit SRP68.

| Mol | Chain | Residues | Atoms | | | | | | ZeroOcc | AltConf | Trace |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | C | N | O | S | Se | | | |
| 1 | A | 219 | 1701 | 1071 | 315 | 308 | 3 | 4 | 0 | 0 | 0 |
| 1 | B | 203 | 1575 | 993 | 287 | 288 | 3 | 4 | 0 | 0 | 0 |
| 1 | C | 184 | 1429 | 900 | 260 | 262 | 3 | 4 | 0 | 0 | 0 |
| 1 | D | 195 | 1505 | 949 | 273 | 276 | 3 | 4 | 0 | 0 | 0 |

There are 32 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|---|---|---|---|---|---|
| A | -6 | MSE | - | initiating methionine | UNP G0S5V2 |
| A | -5 | SER | - | expression tag | UNP G0S5V2 |
| A | -4 | HIS | - | expression tag | UNP G0S5V2 |
| A | -3 | HIS | - | expression tag | UNP G0S5V2 |
| A | -2 | HIS | - | expression tag | UNP G0S5V2 |
| A | -1 | HIS | - | expression tag | UNP G0S5V2 |
| A | 0 | HIS | - | expression tag | UNP G0S5V2 |
| A | 1 | HIS | - | expression tag | UNP G0S5V2 |
| B | -6 | MSE | - | initiating methionine | UNP G0S5V2 |
| B | -5 | SER | - | expression tag | UNP G0S5V2 |
| B | -4 | HIS | - | expression tag | UNP G0S5V2 |
| B | -3 | HIS | - | expression tag | UNP G0S5V2 |
| B | -2 | HIS | - | expression tag | UNP G0S5V2 |
| B | -1 | HIS | - | expression tag | UNP G0S5V2 |
| B | 0 | HIS | - | expression tag | UNP G0S5V2 |
| B | 1 | HIS | - | expression tag | UNP G0S5V2 |
| C | -6 | MSE | - | initiating methionine | UNP G0S5V2 |
| C | -5 | SER | - | expression tag | UNP G0S5V2 |
| C | -4 | HIS | - | expression tag | UNP G0S5V2 |
| C | -3 | HIS | - | expression tag | UNP G0S5V2 |
| C | -2 | HIS | - | expression tag | UNP G0S5V2 |

*Continued on next page...*

*Continued from previous page...*

| Chain | Residue | Modelled | Actual | Comment | Reference |
|---|---|---|---|---|---|
| C | -1 | HIS | - | expression tag | UNP G0S5V2 |
| C | 0 | HIS | - | expression tag | UNP G0S5V2 |
| C | 1 | HIS | - | expression tag | UNP G0S5V2 |
| D | -6 | MSE | - | initiating methionine | UNP G0S5V2 |
| D | -5 | SER | - | expression tag | UNP G0S5V2 |
| D | -4 | HIS | - | expression tag | UNP G0S5V2 |
| D | -3 | HIS | - | expression tag | UNP G0S5V2 |
| D | -2 | HIS | - | expression tag | UNP G0S5V2 |
| D | -1 | HIS | - | expression tag | UNP G0S5V2 |
| D | 0 | HIS | - | expression tag | UNP G0S5V2 |
| D | 1 | HIS | - | expression tag | UNP G0S5V2 |

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: $O_4P$).




| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
|---|---|---|---|---|---|---|---|
| | | | Total | O | P | | |
| 2 | A | 1 | 5 | 4 | 1 | 0 | 0 |
| 2 | B | 1 | 5 | 4 | 1 | 0 | 0 |
| 2 | B | 1 | 5 | 4 | 1 | 0 | 0 |
| 2 | C | 1 | 5 | 4 | 1 | 0 | 0 |
| 2 | D | 1 | 5 | 4 | 1 | 0 | 0 |

- Molecule 3 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|---|---|---|---|---|---|---|
| | | | Total | O | | |
| 3 | A | 3 | 3 | 3 | 0 | 0 |
| | | | Total | O | | |
| 3 | B | 3 | 3 | 3 | 0 | 0 |
| | | | Total | O | | |
| 3 | C | 4 | 4 | 4 | 0 | 0 |
| | | | Total | O | | |
| 3 | D | 2 | 2 | 2 | 0 | 0 |

# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Signal recognition particle subunit SRP68

Chain A:




• Molecule 1: Signal recognition particle subunit SRP68

Chain B:




• Molecule 1: Signal recognition particle subunit SRP68

Chain C:




• Molecule 1: Signal recognition particle subunit SRP68

Chain D:


MSE SER HIS HIS HIS HO I3 V7 V8 E12 L16 Y17 Y23 L27 S28 L31 L32 K36 K37 L38 N39 I40 ALA THR LYS ASN ARG GLY LYS PHE HIS PRO LYS THR ALA I54 T55 P56 R67 L68 Q69 L70 A76 A86 N90 THR LYS
G93 R97 H101 R105 R111 D122 G123 R143 K150 Q151 N152 W153 C156 S159 R164 A169 L170 A171 T172 S173 S174 K175 G176 D177 L183 T186 I187 D188 P189 R192 Y196 K199 I200 PRO R202 T203 L204 P205 A210 HIS ARG ALA PHE
GLU GLN SER

# 4 Data and refinement statistics (i)

| Property | Value | Source |
|---|---|---|
| Space group | P 21 21 21 | Depositor |
| Cell constants<br>a, b, c, $\alpha$, $\beta$, $\gamma$ | 46.84Å 95.05Å 223.86Å<br>90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 42.02 – 2.70<br>58.69 – 2.70 | Depositor<br>EDS |
| % Data completeness<br>(in resolution range) | 99.5 (42.02-2.70)<br>99.4 (58.69-2.70) | Depositor<br>EDS |
| $R_{merge}$ | (Not available) | Depositor |
| $R_{sym}$ | (Not available) | Depositor |
| $< I/\sigma(I) >$ [1] | 2.08 (at 2.69Å) | Xtriage |
| Refinement program | PHENIX (phenix.refine: 1.8.1_1168) | Depositor |
| R, $R_{free}$ | 0.200 , 0.253<br>0.203 , 0.256 | Depositor<br>DCC |
| $R_{free}$ test set | 1441 reflections (5.10%) | wwPDB-VP |
| Wilson B-factor (Å$^2$) | 63.8 | Xtriage |
| Anisotropy | 0.518 | Xtriage |
| Bulk solvent $k_{sol}$(e/Å$^3$), $B_{sol}$(Å$^2$) | 0.33 , 77.7 | EDS |
| L-test for twinning[2] | $< \|L\| > = 0.49$, $< L^2 > = 0.32$ | Xtriage |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| $F_o$,$F_c$ correlation | 0.94 | EDS |
| Total number of atoms | 6247 | wwPDB-VP |
| Average B, all atoms (Å$^2$) | 108.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|---|---|---|---|---|---|
| | | RMSZ | $\#\|Z\|>5$ | RMSZ | $\#\|Z\|>5$ |
| 1 | A | 0.50 | 1/1728 (0.1%) | 0.68 | 2/2324 (0.1%) |
| 1 | B | 0.44 | 0/1596 | 0.62 | 0/2146 |
| 1 | C | 0.47 | 0/1448 | 0.61 | 0/1944 |
| 1 | D | 0.44 | 0/1522 | 0.61 | 0/2043 |
| All | All | 0.46 | 1/6294 (0.0%) | 0.63 | 2/8457 (0.0%) |

All (1) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|---|---|---|---|---|---|---|---|
| 1 | A | 136 | ARG | CG-CD | -5.23 | 1.38 | 1.51 |

All (2) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^o$) | Ideal($^o$) |
|---|---|---|---|---|---|---|---|
| 1 | A | 136 | ARG | NE-CZ-NH1 | -5.45 | 117.57 | 120.30 |
| 1 | A | 136 | ARG | CD-NE-CZ | -5.43 | 116.00 | 123.60 |

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|---|---|---|---|---|---|---|
| 1 | A | 1701 | 0 | 1746 | 27 | 1 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|---|---|---|---|---|---|---|
| 1 | B | 1575 | 0 | 1613 | 17 | 1 |
| 1 | C | 1429 | 0 | 1459 | 19 | 0 |
| 1 | D | 1505 | 0 | 1548 | 22 | 0 |
| 2 | A | 5 | 0 | 0 | 1 | 0 |
| 2 | B | 10 | 0 | 0 | 4 | 0 |
| 2 | C | 5 | 0 | 0 | 2 | 0 |
| 2 | D | 5 | 0 | 0 | 3 | 0 |
| 3 | A | 3 | 0 | 0 | 0 | 0 |
| 3 | B | 3 | 0 | 0 | 0 | 0 |
| 3 | C | 4 | 0 | 0 | 1 | 0 |
| 3 | D | 2 | 0 | 0 | 0 | 0 |
| All | All | 6247 | 0 | 6366 | 80 | 1 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|---|---|---|---|
| 1:D:111:ARG:NH1 | 2:D:301:PO4:O4 | 2.07 | 0.88 |
| 1:D:164:ARG:NH1 | 1:D:188:ASP:OD1 | 2.17 | 0.77 |
| 1:D:143:ARG:NH2 | 2:D:301:PO4:O3 | 2.18 | 0.76 |
| 1:A:206:ILE:HD12 | 1:B:164:ARG:HD3 | 1.68 | 0.76 |
| 1:C:164:ARG:NH1 | 1:C:188:ASP:OD1 | 2.20 | 0.74 |
| 1:A:140:ALA:HB3 | 1:A:163:CYS:HB3 | 1.75 | 0.69 |
| 1:C:5:LYS:HG2 | 1:C:178:ILE:CD1 | 2.29 | 0.63 |
| 1:A:202:ARG:HG2 | 2:B:301:PO4:O2 | 1.98 | 0.62 |
| 1:A:51:LYS:HG2 | 1:A:52:THR:HG23 | 1.79 | 0.62 |
| 1:C:5:LYS:HG3 | 3:C:403:HOH:O | 2.00 | 0.62 |
| 1:A:83:MSE:HE2 | 1:A:98:THR:HG23 | 1.84 | 0.60 |
| 1:D:171:ALA:O | 1:D:175:LYS:N | 2.33 | 0.60 |
| 1:C:143:ARG:NH2 | 2:C:301:PO4:O1 | 2.35 | 0.59 |
| 1:A:202:ARG:O | 1:B:192:ARG:NH2 | 2.36 | 0.57 |
| 1:D:189:PRO:HA | 1:D:192:ARG:HD2 | 1.86 | 0.57 |
| 1:D:204:LEU:HG | 1:D:205:PRO:HD2 | 1.86 | 0.57 |
| 1:D:170:LEU:O | 1:D:173:SER:OG | 2.19 | 0.57 |
| 1:D:8:VAL:O | 1:D:12:GLU:HG3 | 2.04 | 0.57 |
| 1:A:4:THR:HG21 | 1:A:167:TYR:OH | 2.03 | 0.56 |
| 1:D:23:TYR:CZ | 1:D:27:LEU:HD11 | 2.41 | 0.56 |
| 1:D:32:LEU:HG | 1:D:36:LYS:HE3 | 1.87 | 0.56 |
| 1:A:16:LEU:HD22 | 1:A:17:TYR:CE2 | 2.41 | 0.56 |

*Continued on next page...*

*Continued from previous page...*

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|---|---|---|---|
| 1:C:164:ARG:O | 1:C:168:THR:HG22 | 2.06 | 0.55 |
| 1:A:174:SER:C | 1:A:176:GLY:H | 2.09 | 0.55 |
| 1:C:5:LYS:HG2 | 1:C:178:ILE:HD12 | 1.88 | 0.55 |
| 1:D:54:ILE:HD12 | 1:D:69:GLN:HG3 | 1.89 | 0.55 |
| 1:A:171:ALA:O | 1:A:175:LYS:N | 2.39 | 0.54 |
| 1:A:171:ALA:O | 1:A:176:GLY:N | 2.40 | 0.54 |
| 1:A:121:SER:HA | 1:A:132:ILE:HD13 | 1.92 | 0.52 |
| 1:C:111:ARG:HG3 | 2:C:301:PO4:O1 | 2.10 | 0.52 |
| 1:B:212:ARG:NE | 2:B:301:PO4:O3 | 2.39 | 0.52 |
| 1:B:16:LEU:HB3 | 1:B:17:TYR:CD2 | 2.45 | 0.51 |
| 1:B:212:ARG:HE | 2:B:301:PO4:P | 2.33 | 0.51 |
| 1:D:76:ALA:HA | 1:D:105:ARG:HH21 | 1.75 | 0.51 |
| 1:D:56:PRO:HG2 | 1:D:123:GLY:HA3 | 1.93 | 0.50 |
| 1:A:72:THR:HG22 | 1:A:112:ILE:HD13 | 1.92 | 0.50 |
| 1:A:54:ILE:HD12 | 1:A:69:GLN:HG3 | 1.93 | 0.50 |
| 1:B:187:ILE:O | 1:B:191:MSE:HG3 | 2.10 | 0.50 |
| 1:B:108:LYS:HE3 | 1:B:112:ILE:HD11 | 1.94 | 0.50 |
| 1:A:23:TYR:CZ | 1:A:27:LEU:HD11 | 2.48 | 0.49 |
| 1:B:151:GLN:HG2 | 1:B:153:TRP:CZ2 | 2.49 | 0.48 |
| 1:A:23:TYR:CE2 | 1:A:27:LEU:HD11 | 2.49 | 0.48 |
| 1:D:151:GLN:HG2 | 1:D:153:TRP:CZ2 | 2.48 | 0.48 |
| 1:D:67:ARG:HH11 | 1:D:70:LEU:HD12 | 1.79 | 0.47 |
| 1:B:83:MSE:SE | 1:B:105:ARG:HE | 2.46 | 0.47 |
| 1:C:79:HIS:HD2 | 1:C:105:ARG:CZ | 2.28 | 0.47 |
| 1:B:208:THR:HG23 | 1:C:158:LYS:HE2 | 1.97 | 0.47 |
| 1:B:3:ILE:HG23 | 1:B:70:LEU:HD21 | 1.97 | 0.47 |
| 1:A:107:GLU:HB3 | 2:A:301:PO4:O1 | 2.15 | 0.47 |
| 1:C:114:GLU:HG3 | 1:C:139:ALA:HB1 | 1.97 | 0.47 |
| 1:C:140:ALA:HB3 | 1:C:163:CYS:HB3 | 1.96 | 0.46 |
| 1:D:196:TYR:HA | 1:D:199:LYS:HE3 | 1.98 | 0.46 |
| 1:D:186:THR:C | 1:D:189:PRO:HD2 | 2.36 | 0.46 |
| 1:D:67:ARG:HD3 | 1:D:67:ARG:HA | 1.74 | 0.46 |
| 1:B:204:LEU:HD12 | 1:B:205:PRO:HD2 | 1.97 | 0.46 |
| 1:A:45:ARG:HD3 | 1:A:45:ARG:HA | 1.64 | 0.45 |
| 1:C:8:VAL:O | 1:C:12:GLU:HG3 | 2.17 | 0.45 |
| 1:C:79:HIS:CD2 | 1:C:105:ARG:NE | 2.84 | 0.45 |
| 1:B:111:ARG:HD3 | 2:B:302:PO4:O2 | 2.17 | 0.44 |
| 1:D:31:LEU:HD12 | 1:D:31:LEU:HA | 1.87 | 0.44 |
| 1:A:121:SER:OG | 1:A:122:ASP:N | 2.50 | 0.44 |
| 1:A:174:SER:C | 1:A:176:GLY:N | 2.71 | 0.44 |
| 1:B:140:ALA:HB3 | 1:B:163:CYS:HB3 | 1.99 | 0.43 |

*Continued on next page...*

*Continued from previous page...*

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|---|---|---|---|
| 1:A:4:THR:HG23 | 1:A:134:ASP:OD1 | 2.18 | 0.43 |
| 1:C:15:LEU:HD21 | 1:C:20:TYR:CE1 | 2.53 | 0.43 |
| 1:C:72:THR:HG22 | 1:C:112:ILE:HD13 | 2.00 | 0.43 |
| 1:A:38:LEU:HD21 | 1:A:64:GLU:HB2 | 2.01 | 0.43 |
| 1:C:150:LYS:HE2 | 2:D:301:PO4:O1 | 2.19 | 0.43 |
| 1:C:107:GLU:OE1 | 1:D:150:LYS:NZ | 2.52 | 0.42 |
| 1:B:136:ARG:NH1 | 1:D:169:ALA:HA | 2.35 | 0.42 |
| 1:B:178:ILE:HG13 | 1:B:179:PHE:N | 2.33 | 0.42 |
| 1:C:199:LYS:HD3 | 1:C:199:LYS:HA | 1.83 | 0.42 |
| 1:A:120:LEU:HD22 | 1:A:127:ALA:HB2 | 2.03 | 0.41 |
| 1:A:7:VAL:HG21 | 1:A:138:TYR:CE1 | 2.55 | 0.41 |
| 1:B:2:ASP:OD1 | 1:B:5:LYS:HD2 | 2.19 | 0.41 |
| 1:A:120:LEU:HD12 | 1:A:135:ALA:HB2 | 2.03 | 0.41 |
| 1:A:198:ALA:O | 1:A:202:ARG:NH1 | 2.53 | 0.41 |
| 1:C:2:ASP:CG | 1:C:5:LYS:HD2 | 2.42 | 0.40 |
| 1:D:199:LYS:HE3 | 1:D:199:LYS:HB2 | 1.80 | 0.40 |
| 1:A:175:LYS:HB3 | 1:A:175:LYS:HE3 | 1.81 | 0.40 |

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|---|---|---|---|
| 1:A:108:LYS:NZ | 1:B:19:ASP:OD2[3_554] | 2.10 | 0.10 |

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles |
|---|---|---|---|---|---|---|
| 1 | A | 217/224 (97%) | 203 (94%) | 12 (6%) | 2 (1%) | 17 40 |
| 1 | B | 197/224 (88%) | 192 (98%) | 5 (2%) | 0 | 100 100 |
| 1 | C | 178/224 (80%) | 172 (97%) | 6 (3%) | 0 | 100 100 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|---|---|---|---|---|---|---|---|
| 1 | D | 187/224 (84%) | 180 (96%) | 7 (4%) | 0 | 100 | 100 |
| All | All | 779/896 (87%) | 747 (96%) | 30 (4%) | 2 (0%) | 41 | 66 |

All (2) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | A | 90 | ASN |
| 1 | A | 44 | ASN |

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|---|---|---|---|---|---|---|
| 1 | A | 173/173 (100%) | 167 (96%) | 6 (4%) | 36 | 65 |
| 1 | B | 160/173 (92%) | 154 (96%) | 6 (4%) | 33 | 62 |
| 1 | C | 145/173 (84%) | 142 (98%) | 3 (2%) | 53 | 80 |
| 1 | D | 153/173 (88%) | 149 (97%) | 4 (3%) | 46 | 75 |
| All | All | 631/692 (91%) | 612 (97%) | 19 (3%) | 41 | 70 |

All (19) residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | A | 43 | LYS |
| 1 | A | 48 | PHE |
| 1 | A | 49 | HIS |
| 1 | A | 105 | ARG |
| 1 | A | 122 | ASP |
| 1 | A | 156 | CYS |
| 1 | B | 1 | HIS |
| 1 | B | 16 | LEU |
| 1 | B | 91 | THR |
| 1 | B | 92 | LYS |
| 1 | B | 128 | SER |
| 1 | B | 156 | CYS |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | C | 79 | HIS |
| 1 | C | 122 | ASP |
| 1 | C | 156 | CYS |
| 1 | D | 67 | ARG |
| 1 | D | 156 | CYS |
| 1 | D | 159 | SER |
| 1 | D | 175 | LYS |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | C | 79 | HIS |

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Counts | RMSZ | $\#\|Z\| > 2$ | Counts | RMSZ | $\#\|Z\| > 2$ |
| 2 | PO4 | B | 301 | - | 4,4,4 | 1.03 | 0 | 6,6,6 | 0.69 | 0 |

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Counts | RMSZ | $\#\|Z\| > 2$ | Counts | RMSZ | $\#\|Z\| > 2$ |
| 2 | PO4 | D | 301 | - | 4,4,4 | 1.14 | 0 | 6,6,6 | 0.52 | 0 |
| 2 | PO4 | A | 301 | - | 4,4,4 | 0.82 | 0 | 6,6,6 | 0.42 | 0 |
| 2 | PO4 | C | 301 | - | 4,4,4 | 0.86 | 0 | 6,6,6 | 0.77 | 0 |
| 2 | PO4 | B | 302 | - | 4,4,4 | 1.00 | 0 | 6,6,6 | 0.54 | 0 |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 10 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|---|---|---|---|---|---|
| 2 | B | 301 | PO4 | 3 | 0 |
| 2 | D | 301 | PO4 | 3 | 0 |
| 2 | A | 301 | PO4 | 1 | 0 |
| 2 | C | 301 | PO4 | 2 | 0 |
| 2 | B | 302 | PO4 | 1 | 0 |

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, $95^{th}$ percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å$^2$) | Q<0.9 |
|---|---|---|---|---|---|---|
| 1 | A | 215/224 (95%) | 0.57 | 15 (6%) 16 14 | 59, 89, 167, 217 | 0 |
| 1 | B | 199/224 (88%) | 0.76 | 27 (13%) 3 2 | 65, 110, 183, 218 | 0 |
| 1 | C | 180/224 (80%) | 0.33 | 4 (2%) 62 63 | 56, 89, 145, 182 | 0 |
| 1 | D | 191/224 (85%) | 0.73 | 22 (11%) 4 4 | 65, 112, 181, 218 | 0 |
| All | All | 785/896 (87%) | 0.60 | 68 (8%) 10 8 | 56, 99, 175, 218 | 0 |

All (68) RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|---|---|---|---|---|
| 1 | D | 17 | TYR | 6.5 |
| 1 | D | 172 | THR | 5.9 |
| 1 | A | 45 | ARG | 5.7 |
| 1 | B | 66 | VAL | 5.7 |
| 1 | D | 176 | GLY | 5.2 |
| 1 | A | 49 | HIS | 4.9 |
| 1 | D | 40 | ILE | 4.8 |
| 1 | D | 122 | ASP | 4.6 |
| 1 | D | 38 | LEU | 4.5 |
| 1 | D | 174 | SER | 3.6 |
| 1 | B | 38 | LEU | 3.3 |
| 1 | A | 89 | ALA | 3.3 |
| 1 | A | 92 | LYS | 3.3 |
| 1 | C | 179 | PHE | 3.2 |
| 1 | A | 173 | SER | 3.1 |
| 1 | D | 7 | VAL | 3.1 |
| 1 | B | 31 | LEU | 3.1 |
| 1 | D | 177 | ASP | 3.1 |
| 1 | B | 215 | GLU | 3.1 |
| 1 | B | 27 | LEU | 3.0 |
| 1 | B | 61 | GLU | 3.0 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type | RSRZ |
|---|---|---|---|---|
| 1 | A | 76 | ALA | 3.0 |
| 1 | D | 123 | GLY | 3.0 |
| 1 | B | 20 | TYR | 3.0 |
| 1 | D | 97 | ARG | 3.0 |
| 1 | C | 175 | LYS | 2.9 |
| 1 | D | 173 | SER | 2.9 |
| 1 | B | 54 | ILE | 2.8 |
| 1 | B | 23 | TYR | 2.8 |
| 1 | B | 16 | LEU | 2.8 |
| 1 | B | 70 | LEU | 2.7 |
| 1 | C | 15 | LEU | 2.7 |
| 1 | D | 16 | LEU | 2.7 |
| 1 | B | 65 | TYR | 2.6 |
| 1 | B | 132 | ILE | 2.6 |
| 1 | B | 34 | CYS | 2.5 |
| 1 | A | 91 | THR | 2.5 |
| 1 | A | 59 | ILE | 2.5 |
| 1 | B | 25 | THR | 2.5 |
| 1 | B | 133 | LEU | 2.5 |
| 1 | D | 101 | HIS | 2.5 |
| 1 | D | 183 | LEU | 2.5 |
| 1 | A | 93 | GLY | 2.5 |
| 1 | B | 73 | ALA | 2.4 |
| 1 | A | 71 | LEU | 2.4 |
| 1 | A | 47 | LYS | 2.4 |
| 1 | D | 86 | ALA | 2.4 |
| 1 | D | 27 | LEU | 2.3 |
| 1 | B | 59 | ILE | 2.3 |
| 1 | A | 87 | HIS | 2.3 |
| 1 | D | 32 | LEU | 2.3 |
| 1 | D | 202 | ARG | 2.3 |
| 1 | B | 106 | LEU | 2.2 |
| 1 | A | 27 | LEU | 2.2 |
| 1 | B | 91 | THR | 2.2 |
| 1 | D | 105 | ARG | 2.2 |
| 1 | B | 26 | GLN | 2.1 |
| 1 | D | 28 | SER | 2.1 |
| 1 | D | 3 | ILE | 2.1 |
| 1 | A | 98 | THR | 2.1 |
| 1 | C | 177 | ASP | 2.1 |
| 1 | B | 120 | LEU | 2.1 |
| 1 | B | 167 | TYR | 2.1 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type | RSRZ |
|---|---|---|---|---|
| 1 | B | 53 | ALA | 2.0 |
| 1 | B | 121 | SER | 2.0 |
| 1 | B | 173 | SER | 2.0 |
| 1 | A | 16 | LEU | 2.0 |
| 1 | B | 64 | GLU | 2.0 |

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, $95^{th}$ percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors(Å$^2$) | Q<0.9 |
|---|---|---|---|---|---|---|---|---|
| 2 | PO4 | A | 301 | 5/5 | 0.82 | 0.30 | 137,142,144,146 | 0 |
| 2 | PO4 | B | 302 | 5/5 | 0.87 | 0.43 | 142,143,146,146 | 0 |
| 2 | PO4 | D | 301 | 5/5 | 0.89 | 0.51 | 110,111,113,119 | 0 |
| 2 | PO4 | C | 301 | 5/5 | 0.90 | 0.56 | 116,118,123,124 | 0 |
| 2 | PO4 | B | 301 | 5/5 | 0.94 | 0.47 | 125,128,133,133 | 0 |

## 6.5 Other polymers (i)

There are no such residues in this entry.