wwPDB X-ray Structure Validation Summary Report (i)

Oct 4, 2023 – 09:37 AM EDT

PDB ID	:	6PPS
Title	:	A blue light illuminated LOV-PAS construct from the LOV-HK sensory protein from Brucella abortus (construct 15-273)
Authors	:	Rinaldi, J.; Fernandez, I.; Shin, H.; Gunawardana, S.; Otero, L.H.; Cerutti, M.L.; Yang, X.; Klinke, S.; Goldbaum, F.A.
Deposited on	:	2019-07-08
Resolution	:	2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	82% 9% • 9%
1	B	279	2% 80% 12% 8%
1	C	279	15% 73% 18% • 8%
1	D	279	85% 7% • 7%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 8410 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Blue-light-activated histidine kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	255	2037	1293	357	379	8	0	0	0
1	B	257	2052	1301	360	383	8	0	0	0
1	C	256	2047	1299	360	380	8	0	0	0
1	D	259	2077	1317	369	383	8	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	274	HIS	-	expression tag	UNP Q2YKK7
A	275	HIS	-	expression tag	UNP Q2YKK7
A	276	HIS	-	expression tag	UNP Q2YKK7
A	277	HIS	-	expression tag	UNP Q2YKK7
A	278	HIS	-	expression tag	UNP Q2YKK7
A	279	HIS	-	expression tag	UNP Q2YKK7
B	274	HIS	-	expression tag	UNP Q2YKK7
B	275	HIS	-	expression tag	UNP Q2YKK7
B	276	HIS	-	expression tag	UNP Q2YKK7
B	277	HIS	-	expression tag	UNP Q2YKK7
B	278	HIS	-	expression tag	UNP Q2YKK7
B	279	HIS	-	expression tag	UNP Q2YKK7
C	274	HIS	-	expression tag	UNP Q2YKK7
C	275	HIS	-	expression tag	UNP Q2YKK7
C	276	HIS	-	expression tag	UNP Q2YKK7
C	277	HIS	-	expression tag	UNP Q2YKK7
C	278	HIS	-	expression tag	UNP Q2YKK7
C	279	HIS	-	expression tag	UNP Q2YKK7
D	274	HIS	-	expression tag	UNP Q2YKK7
D	275	HIS	-	expression tag	UNP Q2YKK7
D	276	HIS	-	expression tag	UNP Q2YKK7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	277	HIS	-	expression tag	UNP Q2YKK7
D	278	HIS	-	expression tag	UNP Q2YKK7
D	279	HIS	-	expression tag	UNP Q2YKK7

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: $C_{17}H_{21}N_4O_9P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	C	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	D	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	20	Total	O	0	0
			20	20		
3	B	15	Total	O	0	0
			15	15		
3	C	16	Total	O	0	0
			16	16		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	D	22	22	22	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Blue-light-activated histidine kinase

Chain A:



• Molecule 1: Blue-light-activated histidine kinase

Chain B:



• Molecule 1: Blue-light-activated histidine kinase

Chain C:



• Molecule 1: Blue-light-activated histidine kinase

Chain D:


MET
ALA
ILE
ASP
LEU
ARG
PRO
PHE
ILE
PRO
PHE
GLY
ARG
GLY
ALA
LEU
SER
GLN
A19
T37
D96
H114
F128
R141
E144
R158
V167
E168
V169
I172
F179
C186
E189
C190
R191
R192
I193
H224
V227
R246
A254
E258
V262
H277
HIS
HIS

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	110.19Å 56.89Å 116.05Å 90.00° 103.16° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 44.26 – 2.79	Depositor EDS
% Data completeness (in resolution range)	99.3 (50.00-2.80) 91.1 (44.26-2.79)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.04 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.209 , 0.254 0.212 , 0.254	Depositor DCC
R_{free} test set	2005 reflections (5.71%)	wwPDB-VP
Wilson B-factor (Å^2)	64.6	Xtriage
Anisotropy	0.649	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.26 , 46.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	8410	wwPDB-VP
Average B, all atoms (Å^2)	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.19 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.8992e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.24	0/2081	0.39	0/2825
1	B	0.22	0/2096	0.39	0/2845
1	C	0.28	0/2092	0.44	0/2840
1	D	0.24	0/2125	0.40	0/2885
All	All	0.25	0/8394	0.41	0/11395

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2037	0	2033	18	0
1	B	2052	0	2046	20	0
1	C	2047	0	2040	36	0
1	D	2077	0	2061	12	0
2	A	31	0	19	1	0
2	B	31	0	19	3	0
2	C	31	0	19	1	0
2	D	31	0	19	0	0
3	A	20	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	15	0	0	0	0
3	C	16	0	0	0	0
3	D	22	0	0	0	0
All	All	8410	0	8256	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 80 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:269:VAL:HG23	1:C:271:GLU:H	1.50	0.77
1:B:259:ASN:O	1:B:259:ASN:ND2	2.17	0.77
1:A:240:THR:HG23	1:A:242:LEU:H	1.56	0.71
1:A:172:ILE:HD11	1:B:172:ILE:HD11	1.74	0.68
1:D:96:ASP:OD1	1:D:114:HIS:ND1	2.29	0.66

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/279 (91%)	246 (97%)	7 (3%)	0	100	100
1	B	255/279 (91%)	245 (96%)	9 (4%)	1 (0%)	34	66
1	C	254/279 (91%)	242 (95%)	11 (4%)	1 (0%)	34	66
1	D	257/279 (92%)	252 (98%)	5 (2%)	0	100	100
All	All	1019/1116 (91%)	985 (97%)	32 (3%)	2 (0%)	47	78

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	270	THR
1	B	270	THR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/242 (92%)	219 (99%)	3 (1%)	67	90
1	B	224/242 (93%)	218 (97%)	6 (3%)	44	78
1	C	223/242 (92%)	215 (96%)	8 (4%)	35	69
1	D	226/242 (93%)	223 (99%)	3 (1%)	69	91
All	All	895/968 (92%)	875 (98%)	20 (2%)	52	83

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	258	GLU
1	D	172	ILE
1	D	224	HIS
1	D	189	GLU
1	B	258	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	274	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	FMN	D	300	-	33,33,33	1.08	2 (6%)	48,50,50	1.21	8 (16%)
2	FMN	B	300	-	33,33,33	1.08	2 (6%)	48,50,50	1.20	8 (16%)
2	FMN	C	300	-	33,33,33	1.07	2 (6%)	48,50,50	1.19	9 (18%)
2	FMN	A	300	-	33,33,33	1.06	2 (6%)	48,50,50	1.19	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	D	300	-	-	1/18/18/18	0/3/3/3
2	FMN	B	300	-	-	3/18/18/18	0/3/3/3
2	FMN	C	300	-	-	0/18/18/18	0/3/3/3
2	FMN	A	300	-	-	0/18/18/18	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	300	FMN	C4A-N5	3.94	1.38	1.30
2	C	300	FMN	C4A-N5	3.84	1.38	1.30
2	D	300	FMN	C4A-N5	3.83	1.38	1.30
2	A	300	FMN	C4A-N5	3.80	1.38	1.30
2	D	300	FMN	C10-N1	2.58	1.38	1.33

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	C	300	FMN	C4-N3-C2	-3.10	119.92	125.64
2	D	300	FMN	C4-N3-C2	-3.07	119.96	125.64
2	A	300	FMN	C4-N3-C2	-3.02	120.06	125.64
2	B	300	FMN	C4-N3-C2	-2.96	120.18	125.64
2	D	300	FMN	C4A-C4-N3	2.60	119.80	113.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	300	FMN	C3'-C4'-C5'-O5'
2	D	300	FMN	C4'-C5'-O5'-P
2	B	300	FMN	C5'-O5'-P-O2P
2	B	300	FMN	O4'-C4'-C5'-O5'

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	300	FMN	3	0
2	C	300	FMN	1	0
2	A	300	FMN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FMN D 300



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN B 300



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN C 300



Bond lengths



Bond angles



Torsions



Rings

Ligand FMN A 300



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	255/279 (91%)	-0.17	0 100 100	54, 79, 124, 193	0
1	B	257/279 (92%)	0.06	6 (2%) 60 51	56, 95, 152, 190	0
1	C	256/279 (91%)	0.78	42 (16%) 1 1	68, 127, 196, 216	0
1	D	259/279 (92%)	-0.08	1 (0%) 92 91	46, 77, 134, 195	0
All	All	1027/1116 (92%)	0.15	49 (4%) 30 21	46, 93, 174, 216	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	245	THR	8.9
1	C	236	TYR	6.4
1	C	177	ARG	5.9
1	C	212	VAL	5.6
1	C	274	HIS	5.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	FMN	C	300	31/31	0.94	0.21	84,107,126,130	0
2	FMN	B	300	31/31	0.96	0.17	58,91,122,135	0
2	FMN	A	300	31/31	0.96	0.22	50,73,106,130	0
2	FMN	D	300	31/31	0.97	0.22	46,68,88,102	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FMN C 300:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around FMN B 300:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FMN A 300:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FMN D 300:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.