Full wwPDB X-ray Structure Validation Report (i)

Feb 15, 2024 – 04:43 PM EST

PDB ID	:	3Q8S
Title	:	Human DNA polymerase iota incorporating dTTP opposite 8-oxo-guanine
Authors	:	Kirouac, K.N.; Ling, H.
Deposited on	:	2011-01-06
Resolution	:	2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	420	11%; 60%; 26%; • ; 10%
2	T	11	27%; 18%; 27%; 27%
3	P	7	29%; 57%; 14%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 3581 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase iota.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	380	2962	1872	516	552	22	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(*TP*CP*AP*(8OG)P*GP*GP*GP*TP*CP*CP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	T	8	190	88	35	58	9	0	1	0

- Molecule 3 is a DNA chain called DNA (5'-D(P*AP*GP*GP*AP*CP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	P	7	142	67	29	39	7	0	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	B	2	2	2	0	0

- Molecule 5 is THYMIDINE-5'-TRIPHOSPHATE (three-letter code: TTP) (formula: $C_{10}H_{17}N_2O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	B	1	29	10	2	14	3	0	1

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	B	231	231	231	0	0
			Total	O		
6	T	13	13	13	0	0
			Total	O		
6	P	12	12	12	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA polymerase iota

Chain B: 11% 60% 26% • 10%



• Molecule 2: DNA (5'-D(*TP*CP*AP*(8OG)P*GP*GP*GP*TP*CP*CP*T)-3')

Chain T: 27% 18% 27% 27%



• Molecule 3: DNA (5'-D(P*AP*GP*GP*AP*CP*CP*C)-3')

Chain P: 29% 57% 14%



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α, β, γ	97.49Å 97.49Å 201.90Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.48 – 2.09 39.48 – 2.09	Depositor EDS
% Data completeness (in resolution range)	94.5 (39.48-2.09) 93.2 (39.48-2.09)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.15 (at 2.08Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.227 , 0.266 0.223 , 0.260	Depositor DCC
R_{free} test set	1374 reflections (4.00%)	wwPDB-VP
Wilson B-factor (Å^2)	34.8	Xtriage
Anisotropy	0.538	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 54.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	3581	wwPDB-VP
Average B, all atoms (Å^2)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 8OG, TTP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	B	0.31	0/3005	0.51	1/4059 (0.0%)
2	T	0.80	0/160	1.59	3/245 (1.2%)
3	P	0.67	0/159	1.26	1/242 (0.4%)
All	All	0.37	0/3324	0.67	5/4546 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	T	843	DG	O4'-C4'-C3'	-7.74	101.36	106.00
1	B	36	ASP	CB-CA-C	-6.06	98.28	110.40
2	T	847	DT	O4'-C1'-N1	-5.94	103.84	108.00
3	P	872	DC	O4'-C4'-C3'	-5.75	102.20	104.50
2	T	844	DT	O4'-C1'-N1	5.64	111.95	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2962	0	3006	115	0
2	T	190	0	100	2	1
3	P	142	0	76	3	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	2	0	0	0	0
5	B	29	0	13	0	0
6	B	231	0	0	13	0
6	P	12	0	0	0	0
6	T	13	0	0	0	0
All	All	3581	0	3195	119	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (119) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:41:GLN:HE22	1:B:194:THR:H	1.09	0.98
1:B:149:SER:N	1:B:150:ALA:HB3	1.81	0.94
1:B:36:ASP:OD2	1:B:194:THR:HG23	1.69	0.92
1:B:149:SER:HB2	1:B:150:ALA:HB2	1.54	0.90
1:B:290:ASN:HB2	6:B:519:HOH:O	1.74	0.88
1:B:164:ASN:H	1:B:170:HIS:HD2	1.22	0.84
1:B:236:ILE:HD11	1:B:261:LEU:HD22	1.60	0.83
1:B:153:VAL:HG22	1:B:174:LEU:HD22	1.62	0.80
1:B:184:ARG:HE	1:B:218:GLN:HE21	1.30	0.79
1:B:90:GLN:NE2	1:B:90:GLN:H	1.79	0.79
1:B:149:SER:H	1:B:150:ALA:HB3	1.48	0.78
1:B:90:GLN:H	1:B:90:GLN:HE21	1.28	0.77
1:B:358:GLU:HB2	1:B:394:MET:HE1	1.68	0.76
1:B:372:LEU:H	1:B:372:LEU:HD12	1.50	0.76
1:B:342:VAL:HG23	1:B:364:ILE:HG12	1.72	0.72
1:B:347:ARG:HE	1:B:404:THR:CG2	2.02	0.71
1:B:371:LYS:HG2	1:B:372:LEU:N	2.07	0.70
1:B:207:LYS:NZ	3:P:873:DC:OP1	2.24	0.70
1:B:127:GLU:OE1	1:B:207:LYS:HE3	1.93	0.68
1:B:59:GLN:OE1	1:B:64:VAL:HG11	1.93	0.68
1:B:131:ASP:OD2	1:B:133:THR:HG23	1.93	0.67
1:B:41:GLN:HE22	1:B:194:THR:N	1.89	0.66
1:B:365:PRO:HB2	1:B:368:VAL:HG23	1.78	0.66
1:B:153:VAL:HG21	1:B:163:ILE:HD11	1.79	0.65
1:B:371:LYS:HG2	1:B:372:LEU:H	1.62	0.65
1:B:359:SER:O	1:B:360:ARG:HD3	1.97	0.65
1:B:348:ARG:HD2	1:B:358:GLU:OE1	1.96	0.65
1:B:170:HIS:HE1	1:B:224:GLU:OE2	1.80	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:236:ILE:HD13	1:B:250:LEU:HD13	1.79	0.63
1:B:110:THR:HG21	1:B:122:ARG:HH11	1.63	0.63
1:B:275:ILE:HG12	6:B:582:HOH:O	1.98	0.63
1:B:347:ARG:HE	1:B:404:THR:HG22	1.62	0.63
1:B:156:HIS:HB2	1:B:219:THR:HB	1.81	0.62
1:B:237:LYS:HE3	1:B:247:ALA:HB3	1.81	0.62
1:B:382:PRO:HG3	6:B:586:HOH:O	1.99	0.62
1:B:304:GLU:HG3	1:B:328:LEU:HG	1.82	0.62
1:B:158:TYR:OH	1:B:228:HIS:HD2	1.83	0.61
1:B:184:ARG:HE	1:B:218:GLN:NE2	1.97	0.61
1:B:112:LEU:HD23	1:B:112:LEU:C	2.19	0.61
1:B:149:SER:N	1:B:150:ALA:CB	2.60	0.61
1:B:110:THR:CG2	1:B:122:ARG:HH11	2.15	0.59
1:B:47:ASN:HB3	1:B:50:LEU:HD12	1.83	0.59
1:B:347:ARG:HH11	1:B:404:THR:HG23	1.68	0.58
2:T:840[B]:8OG:H5'	2:T:840[B]:8OG:O8	2.02	0.58
1:B:131:ASP:OD2	1:B:133:THR:CG2	2.52	0.57
1:B:110:THR:HG21	1:B:122:ARG:HG2	1.87	0.56
1:B:397:VAL:HB	6:B:504:HOH:O	2.05	0.56
1:B:149:SER:O	1:B:165:LEU:HD13	2.04	0.56
1:B:381:THR:HG22	6:B:486:HOH:O	2.04	0.56
1:B:343:ARG:NH2	1:B:361:GLN:HE22	2.03	0.56
1:B:345:ILE:HB	1:B:407:SER:HB3	1.88	0.56
1:B:35:LEU:HD11	1:B:187:MET:HE1	1.89	0.55
1:B:261:LEU:HD23	1:B:285:SER:HB3	1.88	0.55
1:B:342:VAL:CG2	1:B:364:ILE:HG12	2.37	0.55
1:B:47:ASN:HD22	1:B:47:ASN:C	2.11	0.55
1:B:74:GLY:HA3	1:B:87:LYS:HD3	1.88	0.54
1:B:170:HIS:CE1	1:B:224:GLU:OE2	2.61	0.54
1:B:184:ARG:NE	1:B:218:GLN:HE21	2.02	0.54
1:B:189:ASN:HB3	6:B:612:HOH:O	2.07	0.54
1:B:67:ASN:HD22	1:B:67:ASN:C	2.12	0.54
1:B:358:GLU:CB	1:B:394:MET:HE1	2.38	0.53
1:B:35:LEU:HD21	1:B:109:VAL:HG21	1.91	0.53
1:B:189:ASN:HB3	6:B:646:HOH:O	2.10	0.52
1:B:237:LYS:HE2	1:B:244:TYR:HA	1.91	0.52
1:B:272:GLU:O	1:B:273:LEU:HD23	2.09	0.52
1:B:342:VAL:HG21	1:B:383:MET:HE1	1.92	0.52
1:B:341:THR:HG23	1:B:412:ASN:HB3	1.92	0.51
1:B:152:THR:HG22	6:B:456:HOH:O	2.10	0.51
1:B:341:THR:HG23	1:B:412:ASN:CB	2.41	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:164:ASN:H	1:B:170:HIS:CD2	2.14	0.51
1:B:149:SER:CA	1:B:150:ALA:CB	2.88	0.50
1:B:112:LEU:HD23	1:B:112:LEU:O	2.13	0.49
1:B:358:GLU:HB2	1:B:394:MET:CE	2.40	0.49
1:B:105:MET:HE2	1:B:193:LEU:HD11	1.95	0.48
1:B:149:SER:H	1:B:150:ALA:CB	2.23	0.48
1:B:100:THR:O	1:B:104:GLU:HG3	2.14	0.47
1:B:310:LYS:HB2	1:B:402:HIS:CE1	2.49	0.47
1:B:103:ARG:NH2	6:B:623:HOH:O	2.46	0.47
1:B:40:ALA:O	1:B:44:MET:HG3	2.15	0.47
1:B:35:LEU:HD23	1:B:38:PHE:CD1	2.49	0.47
1:B:170:HIS:O	1:B:174:LEU:HG	2.15	0.46
3:P:870:DA:H2'	3:P:871:DC:C6	2.49	0.46
1:B:47:ASN:ND2	1:B:49:GLU:CG	2.79	0.46
1:B:309:LYS:O	1:B:402:HIS:HE1	1.97	0.46
2:T:844:DT:H2'	2:T:845:DC:C6	2.50	0.46
1:B:47:ASN:ND2	1:B:49:GLU:HG2	2.30	0.46
1:B:43:GLU:HG3	1:B:94:VAL:HG21	1.97	0.46
3:P:871:DC:H2'	3:P:872:DC:C6	2.50	0.46
1:B:371:LYS:HD2	1:B:371:LYS:H	1.81	0.46
1:B:149:SER:CB	1:B:150:ALA:HB2	2.34	0.46
1:B:153:VAL:HG22	1:B:174:LEU:CD2	2.39	0.45
1:B:381:THR:N	1:B:382:PRO:HD2	2.32	0.45
1:B:384:VAL:O	1:B:388:MET:HG2	2.18	0.44
1:B:47:ASN:HD21	1:B:49:GLU:CG	2.30	0.44
1:B:275:ILE:C	1:B:275:ILE:HD13	2.38	0.44
1:B:68:TYR:O	1:B:72:LYS:HD3	2.18	0.43
1:B:371:LYS:HD2	1:B:371:LYS:N	2.33	0.43
1:B:292:PRO:HG2	1:B:294:ILE:HD11	2.00	0.43
1:B:45:ILE:O	1:B:48:PRO:HD3	2.18	0.43
1:B:340:HIS:HB3	6:B:617:HOH:O	2.19	0.43
1:B:41:GLN:HA	1:B:44:MET:HE2	2.01	0.43
1:B:52:ASP:HA	6:B:536:HOH:O	2.19	0.42
1:B:48:PRO:O	1:B:51:LYS:HG2	2.19	0.42
1:B:340:HIS:CE1	1:B:414:LYS:HD2	2.55	0.42
1:B:41:GLN:O	1:B:45:ILE:HG13	2.18	0.41
1:B:112:LEU:C	1:B:112:LEU:CD2	2.89	0.41
1:B:55:LEU:H	1:B:67:ASN:ND2	2.19	0.41
1:B:339:PRO:HA	1:B:413:LEU:HD12	2.01	0.41
1:B:322:GLU:HG2	1:B:384:VAL:HG11	2.03	0.41
1:B:129:PHE:CD2	1:B:203:LYS:HG2	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:THR:CG2	6:B:600:HOH:O	2.69	0.41
1:B:203:LYS:HE2	6:B:631:HOH:O	2.21	0.41
1:B:343:ARG:HB3	1:B:409:CYS:HB3	2.01	0.41
1:B:405:LEU:HD23	1:B:405:LEU:C	2.41	0.41
1:B:105:MET:CE	1:B:193:LEU:HD11	2.51	0.41
1:B:320:LYS:O	1:B:323:GLU:HB3	2.21	0.41
1:B:330:ASN:O	1:B:333:CYS:HB3	2.21	0.41
1:B:139:ARG:O	1:B:143:LEU:HD13	2.22	0.40
1:B:161:GLN:NE2	1:B:224:GLU:HG2	2.36	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:T:847:DT:O3'	3:P:867:DA:P[12_544]	2.04	0.16

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	374/420 (89%)	361 (96%)	12 (3%)	1 (0%)	41 41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	150	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	335/376 (89%)	308 (92%)	27 (8%)	11 8

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	47	ASN
1	B	67	ASN
1	B	69	GLU
1	B	90	GLN
1	B	110	THR
1	B	126	ASP
1	B	133	THR
1	B	134	GLU
1	B	166	LEU
1	B	170	HIS
1	B	194	THR
1	B	204	LEU
1	B	219	THR
1	B	221	LEU
1	B	227	GLN
1	B	275	ILE
1	B	279	GLN
1	B	280	ARG
1	B	305	GLU
1	B	308	PHE
1	B	341	THR
1	B	360	ARG
1	B	371	LYS
1	B	381	THR
1	B	396	ASN
1	B	404	THR
1	B	413	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	41	GLN
1	B	47	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	58	GLN
1	B	67	ASN
1	B	90	GLN
1	B	95	ASN
1	B	156	HIS
1	B	170	HIS
1	B	217	GLN
1	B	218	GLN
1	B	228	HIS
1	B	330	ASN
1	B	334	GLN
1	B	340	HIS
1	B	361	GLN
1	B	396	ASN
1	B	402	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	8OG	T	840[B]	2	22,25,26	2.26	7 (31%)	30,37,40	3.18	9 (30%)
2	8OG	T	840[A]	2	22,25,26	2.24	7 (31%)	30,37,40	3.36	10 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8OG	T	840[B]	2	-	1/7/21/22	0/3/3/3
2	8OG	T	840[A]	2	-	1/7/21/22	0/3/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	T	840[B]	8OG	C5-C4	7.78	1.48	1.37
2	T	840[A]	8OG	C5-C4	7.62	1.48	1.37
2	T	840[B]	8OG	C8-N9	-3.69	1.34	1.40
2	T	840[A]	8OG	C8-N9	-3.57	1.34	1.40
2	T	840[B]	8OG	C4-N9	-2.61	1.34	1.39
2	T	840[A]	8OG	C4-N9	-2.61	1.34	1.39
2	T	840[A]	8OG	C6-N1	-2.52	1.34	1.38
2	T	840[B]	8OG	C5-C6	2.51	1.48	1.42
2	T	840[B]	8OG	C6-N1	-2.50	1.34	1.38
2	T	840[A]	8OG	C8-N7	-2.38	1.33	1.38
2	T	840[A]	8OG	C5-N7	-2.29	1.34	1.37
2	T	840[B]	8OG	C8-N7	-2.07	1.34	1.38
2	T	840[B]	8OG	C5-N7	-2.05	1.34	1.37
2	T	840[A]	8OG	C5-C6	2.02	1.47	1.42

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	T	840[A]	8OG	C5-C4-N3	-9.23	118.27	127.80
2	T	840[B]	8OG	C5-C4-N3	-9.16	118.34	127.80
2	T	840[A]	8OG	N7-C8-N9	8.93	117.18	106.58
2	T	840[B]	8OG	N7-C8-N9	8.62	116.80	106.58
2	T	840[A]	8OG	N9-C4-N3	8.49	135.52	125.81
2	T	840[B]	8OG	N9-C4-N3	7.70	134.61	125.81
2	T	840[B]	8OG	C2-N3-C4	5.00	121.21	112.30
2	T	840[A]	8OG	C2-N3-C4	4.90	121.02	112.30
2	T	840[A]	8OG	C5-N7-C8	-4.28	103.31	109.47
2	T	840[B]	8OG	C5-N7-C8	-3.78	104.02	109.47
2	T	840[A]	8OG	O8-C8-N7	-3.39	120.49	126.64
2	T	840[A]	8OG	O6-C6-C5	-3.28	119.71	127.24
2	T	840[B]	8OG	O8-C8-N7	-3.16	120.91	126.64
2	T	840[A]	8OG	O4'-C1'-N9	2.72	111.03	108.29
2	T	840[B]	8OG	O8-C8-N9	-2.67	122.27	125.99
2	T	840[A]	8OG	O8-C8-N9	-2.65	122.30	125.99
2	T	840[B]	8OG	O6-C6-C5	-2.60	121.28	127.24
2	T	840[B]	8OG	C5-C6-N1	2.02	118.32	112.31
2	T	840[A]	8OG	C5-C6-N1	2.01	118.31	112.31

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	T	840[A]	8OG	O4'-C4'-C5'-O5'
2	T	840[B]	8OG	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	T	840[B]	8OG	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	TTP	B	423[A]	4	26,30,30	1.24	5 (19%)	39,47,47	2.00	9 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TTP	B	423[A]	4	-	4/22/34/34	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	423[A]	TTP	C6-C5	2.73	1.39	1.34
5	B	423[A]	TTP	C4-N3	-2.69	1.33	1.38
5	B	423[A]	TTP	C6-N1	-2.30	1.34	1.38
5	B	423[A]	TTP	C2-N3	-2.06	1.34	1.38
5	B	423[A]	TTP	C4-C5	2.01	1.48	1.44

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	B	423[A]	TTP	C4-N3-C2	-5.31	120.47	127.35
5	B	423[A]	TTP	N3-C2-N1	5.09	121.65	114.89
5	B	423[A]	TTP	O4-C4-C5	-4.46	119.74	124.90
5	B	423[A]	TTP	C5-C4-N3	4.43	119.09	115.31
5	B	423[A]	TTP	C5-C6-N1	-3.93	119.29	123.34
5	B	423[A]	TTP	PB-O3B-PG	-2.70	123.55	132.83
5	B	423[A]	TTP	O2-C2-N1	-2.69	119.21	122.79
5	B	423[A]	TTP	PB-O3A-PA	-2.47	124.34	132.83
5	B	423[A]	TTP	O4'-C1'-N1	2.05	111.53	107.86

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	423[A]	TTP	PB-O3B-PG-O2G
5	B	423[A]	TTP	C5'-O5'-PA-O3A
5	B	423[A]	TTP	C5'-O5'-PA-O1A
5	B	423[A]	TTP	PB-O3B-PG-O1G

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	B	380/420 (90%)	0.59	46 (12%) 4 5	23, 50, 102, 156	0
2	T	7/11 (63%)	0.12	0 100 100	27, 33, 39, 39	0
3	P	7/7 (100%)	-0.08	0 100 100	26, 33, 42, 46	0
All	All	394/438 (89%)	0.57	46 (11%) 4 6	23, 48, 101, 156	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	333	CYS	6.8
1	B	372	LEU	6.3
1	B	350	SER	6.0
1	B	349	TYR	5.9
1	B	399	MET	5.2
1	B	401	PHE	5.2
1	B	400	PRO	5.0
1	B	73	LEU	4.9
1	B	61	TYR	4.9
1	B	377	TYR	4.7
1	B	336	GLY	4.1
1	B	376	ASN	4.1
1	B	370	GLN	3.9
1	B	82	ARG	3.7
1	B	144	GLN	3.7
1	B	367	HIS	3.5
1	B	143	LEU	3.3
1	B	397	VAL	3.2
1	B	310	LYS	3.1
1	B	371	LYS	3.1
1	B	368	VAL	3.1
1	B	403	LEU	3.1
1	B	51	LYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	378	ASP	3.0
1	B	244	TYR	2.9
1	B	395	VAL	2.9
1	B	149	SER	2.9
1	B	50	LEU	2.7
1	B	398	LYS	2.7
1	B	348	ARG	2.7
1	B	52	ASP	2.7
1	B	309	LYS	2.6
1	B	391	PHE	2.6
1	B	141	GLN	2.5
1	B	311	CYS	2.5
1	B	142	GLN	2.5
1	B	404	THR	2.5
1	B	62	LEU	2.4
1	B	75	VAL	2.4
1	B	402	HIS	2.3
1	B	147	GLU	2.3
1	B	334	GLN	2.3
1	B	86	GLU	2.2
1	B	319	ASN	2.2
1	B	312	SER	2.2
1	B	396	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	8OG	T	840[A]	23/24	0.90	0.17	38,43,54,60	23
2	8OG	T	840[B]	23/24	0.90	0.17	36,42,52,54	23

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	MG	B	422	1/1	0.95	0.18	24,24,24,24	0
5	TTP	B	423[A]	29/29	0.96	0.19	25,33,36,41	29
4	MG	B	421	1/1	0.99	0.09	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around TTP B 423 (A):

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.