wwPDB X-ray Structure Validation Summary Report (i)

Jun 22, 2024 – 01:04 PM EDT

PDB ID	:	4R07
Title	:	Crystal structure of human TLR8 in complex with ORN06
Authors	:	Tanji, H.; Ohto, U.; Shibata, T.; Taoka, M.; Yamauchi, Y.; Isobe, T.; Miyake, K.; Shimizu, T.
Deposited on	:	2014-07-30
Resolution	:	2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	811	2%, 79%, 12%, •, 8%
1	B	811	2%, 79%, 10%, •, 9%
1	C	811	7%, 75%, 14%, •, 9%
1	D	811	6%, 75%, 13%, •, 9%
2	E	4	75%, 25%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	3	33% 67%
3	H	3	100%
3	I	3	100%
3	L	3	100%
3	M	3	67% 33%
3	N	3	67% 33%
3	O	3	100%
4	G	2	100%
4	J	2	100%
4	P	2	100%
5	K	4	75% 25%

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 26233 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toll-like receptor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	747	6014	3847	1019	1129	19	0	0	0
1	B	738	5949	3808	1009	1113	19	0	0	0
1	C	738	5946	3806	1007	1114	19	0	0	0
1	D	735	5925	3794	1004	1108	19	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	ARG	-	expression tag	UNP Q9NR97
A	23	SER	-	expression tag	UNP Q9NR97
A	24	PRO	-	expression tag	UNP Q9NR97
A	25	TRP	-	expression tag	UNP Q9NR97
A	828	GLU	-	expression tag	UNP Q9NR97
A	829	PHE	-	expression tag	UNP Q9NR97
A	830	LEU	-	expression tag	UNP Q9NR97
A	831	VAL	-	expression tag	UNP Q9NR97
A	832	PRO	-	expression tag	UNP Q9NR97
A	833	ARG	-	expression tag	UNP Q9NR97
B	22	ARG	-	expression tag	UNP Q9NR97
B	23	SER	-	expression tag	UNP Q9NR97
B	24	PRO	-	expression tag	UNP Q9NR97
B	25	TRP	-	expression tag	UNP Q9NR97
B	828	GLU	-	expression tag	UNP Q9NR97
B	829	PHE	-	expression tag	UNP Q9NR97
B	830	LEU	-	expression tag	UNP Q9NR97
B	831	VAL	-	expression tag	UNP Q9NR97
B	832	PRO	-	expression tag	UNP Q9NR97
B	833	ARG	-	expression tag	UNP Q9NR97
C	22	ARG	-	expression tag	UNP Q9NR97

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	23	SER	-	expression tag	UNP Q9NR97
C	24	PRO	-	expression tag	UNP Q9NR97
C	25	TRP	-	expression tag	UNP Q9NR97
C	828	GLU	-	expression tag	UNP Q9NR97
C	829	PHE	-	expression tag	UNP Q9NR97
C	830	LEU	-	expression tag	UNP Q9NR97
C	831	VAL	-	expression tag	UNP Q9NR97
C	832	PRO	-	expression tag	UNP Q9NR97
C	833	ARG	-	expression tag	UNP Q9NR97
D	22	ARG	-	expression tag	UNP Q9NR97
D	23	SER	-	expression tag	UNP Q9NR97
D	24	PRO	-	expression tag	UNP Q9NR97
D	25	TRP	-	expression tag	UNP Q9NR97
D	828	GLU	-	expression tag	UNP Q9NR97
D	829	PHE	-	expression tag	UNP Q9NR97
D	830	LEU	-	expression tag	UNP Q9NR97
D	831	VAL	-	expression tag	UNP Q9NR97
D	832	PRO	-	expression tag	UNP Q9NR97
D	833	ARG	-	expression tag	UNP Q9NR97

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	4	50	28	2	20	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	F	3	39	22	2	15	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	H	3	39	22	2	15	0	0	0
3	I	3	39	22	2	15	0	0	0
3	L	3	39	22	2	15	0	0	0
3	M	3	39	22	2	15	0	0	0
3	N	3	39	22	2	15	0	0	0
3	O	3	39	22	2	15	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	2	28	16	2	10	0	0	0
4	J	2	28	16	2	10	0	0	0
4	P	2	28	16	2	10	0	0	0

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	K	4	50	28	2	20	0	0	0

- Molecule 6 is URIDINE (three-letter code: URI) (formula: $C_9H_{12}N_2O_6$).

URI



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	A	1	17	9	2	6	0	0
			Total	C	N	O		
6	B	1	17	9	2	6	0	0
			Total	C	N	O		
6	C	1	17	9	2	6	0	0
			Total	C	N	O		
6	D	1	17	9	2	6	0	0

- Molecule 7 is 3'-O-[(R)-{[(2R,3aR,4R,6R,6aR)-6-(2-amino-6-oxo-1,6-dihydro-9H-purin-9-yl)-2-hydroxy-2-oxidotetrahydrofuro[3,4-d][1,3,2]dioxaphosphol-4-yl]methoxy}(hydroxy)phosphoryl]uridine 5'-(dihydrogen phosphate) (three-letter code: UCG) (formula: $C_{19}H_{24}N_7O_{18}P_3$).

UCG



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	P	0	0
			47	19	7	18	3		
7	B	1	Total	C	N	O	P	0	0
			47	19	7	18	3		
7	C	1	Total	C	N	O	P	0	0
			47	19	7	18	3		
7	D	1	Total	C	N	O	P	0	0
			47	19	7	18	3		

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	B	1	Total	C	N	O	0	0
			14	8	1	5		
8	B	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
8	B	1	14	8	1	5	0	0
			Total	C	N	O		
8	B	1	14	8	1	5	0	0
			Total	C	N	O		
8	B	1	14	8	1	5	0	0
			Total	C	N	O		
8	C	1	14	8	1	5	0	0
			Total	C	N	O		
8	C	1	14	8	1	5	0	0
			Total	C	N	O		
8	C	1	14	8	1	5	0	0
			Total	C	N	O		
8	C	1	14	8	1	5	0	0
			Total	C	N	O		
8	D	1	14	8	1	5	0	0
			Total	C	N	O		
8	D	1	14	8	1	5	0	0
			Total	C	N	O		
8	D	1	14	8	1	5	0	0
			Total	C	N	O		
8	D	1	14	8	1	5	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
9	A	469	469	469	0	0
			Total	O		
9	B	374	374	374	0	0
			Total	O		
9	C	314	314	314	0	0
			Total	O		
9	D	277	277	277	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Toll-like receptor 8

Chain A:



- Molecule 1: Toll-like receptor 8

Chain B:





• Molecule 1: Toll-like receptor 8

Chain C:



• Molecule 1: Toll-like receptor 8

Chain D:





• Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H: 

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L: 

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N: 67% 33%

NAG1 NAG2 BMA3

• Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O: 100%

NAG1 NAG2 BMA3

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 100%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J: 100%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain P: 100%

NAG1 NAG2

• Molecule 5: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K: 75% 25%

NAG1 NAG2 BMA3 MAN4

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	86.66Å 141.12Å 169.53Å 90.00° 90.59° 90.00°	Depositor
Resolution (Å)	33.74 – 2.00 33.74 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.5 (33.74-2.00) 97.5 (33.74-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$< I/\sigma(I) >$ [1]	1.86 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.188 , 0.234 0.195 , 0.237	Depositor DCC
R_{free} test set	13522 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å^2)	31.7	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 51.8	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.021 for h,-k,-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	26233	wwPDB-VP
Average B, all atoms (Å^2)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.54% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, URI, MAN, UCG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.60	0/6138	0.69	0/8326
1	B	0.57	0/6072	0.71	3/8234 (0.0%)
1	C	0.54	0/6069	0.66	1/8232 (0.0%)
1	D	0.53	0/6048	0.64	1/8202 (0.0%)
All	All	0.56	0/24327	0.68	5/32994 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	3
1	C	0	4
1	D	0	1
All	All	0	9

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	628	ASP	CB-CG-OD1	6.22	123.89	118.30
1	C	621	ASP	CB-CG-OD1	5.17	122.95	118.30
1	D	472	ARG	NE-CZ-NH1	5.08	122.84	120.30
1	B	805	SER	C-N-CD	-5.03	109.54	120.60
1	B	300	ARG	NE-CZ-NH2	-5.02	117.79	120.30

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	418	SER	Peptide
1	B	734	GLU	Peptide
1	B	85	GLY	Peptide
1	B	99	ASN	Peptide
1	C	45	VAL	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6014	0	5989	68	0
1	B	5949	0	5920	70	1
1	C	5946	0	5916	84	0
1	D	5925	0	5895	75	0
2	E	50	0	43	1	0
3	F	39	0	34	0	0
3	H	39	0	34	0	0
3	I	39	0	34	0	0
3	L	39	0	34	0	0
3	M	39	0	34	0	0
3	N	39	0	34	1	0
3	O	39	0	34	0	0
4	G	28	0	25	0	0
4	J	28	0	25	0	0
4	P	28	0	25	0	0
5	K	50	0	43	1	0
6	A	17	0	12	1	0
6	B	17	0	12	1	0
6	C	17	0	12	1	0
6	D	17	0	12	1	0
7	A	47	0	20	4	0
7	B	47	0	20	6	0
7	C	47	0	20	5	0
7	D	47	0	20	1	0
8	A	70	0	65	0	0
8	B	70	0	65	1	0
8	C	56	0	52	0	0
8	D	56	0	52	2	0
9	A	469	0	0	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	B	374	0	0	11	0
9	C	314	0	0	14	0
9	D	277	0	0	12	0
All	All	26233	0	24481	304	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 304 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:75:ILE:HG22	9:D:1225:HOH:O	1.52	1.09
1:A:33:TYR:O	1:A:60:THR:HG21	1.59	1.02
1:A:33:TYR:O	1:A:60:THR:CG2	2.11	0.96
1:D:77:HIS:N	9:D:1225:HOH:O	2.00	0.92
1:B:708:SER:HB3	1:B:735:VAL:HG22	1.52	0.90

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:280:ARG:NH1	1:B:709:ASP:OD2[1_455]	2.03	0.17

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	739/811 (91%)	700 (95%)	37 (5%)	2 (0%)	41 37
1	B	728/811 (90%)	678 (93%)	47 (6%)	3 (0%)	34 30
1	C	728/811 (90%)	665 (91%)	59 (8%)	4 (0%)	29 23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	725/811 (89%)	668 (92%)	53 (7%)	4 (1%)	25	19
All	All	2920/3244 (90%)	2711 (93%)	196 (7%)	13 (0%)	34	30

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	736	SER
1	C	378	VAL
1	D	243	LYS
1	B	284	GLN
1	C	732	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	693/755 (92%)	658 (95%)	35 (5%)	24	19
1	B	685/755 (91%)	649 (95%)	36 (5%)	22	18
1	C	685/755 (91%)	651 (95%)	34 (5%)	24	20
1	D	682/755 (90%)	646 (95%)	36 (5%)	22	18
All	All	2745/3020 (91%)	2604 (95%)	141 (5%)	24	19

5 of 141 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	239	GLU
1	D	300	ARG
1	D	681	TRP
1	B	248	LEU
1	B	243	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	123	ASN
1	D	686	GLN
1	D	809	GLN
1	B	135	ASN
1	B	84	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

35 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NAG	E	1	1,2	14,14,15	0.84	1 (7%)	17,19,21	1.29	3 (17%)
2	NAG	E	2	2	14,14,15	0.77	1 (7%)	17,19,21	1.47	3 (17%)
2	BMA	E	3	2	11,11,12	0.58	0	15,15,17	3.09	6 (40%)
2	MAN	E	4	2	11,11,12	0.80	0	15,15,17	1.36	4 (26%)
3	NAG	F	1	1,3	14,14,15	1.00	1 (7%)	17,19,21	1.00	1 (5%)
3	NAG	F	2	3	14,14,15	0.83	0	17,19,21	1.63	3 (17%)
3	BMA	F	3	3	11,11,12	0.53	0	15,15,17	0.82	0
4	NAG	G	1	4,1	14,14,15	0.98	1 (7%)	17,19,21	1.20	1 (5%)
4	NAG	G	2	4	14,14,15	0.80	1 (7%)	17,19,21	1.89	3 (17%)
3	NAG	H	1	1,3	14,14,15	0.64	0	17,19,21	1.57	4 (23%)
3	NAG	H	2	3	14,14,15	0.78	0	17,19,21	1.60	4 (23%)
3	BMA	H	3	3	11,11,12	0.69	0	15,15,17	1.50	3 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	NAG	I	1	1,3	14,14,15	0.76	0	17,19,21	1.24	2 (11%)
3	NAG	I	2	3	14,14,15	1.02	1 (7%)	17,19,21	1.43	3 (17%)
3	BMA	I	3	3	11,11,12	0.28	0	15,15,17	0.91	1 (6%)
4	NAG	J	1	4,1	14,14,15	0.83	0	17,19,21	1.34	3 (17%)
4	NAG	J	2	4	14,14,15	0.81	1 (7%)	17,19,21	1.46	3 (17%)
5	NAG	K	1	5,1	14,14,15	1.08	1 (7%)	17,19,21	1.76	4 (23%)
5	NAG	K	2	5	14,14,15	0.78	0	17,19,21	1.18	1 (5%)
5	BMA	K	3	5	11,11,12	0.33	0	15,15,17	1.28	1 (6%)
5	MAN	K	4	5	11,11,12	0.85	0	15,15,17	1.48	2 (13%)
3	NAG	L	1	1,3	14,14,15	0.51	0	17,19,21	1.03	1 (5%)
3	NAG	L	2	3	14,14,15	1.02	1 (7%)	17,19,21	1.20	0
3	BMA	L	3	3	11,11,12	0.45	0	15,15,17	1.12	1 (6%)
3	NAG	M	1	1,3	14,14,15	0.74	0	17,19,21	0.88	0
3	NAG	M	2	3	14,14,15	0.52	0	17,19,21	1.31	3 (17%)
3	BMA	M	3	3	11,11,12	0.56	0	15,15,17	1.11	0
3	NAG	N	1	1,3	14,14,15	0.77	0	17,19,21	2.07	5 (29%)
3	NAG	N	2	3	14,14,15	0.76	0	17,19,21	1.21	1 (5%)
3	BMA	N	3	3	11,11,12	0.61	0	15,15,17	2.30	5 (33%)
3	NAG	O	1	1,3	14,14,15	0.65	0	17,19,21	1.56	4 (23%)
3	NAG	O	2	3	14,14,15	1.11	1 (7%)	17,19,21	1.18	2 (11%)
3	BMA	O	3	3	11,11,12	0.50	0	15,15,17	0.81	1 (6%)
4	NAG	P	1	4,1	14,14,15	1.27	2 (14%)	17,19,21	2.34	5 (29%)
4	NAG	P	2	4	14,14,15	0.70	1 (7%)	17,19,21	1.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
2	BMA	E	3	2	-	2/2/19/22	0/1/1/1
2	MAN	E	4	2	-	0/2/19/22	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1
3	BMA	F	3	3	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	1/6/23/26	0/1/1/1
3	NAG	H	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	H	2	3	-	0/6/23/26	0/1/1/1
3	BMA	H	3	3	-	0/2/19/22	0/1/1/1
3	NAG	I	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	I	2	3	-	0/6/23/26	0/1/1/1
3	BMA	I	3	3	-	0/2/19/22	0/1/1/1
4	NAG	J	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	J	2	4	-	0/6/23/26	0/1/1/1
5	NAG	K	1	5,1	-	2/6/23/26	0/1/1/1
5	NAG	K	2	5	-	0/6/23/26	0/1/1/1
5	BMA	K	3	5	-	0/2/19/22	0/1/1/1
5	MAN	K	4	5	-	1/2/19/22	0/1/1/1
3	NAG	L	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	L	2	3	-	0/6/23/26	0/1/1/1
3	BMA	L	3	3	-	0/2/19/22	0/1/1/1
3	NAG	M	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	M	2	3	-	0/6/23/26	0/1/1/1
3	BMA	M	3	3	-	0/2/19/22	0/1/1/1
3	NAG	N	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	N	2	3	-	0/6/23/26	0/1/1/1
3	BMA	N	3	3	-	1/2/19/22	0/1/1/1
3	NAG	O	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	O	2	3	-	0/6/23/26	0/1/1/1
3	BMA	O	3	3	-	0/2/19/22	0/1/1/1
4	NAG	P	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	P	2	4	-	0/6/23/26	0/1/1/1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	L	2	NAG	O5-C1	-3.01	1.38	1.43
4	P	1	NAG	O5-C1	-2.96	1.39	1.43
2	E	1	NAG	O5-C1	-2.73	1.39	1.43
3	F	1	NAG	O5-C1	-2.58	1.39	1.43
5	K	1	NAG	O5-C1	-2.55	1.39	1.43

The worst 5 of 83 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	E	3	BMA	C1-O5-C5	9.30	124.79	112.19
3	N	3	BMA	C1-O5-C5	5.83	120.09	112.19
4	P	1	NAG	O5-C5-C6	-5.60	98.43	107.20
5	K	1	NAG	O5-C1-C2	-5.17	103.12	111.29
4	P	1	NAG	C1-C2-N2	-4.67	102.51	110.49

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	P	1	NAG	O5-C5-C6-O6
3	H	1	NAG	C4-C5-C6-O6
2	E	3	BMA	C4-C5-C6-O6
3	N	1	NAG	C4-C5-C6-O6
3	N	1	NAG	O5-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1	NAG	1	0
3	N	1	NAG	1	0
5	K	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain E



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain F



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain H



Oligosaccharide Chain I



Oligosaccharide Chain L



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain M



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain N



Oligosaccharide Chain O



Oligosaccharide Chain G



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain J



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain P



Bond lengths



Bond angles



Torsions



Rings

Oligosaccharide Chain K



5.6 Ligand geometry (i)

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
8	NAG	C	916	1	14,14,15	0.73	0	17,19,21	1.91	3 (17%)
8	NAG	B	912	1	14,14,15	0.84	1 (7%)	17,19,21	1.65	4 (23%)
7	UCG	C	902	-	45,52,52	1.69	12 (26%)	60,82,82	1.60	11 (18%)
8	NAG	A	916	1	14,14,15	0.76	1 (7%)	17,19,21	2.10	4 (23%)
8	NAG	C	914	1	14,14,15	0.85	0	17,19,21	1.27	2 (11%)
6	URI	D	901	-	18,18,18	1.20	2 (11%)	26,26,26	1.72	6 (23%)
6	URI	A	901	-	18,18,18	1.39	2 (11%)	26,26,26	1.62	6 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
6	URI	B	901	-	18,18,18	1.42	5 (27%)	26,26,26	2.00	7 (26%)
8	NAG	A	915	1	14,14,15	0.50	0	17,19,21	1.29	3 (17%)
8	NAG	A	913	1	14,14,15	0.86	0	17,19,21	1.28	2 (11%)
7	UCG	D	902	-	45,52,52	1.65	12 (26%)	60,82,82	1.77	16 (26%)
8	NAG	D	909	1	14,14,15	0.64	0	17,19,21	1.58	2 (11%)
7	UCG	B	902	-	45,52,52	1.76	10 (22%)	60,82,82	1.78	14 (23%)
8	NAG	A	914	1	14,14,15	0.56	0	17,19,21	1.44	3 (17%)
8	NAG	B	914	1	14,14,15	0.53	0	17,19,21	1.75	3 (17%)
6	URI	C	901	-	18,18,18	1.25	2 (11%)	26,26,26	2.05	9 (34%)
8	NAG	C	910	1	14,14,15	0.50	0	17,19,21	1.49	2 (11%)
8	NAG	C	915	1	14,14,15	0.63	0	17,19,21	0.99	1 (5%)
7	UCG	A	902	-	45,52,52	1.80	11 (24%)	60,82,82	1.88	16 (26%)
8	NAG	D	913	1	14,14,15	0.60	0	17,19,21	1.84	5 (29%)
8	NAG	D	914	1	14,14,15	0.74	0	17,19,21	1.75	3 (17%)
8	NAG	B	909	1	14,14,15	0.86	1 (7%)	17,19,21	0.96	0
8	NAG	D	912	1	14,14,15	0.96	1 (7%)	17,19,21	1.52	4 (23%)
8	NAG	A	910	1	14,14,15	0.59	0	17,19,21	1.61	3 (17%)
8	NAG	B	913	1	14,14,15	0.68	0	17,19,21	1.50	3 (17%)
8	NAG	B	915	1	14,14,15	0.94	1 (7%)	17,19,21	2.26	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	C	916	1	-	0/6/23/26	0/1/1/1
8	NAG	B	912	1	-	0/6/23/26	0/1/1/1
7	UCG	C	902	-	-	3/21/67/67	0/6/6/6
8	NAG	A	916	1	-	2/6/23/26	0/1/1/1
8	NAG	C	914	1	-	0/6/23/26	0/1/1/1
6	URI	D	901	-	-	4/6/22/22	0/2/2/2
6	URI	A	901	-	-	4/6/22/22	0/2/2/2
6	URI	B	901	-	-	4/6/22/22	0/2/2/2
8	NAG	A	915	1	-	2/6/23/26	0/1/1/1
8	NAG	A	913	1	-	0/6/23/26	0/1/1/1
7	UCG	D	902	-	-	5/21/67/67	0/6/6/6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	D	909	1	-	2/6/23/26	0/1/1/1
7	UCG	B	902	-	-	4/21/67/67	0/6/6/6
8	NAG	A	914	1	-	1/6/23/26	0/1/1/1
8	NAG	B	914	1	-	0/6/23/26	0/1/1/1
6	URI	C	901	-	-	6/6/22/22	0/2/2/2
8	NAG	C	910	1	-	0/6/23/26	0/1/1/1
8	NAG	C	915	1	-	0/6/23/26	0/1/1/1
7	UCG	A	902	-	-	5/21/67/67	0/6/6/6
8	NAG	D	913	1	-	2/6/23/26	0/1/1/1
8	NAG	D	914	1	-	2/6/23/26	0/1/1/1
8	NAG	B	909	1	-	2/6/23/26	0/1/1/1
8	NAG	D	912	1	-	0/6/23/26	0/1/1/1
8	NAG	A	910	1	-	1/6/23/26	0/1/1/1
8	NAG	B	913	1	-	2/6/23/26	0/1/1/1
8	NAG	B	915	1	-	0/6/23/26	0/1/1/1

The worst 5 of 61 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	902	UCG	C11-N3	-5.21	1.30	1.37
7	B	902	UCG	C11-N3	-4.72	1.30	1.37
7	B	902	UCG	C7-N1	-4.41	1.30	1.38
7	D	902	UCG	C7-N1	-4.04	1.31	1.38
7	B	902	UCG	C8-N1	-4.03	1.30	1.38

The worst 5 of 136 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
8	B	915	NAG	O5-C1-C2	-6.55	100.94	111.29
8	A	916	NAG	C1-O5-C5	6.34	120.78	112.19
7	A	902	UCG	N1-C8-N	6.31	123.27	114.89
7	D	902	UCG	N1-C8-N	6.13	123.03	114.89
8	C	916	NAG	C1-O5-C5	5.76	119.99	112.19

There are no chirality outliers.

5 of 51 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	902	UCG	C4-O1-P-O6
7	A	902	UCG	C4-O1-P-O7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
7	B	902	UCG	C4-O1-P-O7
7	C	902	UCG	C4-O1-P-O6
7	C	902	UCG	C4-O1-P-O7

There are no ring outliers.

11 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	902	UCG	5	0
6	D	901	URI	1	0
6	A	901	URI	1	0
6	B	901	URI	1	0
7	D	902	UCG	1	0
8	D	909	NAG	1	0
7	B	902	UCG	6	0
8	B	914	NAG	1	0
6	C	901	URI	1	0
7	A	902	UCG	4	0
8	D	913	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand UCG C 902



Ligand UCG D 902



Ligand UCG B 902



Ligand UCG A 902



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	747/811 (92%)	-0.13	17 (2%)	60	59	18, 32, 62, 90	0
1	B	738/811 (90%)	0.02	20 (2%)	54	53	19, 37, 70, 94	0
1	C	738/811 (90%)	0.25	53 (7%)	15	14	19, 43, 78, 112	0
1	D	735/811 (90%)	0.18	48 (6%)	18	18	21, 41, 86, 104	0
All	All	2958/3244 (91%)	0.08	138 (4%)	31	30	18, 38, 76, 112	0

The worst 5 of 138 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	64	TYR	8.1
1	C	761	THR	7.8
1	B	817	LEU	7.0
1	C	733	SER	6.5
1	D	64	TYR	6.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	BMA	H	3	11/12	0.83	0.12	41,50,53,59	0
3	BMA	M	3	11/12	0.83	0.31	57,64,72,75	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	BMA	O	3	11/12	0.88	0.32	47,57,62,64	0
2	MAN	E	4	11/12	0.89	0.15	34,44,49,53	0
5	BMA	K	3	11/12	0.89	0.15	44,50,64,65	0
5	MAN	K	4	11/12	0.89	0.22	60,70,80,95	0
3	BMA	N	3	11/12	0.90	0.10	40,48,56,56	0
2	BMA	E	3	11/12	0.90	0.10	36,42,49,51	0
3	BMA	F	3	11/12	0.91	0.28	40,46,54,56	0
4	NAG	G	2	14/15	0.92	0.15	29,36,42,52	0
3	BMA	I	3	11/12	0.93	0.17	39,45,54,61	0
4	NAG	P	2	14/15	0.93	0.11	33,39,45,49	0
3	BMA	L	3	11/12	0.93	0.30	41,47,51,57	0
3	NAG	M	2	14/15	0.93	0.17	34,40,46,47	0
3	NAG	F	2	14/15	0.94	0.12	26,30,34,36	0
5	NAG	K	2	14/15	0.94	0.09	34,39,48,48	0
3	NAG	L	2	14/15	0.94	0.13	25,30,35,39	0
4	NAG	P	1	14/15	0.94	0.10	27,30,35,49	0
3	NAG	H	2	14/15	0.96	0.08	31,35,41,42	0
4	NAG	G	1	14/15	0.96	0.09	21,23,26,26	0
3	NAG	O	2	14/15	0.96	0.14	31,36,44,52	0
4	NAG	J	1	14/15	0.96	0.13	21,22,29,31	0
4	NAG	J	2	14/15	0.96	0.09	32,36,40,48	0
3	NAG	H	1	14/15	0.97	0.15	26,30,36,52	0
3	NAG	F	1	14/15	0.97	0.07	20,23,25,26	0
3	NAG	M	1	14/15	0.97	0.08	24,26,29,32	0
2	NAG	E	1	14/15	0.97	0.17	19,21,25,29	0
3	NAG	I	2	14/15	0.97	0.09	23,26,33,42	0
3	NAG	N	1	14/15	0.97	0.15	24,26,31,35	0
3	NAG	N	2	14/15	0.97	0.07	28,32,41,41	0
5	NAG	K	1	14/15	0.97	0.14	28,31,36,46	0
2	NAG	E	2	14/15	0.97	0.12	24,28,34,38	0
3	NAG	O	1	14/15	0.97	0.09	27,31,33,35	0
3	NAG	L	1	14/15	0.97	0.08	22,24,26,26	0
3	NAG	I	1	14/15	0.98	0.07	19,22,23,23	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain E:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain F:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain H:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain I:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain L:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain M:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain N:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain O:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain G:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain J:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain P:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain K:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
8	NAG	B	909	14/15	0.84	0.26	51,71,83,93	0
7	UCG	B	902	47/47	0.85	0.20	31,51,91,100	0
8	NAG	D	913	14/15	0.86	0.15	50,58,70,72	0
8	NAG	D	912	14/15	0.89	0.12	43,54,65,68	0
7	UCG	D	902	47/47	0.89	0.14	33,48,79,96	0
8	NAG	A	915	14/15	0.90	0.11	45,52,61,66	0
8	NAG	B	915	14/15	0.90	0.18	43,49,61,64	0
8	NAG	B	912	14/15	0.91	0.21	43,56,64,69	0
7	UCG	A	902	47/47	0.91	0.15	26,41,85,94	0
8	NAG	C	916	14/15	0.92	0.12	49,55,65,65	0
8	NAG	A	913	14/15	0.92	0.13	40,49,56,61	0
8	NAG	A	914	14/15	0.92	0.18	48,58,67,68	0
8	NAG	C	910	14/15	0.94	0.17	43,51,60,64	0
8	NAG	A	916	14/15	0.94	0.18	35,46,57,58	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
8	NAG	B	913	14/15	0.94	0.08	42,49,60,62	0
7	UCG	C	902	47/47	0.94	0.10	34,42,61,70	0
8	NAG	D	914	14/15	0.94	0.15	48,57,62,64	0
8	NAG	B	914	14/15	0.95	0.11	34,41,44,57	0
8	NAG	C	915	14/15	0.95	0.17	33,46,55,57	0
8	NAG	C	914	14/15	0.96	0.20	41,50,66,73	0
6	URI	C	901	17/17	0.97	0.10	26,33,42,44	0
8	NAG	A	910	14/15	0.97	0.10	35,39,42,48	0
8	NAG	D	909	14/15	0.98	0.08	31,39,51,56	0
6	URI	B	901	17/17	0.98	0.07	18,22,36,37	0
6	URI	A	901	17/17	0.98	0.10	19,22,29,31	0
6	URI	D	901	17/17	0.98	0.10	21,24,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UCG B 902:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)







Electron density around UCG D 902:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around UCG A 902:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around UCG C 902:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.