



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 11, 2025 – 10:13 am BST

PDB ID : 9R0L / pdb\_00009r0l  
Title : Three dimensional structure of human carbonic anhydrase XII in complex with sulfonamide  
Authors : Eimonta, V.; Leitans, J.; Tars, K.  
Deposited on : 2025-04-24  
Resolution : 1.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.45.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1556 (1.46-1.46)
Clashscore	180529	1653 (1.46-1.46)
Ramachandran outliers	177936	1635 (1.46-1.46)
Sidechain outliers	177891	1635 (1.46-1.46)
RSRZ outliers	164620	1556 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	<div> <div>5%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div>.</div> </div> </div>
1	B	263	<div> <div>4%</div> <div> <div></div> <div>91%</div> <div>6%</div> <div>..</div> </div> </div>
1	C	263	<div> <div>5%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div>..</div> </div> </div>
1	D	263	<div> <div>6%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div>..</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9246 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbonic anhydrase 12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	1	0
			2083	1326	354	395	8			
1	B	260	Total	C	N	O	S	0	0	0
			2084	1326	355	396	7			
1	C	260	Total	C	N	O	S	0	0	0
			2084	1326	354	397	7			
1	D	260	Total	C	N	O	S	0	1	0
			2087	1328	354	397	8			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	-	expression tag	UNP O43570
B	2	ALA	-	expression tag	UNP O43570
C	2	ALA	-	expression tag	UNP O43570
D	2	ALA	-	expression tag	UNP O43570

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		
2	D	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 3-(cyclooctylamino)-2,6-difluoro-4-((3-hydroxypropyl)sulfonyl)-5-(piperidin-1-yl)benzenesulfonamide (CCD ID: A1H9A) (formula: C<sub>22</sub>H<sub>35</sub>F<sub>2</sub>N<sub>3</sub>O<sub>5</sub>S<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total 34	C 22	F 2	N 3	O 5	S 2	0	0
3	B	1	Total 34	C 22	F 2	N 3	O 5	S 2	0	0
3	C	1	Total 34	C 22	F 2	N 3	O 5	S 2	0	0
3	D	1	Total 34	C 22	F 2	N 3	O 5	S 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	197	Total	O	0	0
			197	197		
4	B	209	Total	O	0	0
			209	209		
4	C	196	Total	O	0	0
			196	196		
4	D	166	Total	O	0	0
			166	166		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Carbonic anhydrase 12




- Molecule 1: Carbonic anhydrase 12



- Molecule 1: Carbonic anhydrase 12



- Molecule 1: Carbonic anhydrase 12



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.57Å 76.70Å 78.87Å 62.61° 79.07° 72.89°	Depositor
Resolution (Å)	44.41 – 1.45 44.41 – 1.45	Depositor EDS
% Data completeness (in resolution range)	96.6 (44.41-1.45) 96.6 (44.41-1.45)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.56 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.184 , 0.209 0.194 , 0.217	Depositor DCC
$R_{free}$ test set	8053 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	10.5	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 33.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9246	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1H9A, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.11	1/2149 (0.0%)	1.19	1/2928 (0.0%)
1	B	1.13	2/2147 (0.1%)	1.21	4/2925 (0.1%)
1	C	1.12	2/2147 (0.1%)	1.22	4/2925 (0.1%)
1	D	1.14	4/2153 (0.2%)	1.20	4/2933 (0.1%)
All	All	1.12	9/8596 (0.1%)	1.20	13/11711 (0.1%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	25	GLY	C-O	7.71	1.31	1.24
1	A	182	GLU	C-O	6.54	1.31	1.23
1	B	132	LYS	C-O	6.14	1.31	1.23
1	D	89	HIS	ND1-CE1	-5.91	1.26	1.32
1	D	32	ASP	C-O	-5.89	1.16	1.24

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	73	ASP	CA-CB-CG	13.02	125.62	112.60
1	B	131	ASN	CB-CA-C	7.36	122.95	111.02
1	D	80	GLN	CB-CA-C	-6.50	100.41	110.81
1	B	4	LYS	CA-C-O	-6.33	110.03	120.80
1	B	131	ASN	N-CA-C	-5.96	104.66	112.41

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2083	0	1974	13	0
1	B	2084	0	1975	16	0
1	C	2084	0	1973	18	0
1	D	2087	0	1978	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	34	0	0	1	0
3	B	34	0	0	0	0
3	C	34	0	0	3	0
3	D	34	0	0	5	0
4	A	197	0	0	2	0
4	B	209	0	0	2	0
4	C	196	0	0	2	0
4	D	166	0	0	6	0
All	All	9246	0	7900	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 60 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:167:GLN:NE2	1:D:230:MET:SD	2.41	0.94
1:D:227[B]:CYS:SG	1:D:235(A):ARG:HB2	2.09	0.92
1:D:131:ASN:HB2	4:D:562:HOH:O	1.80	0.80
1:D:62:ASN:HD21	3:D:302:A1H9A:C15	1.93	0.80
1:B:167:GLN:HE22	1:B:230:MET:HE3	1.51	0.73

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/263 (98%)	252 (97%)	7 (3%)	0	100	100
1	B	258/263 (98%)	254 (98%)	4 (2%)	0	100	100
1	C	258/263 (98%)	252 (98%)	6 (2%)	0	100	100
1	D	259/263 (98%)	254 (98%)	5 (2%)	0	100	100
All	All	1034/1052 (98%)	1012 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/234 (99%)	226 (98%)	5 (2%)	47	16
1	B	231/234 (99%)	227 (98%)	4 (2%)	56	24
1	C	231/234 (99%)	225 (97%)	6 (3%)	41	11
1	D	232/234 (99%)	229 (99%)	3 (1%)	65	35
All	All	925/936 (99%)	907 (98%)	18 (2%)	50	20

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	256	SER
1	D	153	ASP
1	D	73	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	256	SER
1	C	249	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	108	HIS
1	D	199	ASN
1	C	50	GLN
1	D	244	GLN
1	D	75	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	A1H9A	B	302	2	34,36,36	2.36	11 (32%)	42,52,52	2.34	14 (33%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	A1H9A	D	302	2	34,36,36	2.93	8 (23%)	42,52,52	2.56	13 (30%)
3	A1H9A	A	302	2	34,36,36	2.24	11 (32%)	42,52,52	2.71	14 (33%)
3	A1H9A	C	302	2	34,36,36	2.46	9 (26%)	42,52,52	2.64	14 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1H9A	B	302	2	-	9/25/43/43	0/3/3/3
3	A1H9A	D	302	2	-	10/25/43/43	0/3/3/3
3	A1H9A	A	302	2	-	9/25/43/43	0/3/3/3
3	A1H9A	C	302	2	-	7/25/43/43	0/3/3/3

The worst 5 of 39 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	302	A1H9A	C1-C2	7.69	1.51	1.39
3	D	302	A1H9A	C2-C3	7.25	1.51	1.39
3	C	302	A1H9A	C3-S7	-6.89	1.68	1.79
3	D	302	A1H9A	C3-S7	-6.67	1.68	1.79
3	D	302	A1H9A	C5-C4	6.40	1.49	1.39

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302	A1H9A	O20-S19-O21	-9.17	107.72	118.44
3	D	302	A1H9A	C18-N13-C14	7.70	128.51	111.52
3	A	302	A1H9A	C4-C3-C2	-7.67	111.29	116.67
3	B	302	A1H9A	C4-C3-C2	-7.66	111.30	116.67
3	C	302	A1H9A	C18-N13-C14	7.35	127.75	111.52

There are no chirality outliers.

5 of 35 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	A1H9A	C2-C3-S7-O8
3	A	302	A1H9A	C4-C3-S7-N10
3	A	302	A1H9A	C4-C3-S7-O8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	302	A1H9A	C1-C6-S19-O21
3	A	302	A1H9A	C1-C6-S19-O20


There are no ring outliers.

3 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	302	A1H9A	5	0
3	A	302	A1H9A	1	0
3	C	302	A1H9A	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


## Ligand A1H9A B 302




Bond lengths



Bond angles




Torsions




Rings


## Ligand A1H9A D 302




Bond lengths



Bond angles





Torsions



Rings

## Ligand A1H9A A 302





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	260/263 (98%)	0.42	13 (5%) 35 35	5, 12, 24, 35	1 (0%)
1	B	260/263 (98%)	0.22	10 (3%) 44 45	4, 10, 22, 33	0
1	C	260/263 (98%)	0.33	12 (4%) 38 38	5, 12, 23, 37	0
1	D	260/263 (98%)	0.56	16 (6%) 28 27	6, 14, 27, 39	1 (0%)
All	All	1040/1052 (98%)	0.38	51 (4%) 36 35	4, 12, 24, 39	2 (0%)

The worst 5 of 51 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	8	PHE	6.7
1	D	8	PHE	3.9
1	D	250	LEU	3.7
1	D	230	MET	3.4
1	B	247	ASP	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	A1H9A	D	302	34/34	0.78	0.18	24,33,39,43	0
3	A1H9A	C	302	34/34	0.80	0.16	20,26,29,32	0
3	A1H9A	A	302	34/34	0.93	0.11	11,15,24,26	0
3	A1H9A	B	302	34/34	0.93	0.11	10,14,21,30	0
2	ZN	A	301	1/1	1.00	0.02	6,6,6,6	0
2	ZN	B	301	1/1	1.00	0.01	5,5,5,5	0
2	ZN	C	301	1/1	1.00	0.01	6,6,6,6	0
2	ZN	D	301	1/1	1.00	0.03	7,7,7,7	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around A1H9A D 302:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around A1H9A C 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around A1H9A A 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around A1H9A B 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.