



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2025 – 11:33 am BST

PDB ID : 9R3K / pdb_00009r3k
Title : Crystal structure of human MAO B in complex with ((E)-3-(3-nitrophenyl)-1-(3-(trifluoromethyl)phenyl)prop-2-en-1-one (4b)
Authors : Marchese, S.; Binda, C.
Deposited on : 2025-05-05
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.45.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4274 (1.60-1.60)
Clashscore	180529	4682 (1.60-1.60)
Ramachandran outliers	177936	4583 (1.60-1.60)
Sidechain outliers	177891	4582 (1.60-1.60)
RSRZ outliers	164620	4272 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	519	
1	BBB	519	

2 Entry composition i


There are 5 unique types of molecules in this entry. The entry contains 9049 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amine oxidase [flavin-containing] B.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	500	Total	C	N	O	S	0	6	0
			4006	2564	683	732	27			
1	BBB	495	Total	C	N	O	S	0	3	0
			3957	2530	677	724	26			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	AAA	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	BBB	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is ({E})-3-[3-[oxidanyl(oxidanylidene)-\$l^{\wedge}\{4\}\$-azanyl]phenyl]-1-[3-(trifluoromet hyl)phenyl]prop-2-en-1-one (CCD ID: A1JC6) (formula: C₁₆H₁₀F₃NO₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	AAA	1	Total	C	F	N	O	0	0
			23	16	3	1	3		
3	BBB	1	Total	C	F	N	O	0	0
			23	16	3	1	3		

- Molecule 4 is N-DODECYL-N,N-DIMETHYL-3-AMMONIO-1-PROPANESULFONATE (CCD ID: C15) (formula: C₁₇H₃₈NO₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	AAA	1	Total	C	N	O	S	0	0
			15	10	1	3	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	BBB	1	Total	C	N	O	S	0	0
			11	6	1	3	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	430	Total	O	0	0
			430	430		
5	BBB	478	Total	O	0	0
			478	478		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Amine oxidase [flavin-containing] B



- Molecule 1: Amine oxidase [flavin-containing] B



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.22Å 222.16Å 86.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.27 – 1.60 47.27 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.27-1.60) 99.9 (47.27-1.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.89 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.140 , 0.162 0.154 , 0.176	Depositor DCC
R_{free} test set	4135 reflections (2.51%)	wwPDB-VP
Wilson B-factor (Å ²)	16.0	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.009 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.012 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9049	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, C15, A1JC6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.07	3/4121 (0.1%)	1.18	2/5592 (0.0%)
1	BBB	1.10	2/4063 (0.0%)	1.19	7/5513 (0.1%)
All	All	1.09	5/8184 (0.1%)	1.19	9/11105 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	376	GLU	CD-OE2	5.98	1.36	1.25
1	BBB	96	SER	CA-CB	-5.81	1.45	1.53
1	AAA	226	GLY	C-O	5.67	1.31	1.24
1	AAA	38	ARG	C-O	5.41	1.30	1.23
1	BBB	468	GLU	C-O	5.12	1.30	1.23

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	336	ASN	CA-CB-CG	-8.17	104.43	112.60
1	BBB	415	ARG	CB-CA-C	7.07	122.40	111.02
1	BBB	431	HIS	CA-CB-CG	6.18	119.98	113.80
1	BBB	127	ARG	CG-CD-NE	-6.07	98.64	112.00
1	AAA	124	ASP	CA-CB-CG	5.62	118.22	112.60
1	BBB	335	GLY	N-CA-C	-5.49	107.53	115.27
1	AAA	251	ASN	N-CA-C	-5.19	107.12	112.93
1	BBB	349	ALA	N-CA-C	-5.16	105.66	111.28
1	BBB	353	ALA	CA-C-O	-5.11	114.33	120.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	4006	0	4026	27	1
1	BBB	3957	0	3961	25	0
2	AAA	53	0	29	0	0
2	BBB	53	0	29	0	0
3	AAA	23	0	0	0	0
3	BBB	23	0	0	0	0
4	AAA	15	0	21	0	0
4	BBB	11	0	13	0	0
5	AAA	430	0	0	1	0
5	BBB	478	0	0	3	0
All	All	9049	0	8079	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:28:LEU:HD11	1:AAA:454:MET:HE1	1.61	0.82
1:BBB:285[A]:MET:HE2	1:BBB:414:LEU:HD23	1.69	0.75
1:BBB:285[A]:MET:CE	1:BBB:414:LEU:HD23	2.17	0.75
1:AAA:280:MET:CE	5:BBB:1015:HOH:O	2.34	0.74
1:AAA:251:ASN:HD22	1:AAA:251:ASN:H	1.39	0.70
1:BBB:117:ASN:HD22	1:BBB:120:ARG:HH21	1.40	0.68
1:AAA:117:ASN:HD22	1:AAA:120:ARG:HH21	1.40	0.67
1:BBB:251:ASN:HD22	1:BBB:251:ASN:H	1.44	0.66
1:AAA:451:LEU:HD23	1:AAA:454:MET:HE1	1.83	0.61
1:AAA:451:LEU:HD23	1:AAA:454:MET:CE	2.33	0.58
1:AAA:163:GLN:HB3	1:AAA:318:ASP:OD2	2.03	0.58
1:AAA:248:GLU:HB2	1:AAA:254[B]:MET:HE1	1.86	0.57
1:BBB:93:LYS:HE3	5:BBB:812:HOH:O	2.09	0.53
1:AAA:451:LEU:HA	1:AAA:454:MET:CE	2.40	0.52
1:BBB:321:GLU:H	1:BBB:321:GLU:CD	2.17	0.52
1:AAA:280:MET:HE3	5:BBB:1015:HOH:O	2.05	0.52
1:AAA:410:TYR:OH	1:BBB:350:ARG:HD2	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:93:LYS:HA	1:BBB:93:LYS:CE	2.41	0.51
5:AAA:907:HOH:O	1:BBB:149:LYS:HD3	2.13	0.49
1:AAA:106:VAL:HG21	1:AAA:111:THR:HG22	1.94	0.49
1:AAA:321:GLU:H	1:AAA:321:GLU:CD	2.21	0.48
1:BBB:280:MET:HE3	1:BBB:281:MET:HG2	1.97	0.47
1:AAA:280:MET:HE1	1:BBB:353:ALA:HB1	1.97	0.47
1:BBB:233:ARG:HG3	1:BBB:251:ASN:HD21	1.79	0.47
1:AAA:270:MET:HE1	1:BBB:268:LEU:HD23	1.97	0.46
1:AAA:268:LEU:HD23	1:BBB:270:MET:HE1	1.98	0.46
1:BBB:280:MET:HB3	1:BBB:280:MET:HE2	1.72	0.45
1:AAA:121:THR:HG21	1:AAA:155:LEU:HD21	1.98	0.45
1:AAA:450:ILE:HG22	1:AAA:454:MET:HE2	1.99	0.44
1:AAA:106:VAL:HG11	1:AAA:112:TYR:HA	1.98	0.44
1:BBB:309:LYS:NZ	1:BBB:373:GLY:O	2.50	0.43
1:BBB:332:LYS:HB3	1:BBB:333:PRO:HD2	2.00	0.43
1:AAA:495:LEU:HD12	1:AAA:495:LEU:HA	1.93	0.42
1:BBB:146:MET:HE3	1:BBB:146:MET:HB2	1.89	0.42
1:BBB:254:MET:HB2	1:BBB:254:MET:HE2	1.76	0.42
1:BBB:495:LEU:C	1:BBB:496:ILE:HG13	2.45	0.42
1:BBB:285[B]:MET:CE	1:BBB:414:LEU:HD23	2.51	0.41
1:AAA:451:LEU:HA	1:AAA:454:MET:HE3	2.02	0.41
1:AAA:117:ASN:HD22	1:AAA:120:ARG:NH2	2.13	0.41
1:AAA:108:ASN:OD1	1:AAA:108:ASN:C	2.64	0.41
1:AAA:280:MET:HE3	1:AAA:280:MET:HB3	1.56	0.41
1:BBB:17:MET:CE	1:BBB:39:VAL:HG11	2.50	0.41
1:AAA:280:MET:HE1	1:BBB:387:ASN:ND2	2.36	0.41
1:BBB:414:LEU:HD23	1:BBB:414:LEU:HA	1.90	0.41
1:AAA:254[A]:MET:HE2	1:AAA:254[A]:MET:HB3	1.99	0.40
1:AAA:389:CYS:CB	1:BBB:280:MET:HG3	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:475:GLN:NE2	1:AAA:475:GLN:NE2[3_656]	1.79	0.41

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	504/519 (97%)	492 (98%)	12 (2%)	0	100	100
1	BBB	496/519 (96%)	484 (98%)	12 (2%)	0	100	100
All	All	1000/1038 (96%)	976 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	435/443 (98%)	427 (98%)	8 (2%)	54	31
1	BBB	428/443 (97%)	424 (99%)	4 (1%)	75	62
All	All	863/886 (97%)	851 (99%)	12 (1%)	65	43

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	251	ASN
1	AAA	254[A]	MET
1	AAA	254[B]	MET
1	AAA	280	MET
1	AAA	321	GLU
1	AAA	350	ARG
1	AAA	495	LEU
1	AAA	501	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	BBB	93	LYS
1	BBB	251	ASN
1	BBB	280	MET
1	BBB	350	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	A1JC6	AAA	602	-	24,24,24	0.65	0	34,34,34	1.13	2 (5%)
4	C15	BBB	603	-	10,10,21	0.91	1 (10%)	14,15,26	0.98	1 (7%)
2	FAD	BBB	601	1	53,58,58	1.10	2 (3%)	68,89,89	1.71	19 (27%)
2	FAD	AAA	601	1	53,58,58	1.10	3 (5%)	68,89,89	1.50	14 (20%)
4	C15	AAA	603	-	14,14,21	0.86	1 (7%)	18,19,26	0.82	0
3	A1JC6	BBB	602	-	24,24,24	0.90	2 (8%)	34,34,34	1.11	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1JC6	AAA	602	-	-	0/19/19/19	0/2/2/2
4	C15	BBB	603	-	-	1/8/8/21	-
2	FAD	BBB	601	1	-	1/30/50/50	0/6/6/6
2	FAD	AAA	601	1	-	1/30/50/50	0/6/6/6
4	C15	AAA	603	-	-	1/14/14/21	-
3	A1JC6	BBB	602	-	-	0/19/19/19	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	601	FAD	C4X-N5	3.87	1.38	1.30
2	BBB	601	FAD	C4X-N5	3.72	1.38	1.30
4	AAA	603	C15	O3S-S1	2.66	1.57	1.47
2	AAA	601	FAD	C9A-N10	-2.55	1.36	1.41
4	BBB	603	C15	O3S-S1	2.50	1.56	1.47
3	BBB	602	A1JC6	O3-C8	2.20	1.29	1.24
3	BBB	602	A1JC6	C14-C9	-2.19	1.36	1.39
2	BBB	601	FAD	C9A-N10	-2.04	1.37	1.41
2	AAA	601	FAD	C1'-C2'	2.02	1.55	1.52

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	BBB	601	FAD	C4X-C10-N10	4.20	122.62	116.48
2	AAA	601	FAD	C5A-C6A-N6A	3.66	125.91	120.35
3	AAA	602	A1JC6	C11-C12-C13	-3.56	117.02	120.76
2	AAA	601	FAD	C4-N3-C2	-3.45	119.27	125.64
2	BBB	601	FAD	C5A-C6A-N6A	3.40	125.52	120.35
3	BBB	602	A1JC6	C11-C12-C13	-3.34	117.25	120.76
2	BBB	601	FAD	C9A-C9-C8	3.28	125.90	119.30
2	BBB	601	FAD	C4X-C4-N3	3.13	121.15	113.19
2	BBB	601	FAD	C9A-N10-C10	-3.13	115.89	120.77
2	BBB	601	FAD	N3A-C2A-N1A	-2.97	124.03	128.68
2	BBB	601	FAD	O4-C4-C4X	-2.83	119.09	126.60
2	BBB	601	FAD	C10-N1-C2	2.83	122.56	116.90
2	BBB	601	FAD	C4-N3-C2	-2.83	120.42	125.64
3	AAA	602	A1JC6	C10-C9-C14	-2.80	115.92	119.24
2	BBB	601	FAD	O4B-C1B-C2B	-2.75	102.91	106.93

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	601	FAD	C1'-N10-C9A	2.75	125.09	120.51
2	BBB	601	FAD	C5'-C4'-C3'	-2.72	106.94	112.20
2	AAA	601	FAD	C5'-C4'-C3'	-2.69	107.00	112.20
2	BBB	601	FAD	C4-C4X-N5	2.49	121.77	118.23
2	AAA	601	FAD	C9A-C5X-N5	-2.44	119.78	122.43
2	AAA	601	FAD	C9A-C9-C8	2.41	124.16	119.30
2	BBB	601	FAD	C10-C4X-N5	-2.41	119.74	124.86
2	AAA	601	FAD	C4X-C4-N3	2.40	119.29	113.19
2	BBB	601	FAD	C9-C9A-C5X	-2.40	115.58	120.11
3	BBB	602	A1JC6	F2-C15-C13	-2.39	107.67	112.93
2	BBB	601	FAD	C2A-N1A-C6A	2.38	122.83	118.75
2	AAA	601	FAD	C10-C4X-N5	-2.37	119.84	124.86
2	BBB	601	FAD	C5X-C9A-N10	2.30	120.33	117.95
2	AAA	601	FAD	C4X-C10-N10	2.29	119.83	116.48
2	BBB	601	FAD	O3'-C3'-C4'	-2.27	103.34	108.81
2	BBB	601	FAD	C4X-C10-N1	-2.25	119.52	124.73
2	AAA	601	FAD	O4-C4-C4X	-2.21	120.75	126.60
2	AAA	601	FAD	C8M-C8-C7	2.17	125.19	120.74
2	BBB	601	FAD	C4A-C5A-N7A	2.16	111.65	109.40
2	AAA	601	FAD	O3'-C3'-C4'	-2.14	103.65	108.81
4	BBB	603	C15	O1S-S1-C1	-2.12	104.36	106.92
2	AAA	601	FAD	C4A-C5A-N7A	-2.12	107.19	109.40
3	BBB	602	A1JC6	C10-C9-C14	-2.06	116.80	119.24
2	AAA	601	FAD	O4B-C4B-C3B	2.03	109.12	105.11

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	BBB	603	C15	S1-C1-C2-C3
4	AAA	603	C15	S1-C1-C2-C3
2	BBB	601	FAD	O4B-C4B-C5B-O5B
2	AAA	601	FAD	O4B-C4B-C5B-O5B

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.


Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand FAD BBB 601




Ligand FAD AAA 601



Ligand C15 AAA 603



Ligand A1JC6 BBB 602



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	500/519 (96%)	-0.31	10 (2%) 64 67	8, 17, 34, 78	6 (1%)
1	BBB	495/519 (95%)	-0.47	6 (1%) 76 79	9, 15, 31, 71	3 (0%)
All	All	995/1038 (95%)	-0.39	16 (1%) 70 73	8, 16, 33, 78	9 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	501	ILE	6.9
1	BBB	495	LEU	4.8
1	BBB	496	ILE	4.6
1	AAA	498	LEU	4.1
1	BBB	107	TRP	3.9
1	BBB	494	ARG	3.7
1	AAA	499	THR	3.6
1	AAA	500	THR	3.6
1	AAA	252	HIS	3.2
1	AAA	319	GLY	3.0
1	AAA	473	PRO	2.7
1	AAA	107	TRP	2.4
1	BBB	252	HIS	2.3
1	BBB	493	LEU	2.2
1	AAA	495	LEU	2.1
1	AAA	106	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	C15	AAA	603	15/22	0.77	0.17	26,33,61,66	0
4	C15	BBB	603	11/22	0.82	0.14	31,40,71,75	0
3	A1JC6	AAA	602	23/23	0.93	0.10	19,22,30,40	0
3	A1JC6	BBB	602	23/23	0.94	0.08	18,21,26,36	0
2	FAD	AAA	601	53/53	0.99	0.03	10,12,13,15	0
2	FAD	BBB	601	53/53	0.99	0.03	9,11,12,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around C15 AAA 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around C15 BBB 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around A1JC6 AAA 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1JC6 BBB 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD AAA 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FAD BBB 601:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.