# wwPDB X-ray Structure Validation Summary Report (i)

Oct 23, 2021 – 11:15 AM EDT

| | | |
|---|---|---|
| PDB ID | : | 1RDX |
| Title | : | R-STATE STRUCTURE OF THE ARG 243 TO ALA MUTANT OF PIG KIDNEY FRUCTOSE 1,6-BISPHOSPHATASE EXPRESSED IN E. COLI |
| Authors | : | Stec, B.; Abraham, R.; Giroux, E.; Kantrowitz, E.R. |
| Deposited on | : | 1996-05-17 |
| Resolution | : | 2.75 Å(reported) |

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

| | | |
|---|---|---|
| MolProbity | : | 4.02b-467 |
| Mogul | : | 1.8.5 (274361), CSD as541be (2020) |
| Xtriage (Phenix) | : | NOT EXECUTED |
| EDS | : | NOT EXECUTED |
| buster-report | : | 1.1.7 (2018) |
| Percentile statistics | : | 20191225.v01 (using entries in the PDB archive December 25th 2019) |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.23.2 |

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|---|---|---|
| Clashscore | 141614 | 1277 (2.78-2.74) |
| Ramachandran outliers | 138981 | 1257 (2.78-2.74) |
| Sidechain outliers | 138945 | 1257 (2.78-2.74) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

| Mol | Chain | Length | Quality of chain |
|---|---|---|---|
| 1 | A | 337 | 23% 54% 21% • |
| 1 | B | 337 | 20% 60% 18% • |

# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5223 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FRUCTOSE 1,6-BISPHOSPHATASE.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | Total | C | N | O | S | | | |
| 1 | A | 337 | 2566 | 1630 | 431 | 490 | 15 | 0 | 0 | 0 |
| 1 | B | 337 | 2566 | 1630 | 431 | 490 | 15 | 0 | 0 | 0 |

There are 8 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|---|---|---|---|---|---|
| A | 20 | GLN | GLU | conflict | UNP P00636 |
| A | 96 | THR | SER | conflict | UNP P00636 |
| A | 199 | ASN | ASP | conflict | UNP P00636 |
| A | 243 | ALA | ARG | engineered mutation | UNP P00636 |
| B | 20 | GLN | GLU | conflict | UNP P00636 |
| B | 96 | THR | SER | conflict | UNP P00636 |
| B | 199 | ASN | ASP | conflict | UNP P00636 |
| B | 243 | ALA | ARG | engineered mutation | UNP P00636 |

- Molecule 2 is 6-O-phosphono-beta-D-fructofuranose (three-letter code: F6P) (formula: $C_6H_{13}O_9P$).




| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|---|---|---|---|---|---|---|---|---|
| | | | Total | C | O | P | | |
| 2 | A | 1 | 16 | 6 | 9 | 1 | 0 | 0 |
| | | | Total | C | O | P | | |
| 2 | B | 1 | 16 | 6 | 9 | 1 | 0 | 0 |

- Molecule 3 is water.

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|---|---|---|---|---|---|---|
| | | | Total | O | | |
| 3 | A | 29 | 29 | 29 | 0 | 0 |
| | | | Total | O | | |
| 3 | B | 30 | 30 | 30 | 0 | 0 |

# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: FRUCTOSE 1,6-BISPHOSPHATASE

Chain A: 23% 54% 21% •




• Molecule 1: FRUCTOSE 1,6-BISPHOSPHATASE

Chain B: 20% 60% 18% •




F262
M263
Y264
P265
A266
N267
K268
K269
S270
P271
K272
G273
K274
L275
R276
L277
L278
Y279
E280
C281
N282
P283
M284
A285
Y286
V287
M288
E289
K290
A291
G292
G293
L294
A295
T296
T297
G298
K299
E300
A301
V302
L303
D304
I305
V306
P307
T308
D309
I310
H311
Q312
R313
A314
P315
I316
I317
L318
G319
S320
P321

E322
D323
V324
T325
E326
L327
L328
E329
I330
Y331
Q332
K333
H334
A335
A336
K337

# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

| Property | Value | Source |
|---|---|---|
| Space group | P 32 2 1 | Depositor |
| Cell constants<br>a, b, c, $\alpha$, $\beta$, $\gamma$ | 131.31Å 131.31Å 66.40Å<br>90.00° 90.00° 120.00° | Depositor |
| Resolution (Å) | 10.00 – 2.75 | Depositor |
| % Data completeness<br>(in resolution range) | (Not available) (10.00-2.75) | Depositor |
| $R_{merge}$ | 0.14 | Depositor |
| $R_{sym}$ | (Not available) | Depositor |
| Refinement program | X-PLOR | Depositor |
| R, $R_{free}$ | 0.223 , 0.297 | Depositor |
| Estimated twinning fraction | No twinning to report. | Xtriage |
| Total number of atoms | 5223 | wwPDB-VP |
| Average B, all atoms (Å$^2$) | 41.0 | wwPDB-VP |

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: F6P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|---|---|---|---|---|---|
| | | RMSZ | #$\|Z\|$ >5 | RMSZ | #$\|Z\|$ >5 |
| 1 | A | 0.58 | 0/2609 | 0.92 | 5/3531 (0.1%) |
| 1 | B | 0.56 | 0/2609 | 0.89 | 3/3531 (0.1%) |
| All | All | 0.57 | 0/5218 | 0.91 | 8/7062 (0.1%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|---|---|---|---|
| 1 | B | 0 | 1 |

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^o$) | Ideal($^o$) |
|---|---|---|---|---|---|---|---|
| 1 | A | 153 | LEU | CA-CB-CG | 7.19 | 131.84 | 115.30 |
| 1 | A | 169 | SER | N-CA-C | -6.88 | 92.44 | 111.00 |
| 1 | A | 33 | LEU | CA-CB-CG | 6.72 | 130.76 | 115.30 |
| 1 | B | 298 | GLY | N-CA-C | -6.58 | 96.66 | 113.10 |
| 1 | A | 306 | VAL | N-CA-C | -5.77 | 95.42 | 111.00 |

There are no chirality outliers.

All (1) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|---|---|---|---|---|
| 1 | B | 215 | TYR | Sidechain |

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|---|---|---|---|---|---|---|
| 1 | A | 2566 | 0 | 2620 | 396 | 0 |
| 1 | B | 2566 | 0 | 2620 | 427 | 0 |
| 2 | A | 16 | 0 | 11 | 1 | 0 |
| 2 | B | 16 | 0 | 11 | 1 | 0 |
| 3 | A | 29 | 0 | 0 | 2 | 0 |
| 3 | B | 30 | 0 | 0 | 2 | 0 |
| All | All | 5223 | 0 | 5262 | 791 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 76.

The worst 5 of 791 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|---|---|---|---|
| 1:A:297:THR:HB | 1:A:315:PRO:HG2 | 1.21 | 1.15 |
| 1:A:97:GLU:HB3 | 1:A:279:TYR:HE2 | 1.08 | 1.13 |
| 1:A:224:THR:HA | 1:A:227:ILE:HD12 | 1.34 | 1.09 |
| 1:A:174:VAL:HA | 1:A:183:CYS:HA | 1.33 | 1.09 |
| 1:B:114:VAL:HB | 1:B:139:TYR:HB2 | 1.33 | 1.07 |

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles |
|---|---|---|---|---|---|---|
| 1 | A | 335/337 (99%) | 209 (62%) | 84 (25%) | 42 (12%) | 0 0 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles |
|---|---|---|---|---|---|---|
| 1 | B | 335/337 (99%) | 206 (62%) | 97 (29%) | 32 (10%) | 0 0 |
| All | All | 670/674 (99%) | 415 (62%) | 181 (27%) | 74 (11%) | 0 0 |

5 of 74 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | A | 3 | GLN |
| 1 | A | 8 | THR |
| 1 | A | 33 | LEU |
| 1 | A | 34 | LEU |
| 1 | A | 56 | LEU |

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles |
|---|---|---|---|---|---|
| 1 | A | 278/278 (100%) | 220 (79%) | 58 (21%) | 1 1 |
| 1 | B | 278/278 (100%) | 223 (80%) | 55 (20%) | 1 1 |
| All | All | 556/556 (100%) | 443 (80%) | 113 (20%) | 1 1 |

5 of 113 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | A | 331 | TYR |
| 1 | B | 325 | THR |
| 1 | B | 80 | LEU |
| 1 | B | 324 | VAL |
| 1 | B | 254 | ARG |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | B | 332 | GLN |
| 1 | B | 334 | HIS |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type |
|---|---|---|---|
| 1 | B | 158 | ASN |
| 1 | B | 236 | ASN |
| 1 | B | 267 | ASN |

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Counts | RMSZ | $\#\|Z\| > 2$ | Counts | RMSZ | $\#\|Z\| > 2$ |
| 2 | F6P | A | 338 | - | 15,16,16 | 1.07 | 1 (6%) | 17,25,25 | 0.97 | 1 (5%) |
| 2 | F6P | B | 338 | - | 15,16,16 | 1.36 | 1 (6%) | 17,25,25 | 1.45 | 2 (11%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|---|---|---|---|---|---|---|---|
| 2 | F6P | A | 338 | - | - | 4/9/28/28 | 0/1/1/1 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|---|---|---|---|---|---|---|---|
| 2 | F6P | B | 338 | - | - | 1/9/28/28 | 0/1/1/1 |

All (2) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|---|---|---|---|---|---|---|---|
| 2 | B | 338 | F6P | O5-C2 | -4.15 | 1.36 | 1.43 |
| 2 | A | 338 | F6P | O2-C2 | 2.45 | 1.44 | 1.40 |

All (3) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^o$) | Ideal($^o$) |
|---|---|---|---|---|---|---|---|
| 2 | B | 338 | F6P | O2-C2-O5 | -3.50 | 102.74 | 109.50 |
| 2 | B | 338 | F6P | P-O6-C6 | 2.32 | 124.69 | 118.30 |
| 2 | A | 338 | F6P | O6-P-O1P | 2.12 | 112.41 | 106.47 |

There are no chirality outliers.

All (5) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|---|---|---|---|---|
| 2 | A | 338 | F6P | O1-C1-C2-O2 |
| 2 | A | 338 | F6P | O1-C1-C2-C3 |
| 2 | A | 338 | F6P | O1-C1-C2-O5 |
| 2 | B | 338 | F6P | C5-C6-O6-P |
| 2 | A | 338 | F6P | O5-C5-C6-O6 |

There are no ring outliers.

2 monomers are involved in 2 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|---|---|---|---|---|---|
| 2 | A | 338 | F6P | 1 | 0 |
| 2 | B | 338 | F6P | 1 | 0 |

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

**Ligand F6P A 338**




**Ligand F6P B 338**




## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

## 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

## 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.