Full wwPDB X-ray Structure Validation Report (i)

Oct 8, 2023 – 01:19 PM EDT

PDB ID	:	4RE6
Title	:	Acylaminoacyl peptidase complexed with a chloromethylketone inhibitor
Authors	:	Menyhard, D.K.; Orgovan, Z.; Szeltner, Z.; Szamosi, I.; Harmat, V.
Deposited on	:	2014-09-22
Resolution	:	2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	582	.% 91% 8% ••
1	B	582	2% 92% 7% ••
1	C	582	.% 92% 7% ••
1	D	582	2% 88% 10% ••

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 18074 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acylamino-acid-releasing enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	577	4383	2771	769	830	13	0	4	0
1	B	574	4277	2712	737	816	12	0	2	0
1	C	577	4376	2771	769	823	13	0	4	0
1	D	577	4315	2736	745	821	13	0	6	0

- Molecule 2 is N-[(benzyloxy)carbonyl]glycyl-N-[(2S,3R)-4-chloro-3-hydroxy-1-phenylbutan-2-yl]glycinamide (three-letter code: Y3A) (formula: $C_{22}H_{26}ClN_3O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	23	15	3	5	0	0
2	C	1	20	14	3	3	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
3	A	2	2	2	0	0
			Total	Cl		
3	B	2	2	2	0	0
			Total	Cl		
3	C	2	2	2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	174	174	174	0	0
			Total	O		
4	B	154	154	154	0	0
			Total	O		
4	C	204	204	204	0	0
			Total	O		
4	D	142	142	142	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Acylamino-acid-releasing enzyme

Chain A:





• Molecule 1: Acylamino-acid-releasing enzyme

Chain B:





• Molecule 1: Acylamino-acid-releasing enzyme

Chain C:





• Molecule 1: Acylamino-acid-releasing enzyme

Chain D:



S196
L235
K238
S241
R264
R265
E266
G267
E279
N284
R287
R292
L322
P323
E324
R327
G332
E373
F381
Y403
L420
R428
C453
F462
V472
M477
Y478
S481
D482
A483
R486
E490
G495
I506

D510
E514
H520
L532
L540
I551
A554
T560
K566
I567
L568
E580
R581
ARG

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	71.58Å 97.30Å 99.16Å 105.15° 103.96° 100.26°	Depositor
Resolution (Å)	19.85 – 2.55 19.85 – 2.55	Depositor EDS
% Data completeness (in resolution range)	86.5 (19.85-2.55) 86.7 (19.85-2.55)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.66 (at 2.56Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.212 , 0.260 0.213 , 0.258	Depositor DCC
R_{free} test set	3441 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å^2)	34.8	Xtriage
Anisotropy	0.126	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.32 , 45.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.003 for -h,-l,-k	Xtriage
F_o,F_c correlation	0.90	EDS
Total number of atoms	18074	wwPDB-VP
Average B, all atoms (Å^2)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, Y3A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.59	0/4488	0.75	0/6090
1	B	0.58	1/4374 (0.0%)	0.74	3/5941 (0.1%)
1	C	0.60	0/4481	0.76	0/6079
1	D	0.59	0/4424	0.77	4/6011 (0.1%)
All	All	0.59	1/17767 (0.0%)	0.76	7/24121 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	324	GLU	CG-CD	5.61	1.60	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	327	ARG	NE-CZ-NH1	7.24	123.92	120.30
1	D	327	ARG	NE-CZ-NH2	-7.15	116.72	120.30
1	B	324	GLU	OE1-CD-OE2	-5.95	116.16	123.30
1	D	188	ARG	NE-CZ-NH2	-5.49	117.56	120.30
1	B	524	ASP	CB-CG-OD1	5.45	123.21	118.30
1	B	532	LEU	CA-CB-CG	-5.11	103.55	115.30
1	D	324	GLU	OE1-CD-OE2	-5.10	117.17	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4383	0	4327	30	0
1	B	4277	0	4173	28	0
1	C	4376	0	4334	27	0
1	D	4315	0	4217	39	0
2	A	23	0	16	0	0
2	C	20	0	15	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
4	A	174	0	0	2	0
4	B	154	0	0	2	0
4	C	204	0	0	3	0
4	D	142	0	0	2	0
All	All	18074	0	17082	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (118) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:31:VAL:HG23	1:D:74:VAL:O	1.73	0.89
1:D:31:VAL:CG2	1:D:74:VAL:O	2.29	0.81
1:A:13:VAL:HG21	1:B:13:VAL:HG21	1.65	0.78
1:C:174:ARG:NH1	4:C:839:HOH:O	2.17	0.78
1:C:13:VAL:HG21	1:D:13:VAL:HG21	1.68	0.76
1:B:520:HIS:ND1	1:B:532:LEU:HD22	2.01	0.75
1:D:520:HIS:ND1	1:D:532:LEU:HD22	2.04	0.72
1:D:472:VAL:HG12	1:D:506:ILE:HB	1.73	0.71
1:D:580:GLU:HB2	4:D:681:HOH:O	1.91	0.70
1:A:287:ARG:NH1	4:A:753:HOH:O	2.24	0.69
1:B:472:VAL:HG12	1:B:506:ILE:HB	1.75	0.67
1:C:529:LEU:HD23	1:D:540:LEU:HD13	1.77	0.67
1:B:520:HIS:CG	1:B:532:LEU:HD22	2.30	0.66
1:D:520:HIS:CG	1:D:532:LEU:HD22	2.32	0.65
1:D:17:GLU:O	1:D:20:ILE:HG22	1.97	0.64
1:A:529:LEU:HD23	1:B:540:LEU:HD13	1.80	0.64
1:D:31:VAL:HG22	1:D:32:ASP:N	2.14	0.63
1:A:330:ILE:HG23	1:A:351:LEU:HD11	1.80	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22:VAL:HG21	1:B:322:LEU:HD11	1.80	0.63
1:C:330:ILE:HG23	1:C:351:LEU:HD11	1.81	0.62
1:C:472:VAL:HG12	1:C:506:ILE:HB	1.83	0.60
1:C:174:ARG:HD2	4:C:769:HOH:O	2.02	0.60
1:D:554:ALA:HB1	1:D:567:ILE:HD11	1.83	0.59
1:B:520:HIS:CE1	1:B:532:LEU:HD22	2.38	0.58
1:A:472:VAL:HG12	1:A:506:ILE:HB	1.86	0.58
1:D:52:ASP:OD1	1:D:292:ARG:NH2	2.34	0.58
1:C:35:LYS:HD3	1:C:52:ASP:HB2	1.86	0.57
1:D:478:TYR:CE1	1:D:486:ARG:HG3	2.40	0.57
1:D:51:TYR:CE1	1:D:54:GLY:HA2	2.40	0.57
1:D:551:ILE:HG21	1:D:566:LYS:HD3	1.88	0.56
1:D:520:HIS:CE1	1:D:532:LEU:HD22	2.42	0.55
1:B:462:PHE:O	1:B:514:GLU:HG2	2.06	0.55
1:C:216:ARG:NH2	1:C:216:ARG:HB2	2.22	0.55
1:A:32:ASP:OD1	1:A:33:GLY:N	2.39	0.55
1:D:551:ILE:CG2	1:D:566:LYS:HD3	2.37	0.54
1:B:580:GLU:CG	1:B:580:GLU:O	2.56	0.54
1:A:52:ASP:OD1	1:A:292:ARG:NH2	2.40	0.54
1:D:462:PHE:O	1:D:514:GLU:HG2	2.08	0.53
1:C:113:ARG:NH1	4:C:782:HOH:O	2.27	0.52
1:B:580:GLU:O	1:B:580:GLU:HG3	2.08	0.52
1:B:381:PHE:CD1	1:B:568:LEU:HD13	2.45	0.52
1:C:462:PHE:O	1:C:514:GLU:HG2	2.10	0.51
1:B:551:ILE:CG2	1:B:566:LYS:HD2	2.41	0.51
1:D:23:GLU:HG2	1:D:42:SER:HB2	1.94	0.50
1:D:381:PHE:CD1	1:D:568:LEU:HD13	2.47	0.50
1:D:264[A]:ARG:NH2	1:D:267:GLY:HA2	2.27	0.50
1:B:321:GLY:C	1:B:322:LEU:HD12	2.32	0.49
1:D:31:VAL:HG22	1:D:32:ASP:H	1.76	0.49
1:D:188:ARG:NH1	1:D:189:VAL:O	2.46	0.49
1:A:86:GLY:O	1:A:555:GLY:HA3	2.12	0.49
1:C:530:LYS:HB3	1:C:531:PRO:HD3	1.94	0.48
1:D:125:VAL:HB	1:D:159:ILE:HD11	1.95	0.48
1:D:520:HIS:ND1	1:D:532:LEU:CD2	2.76	0.48
1:D:51:TYR:HE1	1:D:54:GLY:HA2	1.77	0.48
1:C:485:PHE:CZ	1:C:526:ARG:HD3	2.49	0.47
1:A:8:GLU:HG3	1:A:11[A]:ARG:HB3	1.96	0.47
1:B:110:LYS:HA	4:B:841:HOH:O	2.14	0.47
1:A:159:ILE:HG12	1:A:164:ILE:HG12	1.96	0.47
1:A:462:PHE:O	1:A:514:GLU:HG2	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:520:HIS:ND1	1:B:532:LEU:CD2	2.73	0.47
1:C:159:ILE:HG12	1:C:164:ILE:HG12	1.96	0.47
1:D:31:VAL:HG22	1:D:74:VAL:O	2.12	0.47
1:A:510[B]:ASP:OD2	1:C:510[B]:ASP:HB3	2.14	0.47
1:A:485:PHE:CZ	1:A:526:ARG:HD3	2.50	0.46
1:A:558:ILE:HG23	1:A:563:ASP:HB2	1.97	0.46
1:B:159:ILE:HG12	1:B:164:ILE:HG12	1.97	0.46
1:C:214:THR:O	1:C:215:ALA:C	2.54	0.46
1:C:13:VAL:O	1:C:17:GLU:HB2	2.15	0.46
1:D:490:GLU:HG3	1:D:495:GLY:HA2	1.96	0.46
1:D:264[A]:ARG:NH2	1:D:373:GLU:OE1	2.49	0.46
1:C:216:ARG:HB2	1:C:216:ARG:CZ	2.46	0.46
2:C:601:Y3A:C15	2:C:601:Y3A:O3	2.63	0.46
1:A:28:GLN:O	1:A:287:ARG:NH1	2.49	0.45
1:A:510[B]:ASP:OD2	1:C:510[B]:ASP:CB	2.64	0.45
1:D:477:MET:O	1:D:481:SER:OG	2.29	0.45
1:A:13:VAL:O	1:A:17:GLU:HB2	2.17	0.45
1:B:188:ARG:HG2	1:B:190:PHE:CZ	2.51	0.44
1:D:28:GLN:O	1:D:287:ARG:NH1	2.51	0.44
1:B:415:PRO:O	1:B:416:CYS:HB2	2.18	0.44
1:C:86:GLY:O	1:C:555:GLY:HA3	2.17	0.44
1:B:28:GLN:O	1:B:287:ARG:NH1	2.50	0.44
1:B:551:ILE:HG21	1:B:566:LYS:HD2	1.99	0.44
1:D:322:LEU:HD12	1:D:327:ARG:CD	2.48	0.44
1:C:71:HIS:O	1:C:74:VAL:HG23	2.17	0.43
1:C:558:ILE:HG23	1:C:563:ASP:HB2	1.99	0.43
1:A:522:GLN:HG2	1:A:552:PRO:HA	2.00	0.43
1:A:217:GLU:HB3	4:A:767:HOH:O	2.18	0.43
1:B:125:VAL:HB	1:B:159:ILE:HD11	1.99	0.43
1:A:496:SER:HB3	1:A:499:ILE:HD12	1.99	0.43
1:D:264[A]:ARG:NH1	1:D:284:ASN:OD1	2.52	0.43
1:D:43:GLU:O	1:D:560:THR:HG22	2.19	0.43
1:D:266:GLU:HA	1:D:403:TYR:CE2	2.53	0.43
1:A:555:GLY:O	1:A:558:ILE:CD1	2.67	0.43
1:B:22:VAL:O	1:B:24:LYS:NZ	2.51	0.43
1:A:530:LYS:HB3	1:A:531:PRO:HD3	2.00	0.43
1:A:214:THR:O	1:A:215:ALA:C	2.58	0.42
1:C:522:GLN:HG2	1:C:552:PRO:HA	2.01	0.42
1:A:475:GLU:OE1	1:A:501:ARG:HG3	2.19	0.42
1:A:482:ASP:O	1:A:486:ARG:HB2	2.20	0.42
1:C:420:LEU:HD22	1:C:453:CYS:SG	2.59	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:566:LYS:HE3	1:B:577:THR:HG21	2.01	0.42
1:C:125:VAL:HB	1:C:159:ILE:HD11	2.01	0.42
1:D:420:LEU:HD22	1:D:453:CYS:SG	2.60	0.42
1:A:420:LEU:HD22	1:A:453:CYS:SG	2.60	0.41
1:B:318:LEU:HD12	1:B:318:LEU:HA	1.89	0.41
1:C:28:GLN:O	1:C:287:ARG:NH1	2.53	0.41
1:B:321:GLY:O	1:B:322:LEU:HD12	2.20	0.41
1:D:235:LEU:O	1:D:238:LYS:NZ	2.42	0.41
1:B:239:ASP:HB2	1:B:275:GLY:O	2.21	0.41
1:B:6:PRO:HG2	4:B:799:HOH:O	2.21	0.41
1:A:8:GLU:HG3	1:A:11[A]:ARG:CB	2.50	0.41
1:A:239:ASP:HB2	1:A:275:GLY:O	2.20	0.41
1:A:509:VAL:HG21	1:A:538:GLU:CG	2.51	0.41
1:C:420:LEU:HA	1:C:450:MET:HE1	2.03	0.41
1:D:8:GLU:OE1	1:D:11:ARG:HD3	2.20	0.41
1:B:8:GLU:OE1	1:B:11:ARG:HD3	2.21	0.40
1:C:90:HIS:O	1:C:111:PRO:HA	2.21	0.40
1:D:428:ARG:NH2	4:D:676:HOH:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	579/582 (100%)	563 (97%)	16 (3%)	0	100	100
1	B	572/582 (98%)	552 (96%)	20 (4%)	0	100	100
1	C	579/582 (100%)	562 (97%)	17 (3%)	0	100	100
1	D	581/582 (100%)	561 (97%)	20 (3%)	0	100	100
All	All	2311/2328 (99%)	2238 (97%)	73 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	458/469 (98%)	451 (98%)	7 (2%)	65	77
1	B	437/469 (93%)	429 (98%)	8 (2%)	59	74
1	C	455/469 (97%)	447 (98%)	8 (2%)	59	74
1	D	440/469 (94%)	430 (98%)	10 (2%)	50	65
All	All	1790/1876 (95%)	1757 (98%)	33 (2%)	60	74

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	HIS
1	A	132	ASP
1	A	196	SER
1	A	287	ARG
1	A	428	ARG
1	A	501	ARG
1	A	522	GLN
1	B	34	ASP
1	B	71	HIS
1	B	196[A]	SER
1	B	196[B]	SER
1	B	241	SER
1	B	287	ARG
1	B	416	CYS
1	B	579	ARG
1	C	71	HIS
1	C	104	GLN
1	C	132	ASP
1	C	149	ARG
1	C	196	SER
1	C	244	ARG
1	C	287	ARG
1	C	522	GLN
1	D	34	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	71	HIS
1	D	196[A]	SER
1	D	196[B]	SER
1	D	241	SER
1	D	279	GLU
1	D	287	ARG
1	D	486	ARG
1	D	510[A]	ASP
1	D	510[B]	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	96	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	Y3A	C	601	1	18,20,32	0.76	0	19,25,40	0.50	0
2	Y3A	A	601	1	21,23,32	0.80	1 (4%)	24,29,40	1.23	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	Y3A	C	601	1	-	6/19/19/29	0/1/1/2
2	Y3A	A	601	1	-	10/22/22/29	0/1/1/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	Y3A	CA3-N3	2.50	1.51	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	601	Y3A	C3-CA3-N3	-3.04	104.64	113.17
2	A	601	Y3A	C13-C14-N31	2.85	121.17	113.17

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	Y3A	O4-C4-CA4-CB4
2	C	601	Y3A	O4-C4-CA4-CB4
2	A	601	Y3A	O3-C3-N4-CA4
2	A	601	Y3A	CA3-C3-N4-CA4
2	A	601	Y3A	O4-C4-CA4-N4
2	C	601	Y3A	O4-C4-CA4-N4
2	A	601	Y3A	N4-C3-CA3-N3
2	C	601	Y3A	O3-C3-N4-CA4
2	A	601	Y3A	C15-C4-CA4-N4
2	A	601	Y3A	C15-C4-CA4-CB4
2	A	601	Y3A	O3-C3-CA3-N3
2	C	601	Y3A	C15-C4-CA4-CB4
2	C	601	Y3A	CB4-CA4-N4-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	601	Y3A	N3-C13-C14-N31
2	A	601	Y3A	O31-C13-C14-N31
2	C	601	Y3A	CA3-C3-N4-CA4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	601	Y3A	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand Y3A C 601





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	577/582 (99%)	-0.31	3 (0%)	91	94	20, 30, 47, 71	0
1	B	574/582 (98%)	-0.09	13 (2%)	60	67	20, 35, 61, 96	0
1	C	577/582 (99%)	-0.26	3 (0%)	91	94	15, 31, 50, 65	0
1	D	577/582 (99%)	-0.11	14 (2%)	59	65	19, 34, 56, 81	0
All	All	2305/2328 (99%)	-0.20	33 (1%)	75	81	15, 32, 54, 96	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	555	GLY	6.6
1	B	556	HIS	5.8
1	D	87	ALA	4.4
1	A	141	GLY	4.3
1	D	85	LYS	4.0
1	D	5	MET	3.9
1	A	5	MET	3.5
1	D	132	ASP	3.5
1	D	54	GLY	3.5
1	D	131	GLU	3.3
1	B	6	PRO	3.1
1	D	142	GLY	3.0
1	B	33	GLY	3.0
1	D	141	GLY	2.7
1	A	341	PHE	2.6
1	C	194	GLU	2.5
1	B	193	GLY	2.4
1	D	173	GLY	2.4
1	D	172	GLY	2.3
1	B	132	ASP	2.3
1	D	84[A]	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	332	GLY	2.3
1	B	194	GLU	2.3
1	B	108	ALA	2.3
1	B	553	ASP	2.3
1	D	483	ALA	2.3
1	B	494	GLY	2.2
1	C	126	PHE	2.2
1	C	125	VAL	2.2
1	D	145[A]	ARG	2.1
1	B	557	ALA	2.1
1	B	498	GLU	2.1
1	B	86	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	Y3A	A	601	23/31	0.83	0.21	26,28,47,51	0
2	Y3A	C	601	20/31	0.83	0.20	24,28,47,52	0
3	CL	C	602	1/1	0.88	0.13	51,51,51,51	0
3	CL	A	602	1/1	0.92	0.13	38,38,38,38	0
3	CL	B	601	1/1	0.93	0.12	40,40,40,40	0
3	CL	B	602	1/1	0.95	0.10	35,35,35,35	0
3	CL	A	603	1/1	0.95	0.05	39,39,39,39	0
3	CL	C	603	1/1	0.95	0.15	44,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Y3A A 601: 2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)	
	

Electron density around Y3A C 601:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.