Full wwPDB X-ray Structure Validation Report (i)

Jan 16, 2024 – 03:38 am GMT

PDB ID	:	6RTH
Title	:	Crystal structure of the ligand-free glycosyltransferase domain from the YGT toxin
Authors	:	Wirth, C.; Bogdanovic, X.; Kao, W.-C.; Hunte, C.
Deposited on	:	2019-05-23
Resolution	:	3.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1026 (3.48-3.32)
Clashscore	141614	1055 (3.48-3.32)
Ramachandran outliers	138981	1038 (3.48-3.32)
Sidechain outliers	138945	1038 (3.48-3.32)
RSRZ outliers	127900	2173 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	521	86% 10% • •
1	B	521	% 85% 10% • 5%

2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 8080 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RTX toxin and Ca2+-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	504	4073	2583	680	798	12	43	0	0
1	B	496	4007	2545	667	783	12	59	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP C4SH25
A	1	SER	-	expression tag	UNP C4SH25
B	0	GLY	-	expression tag	UNP C4SH25
B	1	SER	-	expression tag	UNP C4SH25

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RTX toxin and Ca2+-binding protein

Chain A:

• Molecule 1: RTX toxin and Ca2+-binding protein

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	47.18Å 161.74Å 162.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.40 45.32 – 3.40	Depositor EDS
% Data completeness (in resolution range)	99.1 (30.00-3.40) 99.1 (45.32-3.40)	Depositor EDS
R_{merge}	0.47	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.28 (at 3.40Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.227 , 0.249 0.241 , 0.256	Depositor DCC
R_{free} test set	918 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å^2)	43.1	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.29 , -3.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.47$, $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	0.048 for -h,l,k	Xtriage
F_o,F_c correlation	0.86	EDS
Total number of atoms	8080	wwPDB-VP
Average B, all atoms (Å^2)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.42	0/4148	0.62	0/5616
1	B	0.42	0/4080	0.63	0/5519
All	All	0.42	0/8228	0.63	0/11135

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4073	0	3999	23	0
1	B	4007	0	3914	20	0
All	All	8080	0	7913	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:232:ASN:HB3	1:B:240:VAL:HG21	1.78	0.65
1:A:245:ILE:HG23	1:A:275:ILE:HD12	1.82	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:LEU:HD21	1:A:136:LYS:HD3	1.86	0.58
1:B:90:TYR:HB3	1:B:381:VAL:HG22	1.87	0.57
1:B:250:GLU:HG3	1:B:255:LEU:HD12	1.87	0.56
1:A:250:GLU:HG3	1:A:255:LEU:HD12	1.87	0.56
1:A:425:VAL:O	1:A:429:PHE:HB2	2.07	0.55
1:B:473:SER:HB3	1:B:476:GLN:HB3	1.89	0.54
1:A:176:LEU:HD12	1:A:196:ILE:HG23	1.89	0.54
1:B:361:ASN:HD22	1:B:364:HIS:H	1.56	0.53
1:A:300:ASN:HD22	1:A:300:ASN:N	2.10	0.50
1:B:69:LEU:HD13	1:B:121:ILE:HG23	1.94	0.49
1:A:69:LEU:HD13	1:A:121:ILE:HG23	1.96	0.47
1:A:90:TYR:HB3	1:A:381:VAL:HG22	1.95	0.47
1:A:467:PHE:HD1	1:A:501:LEU:HD23	1.79	0.47
1:B:222:LEU:HD21	1:B:278:LEU:HD22	1.96	0.47
1:A:161:LEU:HD11	1:A:201:ALA:HA	1.97	0.46
1:A:303:ARG:NE	1:A:303:ARG:HA	2.31	0.46
1:A:90:TYR:CE1	1:A:382:PRO:HD3	2.51	0.45
1:A:300:ASN:HD22	1:A:300:ASN:H	1.64	0.45
1:A:250:GLU:HB3	1:A:260:THR:HG21	1.98	0.45
1:B:250:GLU:HB3	1:B:260:THR:HG21	1.98	0.44
1:A:499:LEU:HD11	1:A:519:ALA:HB2	2.00	0.44
1:B:161:LEU:HD11	1:B:201:ALA:HA	1.99	0.44
1:B:67:LYS:O	1:B:71:GLU:HG3	2.18	0.43
1:A:176:LEU:HD22	1:A:327:ILE:HD11	2.01	0.43
1:B:430:SER:C	1:B:432:TRP:H	2.22	0.43
1:A:408:ALA:HB1	1:A:414:VAL:HB	2.00	0.43
1:B:122:ALA:HB1	1:B:130:VAL:HA	1.99	0.43
1:A:67:LYS:O	1:A:71:GLU:HG3	2.19	0.43
1:B:408:ALA:HB1	1:B:414:VAL:HB	2.01	0.43
1:A:48:HIS:O	1:A:154:HIS:HB3	2.18	0.43
1:A:78:LEU:HD11	1:A:89:TYR:CE1	2.54	0.43
1:A:222:LEU:HD21	1:A:278:LEU:HD22	1.99	0.43
1:B:432:TRP:HE3	1:B:432:TRP:HA	1.84	0.42
1:B:432:TRP:HA	1:B:432:TRP:CE3	2.55	0.41
1:B:34:SER:HB3	1:B:255:LEU:HD22	2.01	0.41
1:A:34:SER:HB3	1:A:255:LEU:HD22	2.03	0.41
1:B:444:TYR:O	1:B:448:VAL:HG23	2.20	0.41
1:B:463:ARG:O	1:B:467:PHE:HD2	2.04	0.41
1:A:368:PHE:HD1	1:A:392:PHE:HB3	1.86	0.41
1:B:51:TRP:CZ3	1:B:161:LEU:HB2	2.56	0.40
1:B:460:GLY:HA2	1:B:463:ARG:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	500/521 (96%)	479 (96%)	20 (4%)	1 (0%)	47	78
1	B	484/521 (93%)	464 (96%)	17 (4%)	3 (1%)	25	57
All	All	984/1042 (94%)	943 (96%)	37 (4%)	4 (0%)	34	67

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	233	ASP
1	B	435	SER
1	B	84	ALA
1	A	125	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	450/466 (97%)	431 (96%)	19 (4%)	30	59
1	B	441/466 (95%)	420 (95%)	21 (5%)	25	56
All	All	891/932 (96%)	851 (96%)	40 (4%)	27	58

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	THR
1	A	61	LEU
1	A	83	ASN
1	A	104	VAL
1	A	108	HIS
1	A	138	GLU
1	A	152	GLN
1	A	176	LEU
1	A	205	GLN
1	A	300	ASN
1	A	348	LEU
1	A	361	ASN
1	A	365	GLU
1	A	422	LEU
1	A	429	PHE
1	A	439	ASP
1	A	473	SER
1	A	504	LYS
1	A	514	THR
1	B	21	THR
1	B	61	LEU
1	B	85	THR
1	B	87	ASN
1	B	88	ASP
1	B	104	VAL
1	B	125	LEU
1	B	149	ASP
1	B	176	LEU
1	B	202	LEU
1	B	232	ASN
1	B	277	ARG
1	B	422	LEU
1	B	429	PHE
1	B	432	TRP
1	B	434	ILE
1	B	435	SER
1	B	456	SER
1	B	478	THR
1	B	490	SER
1	B	514	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	164	GLN
1	A	300	ASN
1	A	361	ASN
1	A	476	GLN
1	B	164	GLN
1	B	205	GLN
1	B	361	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	504/521 (96%)	-0.13	0	100	100	17, 41, 66, 88	11 (2%)
1	B	496/521 (95%)	-0.01	4 (0%)	86	85	13, 44, 84, 105	14 (2%)
All	All	1000/1042 (95%)	-0.07	4 (0%)	92	92	13, 42, 77, 105	25 (2%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	475	GLU	2.4
1	B	430	SER	2.3
1	B	232	ASN	2.2
1	B	231	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.