wwPDB X-ray Structure Validation Summary Report (i)

Feb 17, 2024 – 03:00 PM EST

PDB ID	:	3S8G
Title	:	1.8 A structure of ba3 cytochrome c oxidase mutant (A120F) from Thermus thermophilus in lipid environment
Authors	:	Tiefenbrunn, T.; Liu, W.; Chen, Y.; Katritch, V.; Stout, C.D.; Fee, J.A.; Cherezov, V.
Deposited on	:	2011-05-27
Resolution	:	1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	569	4% 90% 7% • •
2	B	168	2% 92% 7% ••
3	C	34	76% 9% 6% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	HAS	A	801	X	-	-	-

2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 6695 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c oxidase subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	554	4399	2988	700	694	17	0	7	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	expression tag	UNP Q5SJ79
A	-5	HIS	-	expression tag	UNP Q5SJ79
A	-4	HIS	-	expression tag	UNP Q5SJ79
A	-3	HIS	-	expression tag	UNP Q5SJ79
A	-2	HIS	-	expression tag	UNP Q5SJ79
A	-1	HIS	-	expression tag	UNP Q5SJ79
A	0	HIS	-	expression tag	UNP Q5SJ79
A	1	HIS	-	expression tag	UNP Q5SJ79
A	120	PHE	ALA	conflict	UNP Q5SJ79

- Molecule 2 is a protein called Cytochrome c oxidase subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	166	1307	846	219	238	4	0	5	0

- Molecule 3 is a protein called Cytochrome c oxidase polypeptide 2A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	31	241	169	37	35	0	0	0

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
4	A	1	43	34	1	4	4	0	0

- Molecule 5 is HEME-AS (three-letter code: HAS) (formula: $C_{54}H_{64}FeN_4O_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
5	A	1	65	54	1	4	6	0	0

- Molecule 6 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cu		
6	A	1	1	1	0	0

- Molecule 7 is PEROXIDE ION (three-letter code: PER) (formula: O_2).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	1	2	2	0	0

- Molecule 8 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: $C_{21}H_{40}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
8	A	1	25	21	4	0	0
8	A	1	25	21	4	0	0
8	A	1	25	21	4	0	0
8	A	1	23	19	4	0	0
8	A	1	25	21	4	0	0
8	A	1	18	14	4	0	0
8	A	1	17	13	4	0	0
8	A	1	8	4	4	0	0
8	A	1	15	11	4	0	0
8	A	1	20	16	4	0	0
8	A	1	25	21	4	0	0
8	A	1	21	17	4	0	0
8	A	1	16	12	4	0	0
8	A	1	25	21	4	0	0
8	A	1	12	8	4	0	0
8	B	1	25	21	4	0	0
8	B	1	25	21	4	0	0
8	B	1	25	21	4	0	0
8	B	1	21	19	2	0	0
8	C	1	14	10	4	0	0

- Molecule 9 is DINUCLEAR COPPER ION (three-letter code: CUA) (formula: Cu_2).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cu		
9	B	1	2	2	0	0

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
10	A	137	137	137	0	0
			Total	O		
10	B	81	81	81	0	0
			Total	O		
10	C	7	7	7	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cytochrome c oxidase subunit 1

Chain A:



- Molecule 2: Cytochrome c oxidase subunit 2

Chain B:





- Molecule 3: Cytochrome c oxidase polypeptide 2A

Chain C:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	144.96Å 98.64Å 95.06Å 90.00° 128.07° 90.00°	Depositor
Resolution (Å)	39.48 – 1.80 39.48 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.8 (39.48-1.80) 97.8 (39.48-1.80)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.16 (at 1.79Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.175 , 0.196 0.175 , 0.197	Depositor DCC
R_{free} test set	4759 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å^2)	21.3	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 66.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.017 for -h+k-l,-l,-k 0.002 for -h-k-l,l,k 0.018 for -h-2*l,-k,l	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	6695	wwPDB-VP
Average B, all atoms (Å^2)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: OLC, CUA, PER, HAS, HEM, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.68	0/4584	0.64	1/6292 (0.0%)
2	B	0.69	0/1364	0.61	0/1862
3	C	0.73	0/247	0.72	0/335
All	All	0.69	0/6195	0.64	1/8489 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	233	HIS	N-CA-CB	5.24	120.04	110.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4399	0	4496	21	0
2	B	1307	0	1277	9	0
3	C	241	0	267	4	0
4	A	43	0	30	3	0
5	A	65	0	62	1	0
6	A	1	0	0	0	0
7	A	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	A	300	0	434	8	0
8	B	96	0	153	7	0
8	C	14	0	17	0	0
9	B	2	0	0	0	0
10	A	137	0	0	2	0
10	B	81	0	0	1	1
10	C	7	0	0	2	0
All	All	6695	0	6736	39	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 39 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:141:ARG:HH12	8:B:170:OLC:H24A	1.19	1.05
1:A:168:ARG:HH22	8:A:572:OLC:H6A	1.26	0.97
2:B:158[B]:GLN:HG2	2:B:159[B]:ASN:OD1	1.80	0.80
3:C:33:ARG:HD2	10:C:36:HOH:O	1.89	0.72
1:A:168:ARG:NH2	8:A:572:OLC:H6A	2.02	0.72

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:B:1062:HOH:O	10:B:1063:HOH:O[2_556]	1.54	0.66

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	558/569 (98%)	542 (97%)	15 (3%)	1 (0%)	47	33
2	B	169/168 (101%)	167 (99%)	2 (1%)	0	100	100
3	C	29/34 (85%)	29 (100%)	0	0	100	100
All	All	756/771 (98%)	738 (98%)	17 (2%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	516	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	453/464 (98%)	437 (96%)	16 (4%)	36	21
2	B	138/138 (100%)	133 (96%)	5 (4%)	35	20
3	C	24/27 (89%)	20 (83%)	4 (17%)	2	0
All	All	615/629 (98%)	590 (96%)	25 (4%)	31	16

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	519	ARG
2	B	37	LEU
3	C	24	LEU
2	B	26	LEU
2	B	158[A]	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	407	ASN
2	B	8	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	40	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 25 ligands modelled in this entry, 1 is monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
8	OLC	A	572	-	14,14,24	0.59	0	15,15,25	0.51	0
8	OLC	A	565	-	24,24,24	0.44	0	25,25,25	0.55	0
9	CUA	B	802	2	0,1,1	-	-	-		
8	OLC	B	169	-	24,24,24	0.46	0	25,25,25	0.62	0
5	HAS	A	801	1,7	69,72,72	2.08	22 (31%)	73,109,109	1.47	13 (17%)
8	OLC	C	35	-	13,13,24	0.60	0	14,14,25	0.64	0
8	OLC	A	567	-	22,22,24	0.47	0	23,23,25	0.72	0
8	OLC	A	568	-	24,24,24	0.46	0	25,25,25	0.55	0
8	OLC	A	573	-	19,19,24	0.51	0	20,20,25	0.70	0
8	OLC	A	570	-	16,16,24	0.51	0	17,17,25	0.60	0
8	OLC	A	569	-	17,17,24	0.57	0	18,18,25	0.63	0
8	OLC	B	171	-	24,24,24	0.46	0	25,25,25	0.52	0
8	OLC	A	566	-	24,24,24	0.45	0	25,25,25	0.59	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
8	OLC	A	578	-	11,11,24	0.62	0	12,12,25	0.56	0
8	OLC	A	576	-	15,15,24	0.55	0	16,16,25	0.60	0
8	OLC	A	575	-	20,20,24	0.47	0	21,21,25	0.69	0
8	OLC	B	172	-	20,20,24	0.47	0	20,20,25	0.60	0
7	PER	A	563	6,5	0,1,1	-	-	-		
8	OLC	A	577	-	24,24,24	0.43	0	25,25,25	0.62	0
8	OLC	A	564	-	24,24,24	0.44	0	25,25,25	0.65	0
4	HEM	A	800	1	41,50,50	2.52	17 (41%)	45,82,82	2.36	15 (33%)
8	OLC	A	574	-	24,24,24	0.41	0	25,25,25	0.69	0
8	OLC	A	571	-	7,7,24	0.44	0	6,7,25	0.45	0
8	OLC	B	170	-	24,24,24	0.53	0	25,25,25	0.63	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	OLC	A	572	-	-	7/14/14/24	-
8	OLC	A	565	-	-	13/24/24/24	-
8	OLC	B	169	-	-	9/24/24/24	-
5	HAS	A	801	1,7	1/1/8/18	4/40/82/82	-
8	OLC	C	35	-	-	8/13/13/24	-
8	OLC	A	567	-	-	8/22/22/24	-
8	OLC	A	568	-	-	13/24/24/24	-
8	OLC	A	573	-	-	10/19/19/24	-
8	OLC	A	570	-	-	4/16/16/24	-
8	OLC	A	569	-	-	6/17/17/24	-
8	OLC	B	171	-	-	14/24/24/24	-
8	OLC	A	566	-	-	10/24/24/24	-
8	OLC	A	578	-	-	5/11/11/24	-
8	OLC	A	576	-	-	8/15/15/24	-
8	OLC	A	575	-	-	8/20/20/24	-
8	OLC	B	172	-	-	9/19/19/24	-
8	OLC	A	577	-	-	8/24/24/24	-
8	OLC	A	564	-	-	15/24/24/24	-
4	HEM	A	800	1	-	3/12/54/54	-
8	OLC	A	574	-	-	15/24/24/24	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	OLC	A	571	-	-	2/6/6/24	-
8	OLC	B	170	-	-	11/24/24/24	-

The worst 5 of 39 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	801	HAS	C3C-C2C	6.37	1.49	1.40
5	A	801	HAS	C3C-CAC	-6.00	1.35	1.47
4	A	800	HEM	C3B-C2B	5.45	1.48	1.37
5	A	801	HAS	CHB-C1D	5.18	1.49	1.38
4	A	800	HEM	CHB-C1B	5.07	1.48	1.35

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	A	800	HEM	C1B-NB-C4B	6.63	111.92	105.07
4	A	800	HEM	CHC-C4B-NB	6.50	131.50	124.43
4	A	800	HEM	CHA-C4D-ND	5.06	130.64	124.38
4	A	800	HEM	C4D-ND-C1D	4.98	110.22	105.07
4	A	800	HEM	CHD-C1D-ND	4.69	129.52	124.43

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	801	HAS	NA

5 of 190 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	800	HEM	C2B-C3B-CAB-CBB
8	A	565	OLC	O20-C21-C22-C24
8	A	567	OLC	C21-C22-C24-O25
8	A	567	OLC	O23-C22-C24-O25
8	A	571	OLC	O19-C1-O20-C21

There are no ring outliers.

12 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	572	OLC	2	0
8	B	169	OLC	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	801	HAS	1	0
8	A	567	OLC	1	0
8	B	171	OLC	1	0
8	A	566	OLC	1	0
8	A	578	OLC	1	0
8	A	576	OLC	1	0
8	A	575	OLC	1	0
4	A	800	HEM	3	0
8	A	574	OLC	1	0
8	B	170	OLC	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand OLC A 572



Ligand OLC A 565



Ligand OLC B 169



Ligand HAS A 801



Ligand OLC C 35



Ligand OLC A 567



Ligand OLC A 568



Ligand OLC A 573

Ligand OLC A 570



Ligand OLC A 569



Ligand OLC B 171



Ligand OLC A 566



Ligand OLC A 578



Ligand OLC A 576



Ligand OLC A 575

Ligand OLC B 172



Ligand OLC A 577



Ligand OLC A 564



Ligand HEM A 800



Bond lengths



Bond angles



Torsions



Rings

Ligand OLC A 574



Bond lengths



Bond angles



Torsions

Rings

Ligand OLC A 571



Ligand OLC B 170



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	554/569 (97%)	-0.22	23 (4%) 36 30	11, 21, 43, 67	0
2	B	166/168 (98%)	-0.39	4 (2%) 59 54	12, 22, 38, 53	0
3	C	31/34 (91%)	-0.70	0 100 100	16, 21, 31, 41	0
All	All	751/771 (97%)	-0.28	27 (3%) 42 37	11, 21, 42, 67	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	495	ARG	5.3
1	A	515	PRO	5.0
1	A	333	PHE	4.9
1	A	496	GLU	4.4
1	A	516	GLU	4.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
8	OLC	A	569	18/25	0.41	0.23	56,61,70,70	0
8	OLC	A	578	12/25	0.44	0.31	74,76,77,78	0
8	OLC	A	568	25/25	0.47	0.30	74,75,83,83	0
8	OLC	A	570	17/25	0.52	0.26	63,65,74,74	0
8	OLC	A	565	25/25	0.57	0.25	66,70,79,80	0
8	OLC	A	573	20/25	0.58	0.24	59,63,64,64	0
8	OLC	A	566	25/25	0.58	0.22	54,64,73,73	0
8	OLC	B	170	25/25	0.58	0.23	60,65,68,69	0
8	OLC	A	576	16/25	0.61	0.30	67,70,72,72	0
8	OLC	C	35	14/25	0.62	0.28	77,78,79,79	0
8	OLC	A	572	15/25	0.63	0.38	54,59,65,65	0
8	OLC	A	577	25/25	0.65	0.22	52,55,58,59	0
8	OLC	A	575	21/25	0.66	0.20	57,63,71,72	0
8	OLC	B	172	21/25	0.68	0.23	59,61,65,66	0
8	OLC	A	564	25/25	0.71	0.21	37,46,64,65	0
8	OLC	A	574	25/25	0.74	0.18	53,56,57,57	0
8	OLC	B	171	25/25	0.76	0.21	59,61,66,67	0
8	OLC	B	169	25/25	0.77	0.18	56,59,61,62	0
8	OLC	A	571	8/25	0.84	0.25	59,61,63,64	0
8	OLC	A	567	23/25	0.89	0.14	28,36,54,55	0
7	PER	A	563	2/2	0.97	0.12	14,14,14,15	0
5	HAS	A	801	65/65	0.99	0.11	8,13,26,30	0
4	HEM	A	800	43/43	0.99	0.08	6,11,14,20	0
6	CU	A	803	1/1	1.00	0.07	15,15,15,15	0
9	CUA	B	802	2/2	1.00	0.06	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OLC A 569:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around OLC A 578:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around OLC A 568:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OLC A 570:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OLC A 565:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OLC A 573:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OLC A 566:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OLC B 170:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OLC A 576:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OLC C 35:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OLC A 572:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around OLC A 577:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around OLC A 575:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around OLC B 172:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around OLC A 564:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OLC A 574:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OLC B 171:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around OLC B 169:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around OLC A 571:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around OLC A 567:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around HAS A 801:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 800:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.