



Full wwPDB X-ray Structure Validation Report i

May 21, 2020 – 08:32 am BST

PDB ID : 6SH7
Title : Crystal structure of the human DEAH-helicase DHX15 in complex with the NKRF G-patch
Authors : Jonas, S.; Studer, M.K.; Ivanovic, L.
Deposited on : 2019-08-05
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see references ①) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5912 (2.24-2.20)
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)
RSRZ outliers	127900	5797 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 5745 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor ATP-dependent RNA helicase DHX15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	665	5362	3406	927	992	37	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	107	GLY	-	expression tag	UNP O43143
A	108	PRO	-	expression tag	UNP O43143
A	109	HIS	-	expression tag	UNP O43143
A	110	MET	-	expression tag	UNP O43143
A	111	LEU	-	expression tag	UNP O43143
A	112	GLU	-	expression tag	UNP O43143

- Molecule 2 is a protein called NF-kappa-B-repressing factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	46	349	213	66	69	1	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	537	GLY	-	expression tag	UNP O15226
B	538	PRO	-	expression tag	UNP O15226
B	539	HIS	-	expression tag	UNP O15226
B	540	MET	-	expression tag	UNP O15226


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	34	Total O 34 34	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-splicing factor ATP-dependent RNA helicase DHX15



- Molecule 2: NF-kappa-B-repressing factor



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	82.24Å 90.02Å 213.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.23 – 2.21 46.23 – 2.21	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.23-2.21) 99.8 (46.23-2.21)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.00 (at 2.22Å)	Xtriage
Refinement program	PHENIX	Depositor
R , R_{free}	0.209 , 0.251 0.209 , 0.251	Depositor DCC
R_{free} test set	1983 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	61.8	Xtriage
Anisotropy	0.186	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 50.8	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5745	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.24	0/5472	0.40	0/7413
2	B	0.24	0/352	0.46	0/467
All	All	0.24	0/5824	0.41	0/7880

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5362	0	5386	30	0
2	B	349	0	348	1	0
3	A	34	0	0	0	0
All	All	5745	0	5734	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:392:LEU:HD12	1:A:393:PRO:HD2	1.75	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:742:THR:OG1	1:A:743:THR:N	2.30	0.65
1:A:384:LYS:HG2	1:A:405:PRO:HD3	1.80	0.62
1:A:256:VAL:HG22	1:A:287:LYS:HB2	1.89	0.54
1:A:768:MET:O	1:A:777:LYS:NZ	2.31	0.54
1:A:193:PRO:HB3	1:A:266:THR:HG21	1.91	0.51
1:A:446:VAL:HG21	1:A:457:LEU:HD12	1.92	0.51
1:A:633:ALA:HA	1:A:636:GLN:HG2	1.93	0.50
1:A:389:TYR:CZ	1:A:392:LEU:HD13	2.46	0.50
1:A:564:GLU:HG3	1:A:671:PHE:HZ	1.77	0.50
1:A:384:LYS:HE2	1:A:405:PRO:HB3	1.94	0.49
1:A:343:HIS:HD2	1:A:415:ARG:HD2	1.76	0.49
1:A:356:LEU:HD13	1:A:361:GLU:HG2	1.94	0.49
1:A:130:ILE:HG21	1:A:208:GLU:HG2	1.94	0.48
1:A:771:PHE:HB3	1:A:777:LYS:HD2	1.94	0.48
2:B:554:ILE:HD12	2:B:554:ILE:H	1.79	0.47
1:A:385:ILE:HA	1:A:417:VAL:HG13	1.97	0.47
1:A:349:GLU:O	1:A:477:ARG:NH1	2.30	0.46
1:A:192:GLN:O	1:A:237:THR:HA	2.16	0.46
1:A:705:GLU:OE1	1:A:707:THR:OG1	2.31	0.45
1:A:514:LEU:HD22	1:A:519:ILE:HD12	1.99	0.45
1:A:259:LEU:HD11	1:A:276:LEU:HD12	2.00	0.43
1:A:389:TYR:O	1:A:397:GLN:NE2	2.51	0.42
1:A:399:ARG:O	1:A:402:GLU:HG2	2.18	0.42
1:A:704:LEU:HB3	1:A:732:GLU:HA	2.00	0.42
1:A:167:THR:HG22	1:A:201:VAL:HG21	2.01	0.42
1:A:564:GLU:HG3	1:A:671:PHE:CZ	2.55	0.42
1:A:585:ASN:OD1	1:A:675:ARG:HB3	2.18	0.42
1:A:719:VAL:HG12	1:A:745:ASN:HB3	2.02	0.41
1:A:544:ASN:OD1	1:A:764:GLN:NE2	2.46	0.40
1:A:600:PHE:CE1	1:A:614:LYS:HG3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	657/689 (95%)	635 (97%)	22 (3%)	0	100	100
2	B	44/67 (66%)	42 (96%)	2 (4%)	0	100	100
All	All	701/756 (93%)	677 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	596/615 (97%)	593 (100%)	3 (0%)	88	94
2	B	36/53 (68%)	36 (100%)	0	100	100
All	All	632/668 (95%)	629 (100%)	3 (0%)	88	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	210	ASP
1	A	498	THR
1	A	742	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	113	GLN
1	A	341	GLN
1	A	343	HIS
1	A	395	GLN
1	A	444	GLN
1	A	495	GLN
1	A	700	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	709	HIS
1	A	788	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	665/689 (96%)	0.25	36 (5%) 25 24	46, 71, 132, 176	0
2	B	46/67 (68%)	1.22	11 (23%) 0 0	78, 106, 147, 167	0
All	All	711/756 (94%)	0.31	47 (6%) 18 17	46, 73, 134, 176	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	412	ALA	6.6
1	A	413	ILE	6.3
1	A	403	PRO	5.0
2	B	595	VAL	4.6
2	B	584	GLU	4.4
1	A	401	PHE	4.2
2	B	586	HIS	4.2
2	B	572	SER	4.0
1	A	384	LYS	3.9
1	A	774	CYS	3.5
1	A	492	THR	3.3
1	A	488	LYS	3.3
1	A	374	ASP	3.2
1	A	373	VAL	3.2
1	A	121	LEU	2.9
1	A	366	CYS	2.9
1	A	120	ASN	2.9
2	B	565	THR	2.8
1	A	185	LYS	2.8
2	B	577	ARG	2.7
1	A	329	ARG	2.7
1	A	470	ALA	2.7
1	A	128	TYR	2.6
1	A	370	LYS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	131	LEU	2.5
1	A	400	ILE	2.4
1	A	773	GLN	2.4
1	A	415	ARG	2.4
2	B	583	LYS	2.4
2	B	594	ASP	2.4
1	A	371	ARG	2.4
1	A	416	LYS	2.3
2	B	596	GLU	2.3
1	A	386	ILE	2.3
1	A	375	ASP	2.3
2	B	580	ILE	2.3
1	A	493	GLU	2.3
1	A	295	LEU	2.2
1	A	405	PRO	2.2
1	A	114	CYS	2.2
1	A	414	GLY	2.1
1	A	395	GLN	2.1
1	A	182	PRO	2.1
2	B	576	ILE	2.1
1	A	782	ARG	2.1
1	A	126	ARG	2.0
1	A	433	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.