Full wwPDB X-ray Structure Validation Report (i)

Sep 3, 2023 – 06:41 PM EDT

PDB ID	:	3SQR
Title	:	Crystal structure of laccase from Botrytis aclada at 1.67 A resolution
Authors	:	Osipov, E.M.; Polyakov, K.M.; Tikhonova, T.V.; Dorovatovsky, P.V.; Ludwig, R.; Kittl, R.; Shleev, S.V.; Popov, V.O.
Deposited on	:	2011-07-06
Resolution	:	1.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	580	4% 87% 6% 7%
2	B	3	100%
3	C	6	100%

2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 4851 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called laccase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	539	4206	2677	687	829	13	0	6	0

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	3	39	22	2	15	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	6	72	40	2	30	0	0	0

- Molecule 4 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cu		
4	A	3	3	3	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
			Total	C	N	O		
5	A	1	14	8	1	5	0	0

- Molecule 6 is alpha-D-mannopyranose (three-letter code: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	A	1	11	6	5	0	0

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
7	A	1	5	4	1	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
8	A	445	445	445	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: laccase

Chain A:



• Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:



• Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	69.67Å 113.50Å 79.91Å 90.00° 108.75° 90.00°	Depositor
Resolution (Å)	75.67 – 1.67 28.13 – 1.67	Depositor EDS
% Data completeness (in resolution range)	97.0 (75.67-1.67) 97.0 (28.13-1.67)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$< I/\sigma(I) >$ [1]	1.68 (at 1.68Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.165 , 0.191 0.175 , 0.202	Depositor DCC
R_{free} test set	3329 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å^2)	21.5	Xtriage
Anisotropy	0.529	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.39 , 45.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	4851	wwPDB-VP
Average B, all atoms (Å^2)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CU, NAG, BMA, SO4, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.01	2/4362 (0.0%)	0.92	1/5988 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	397	TYR	CD1-CE1	-5.64	1.30	1.39
1	A	240	GLU	CD-OE2	5.31	1.31	1.25

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	159	ASP	CB-CG-OD2	-5.66	113.21	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4206	0	3905	22	0
2	B	39	0	34	0	0
3	C	72	0	61	0	0
4	A	3	0	0	0	0
5	A	70	0	65	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	11	0	10	0	0
7	A	5	0	0	0	0
8	A	445	0	0	4	0
All	All	4851	0	4075	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:404:GLN:NE2	8:A:1004:HOH:O	2.13	0.81
1:A:34[A]:TRP:CZ3	8:A:845:HOH:O	2.46	0.67
1:A:300:ASN:ND2	1:A:302:GLU:H	1.96	0.64
1:A:34[A]:TRP:HZ3	8:A:845:HOH:O	1.80	0.62
1:A:136:LEU:HD13	1:A:173:VAL:HG22	1.82	0.62
1:A:156:TYR:HB3	1:A:277:ILE:HD12	1.87	0.57
1:A:332:GLU:HG2	1:A:425:TRP:CH2	2.47	0.50
1:A:97:LEU:HD21	1:A:383[B]:THR:CG2	2.43	0.48
1:A:388:ASN:HA	5:A:559:NAG:H82	1.96	0.47
1:A:205:THR:HG23	1:A:205:THR:O	2.15	0.47
1:A:345:LEU:O	1:A:458:ASN:HA	2.16	0.46
1:A:234:GLY:HA3	1:A:237:SER:OG	2.17	0.44
1:A:173:VAL:HG23	1:A:493:TRP:CH2	2.52	0.44
1:A:205:THR:O	1:A:205:THR:CG2	2.65	0.44
1:A:160:VAL:HG22	1:A:227:ARG:O	2.19	0.43
1:A:300:ASN:HD22	1:A:302:GLU:H	1.63	0.42
1:A:147:ILE:HD11	1:A:162:VAL:HG23	2.02	0.42
1:A:34[A]:TRP:CH2	8:A:845:HOH:O	2.71	0.41
1:A:162:VAL:HA	1:A:229:ARG:O	2.20	0.41
1:A:217:GLU:HA	1:A:309:ILE:O	2.19	0.41
1:A:250:VAL:O	1:A:260:PRO:HA	2.21	0.40
1:A:435:PHE:O	1:A:461:ARG:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	541/580 (93%)	524 (97%)	17 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	459/488 (94%)	458 (100%)	1 (0%)	93 89

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	205	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	166	GLN
1	A	253	ASN
1	A	300	ASN
1	A	319	ASN
1	A	455	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	NAG	B	1	1,2	14,14,15	1.03	1 (7%)	17,19,21	1.86	5 (29%)
2	NAG	B	2	2	14,14,15	1.02	1 (7%)	17,19,21	1.02	0
2	BMA	B	3	2	11,11,12	0.80	0	15,15,17	3.18	6 (40%)
3	NAG	C	1	1,3	14,14,15	1.10	1 (7%)	17,19,21	1.71	2 (11%)
3	NAG	C	2	3	14,14,15	0.38	0	17,19,21	1.17	3 (17%)
3	BMA	C	3	3	11,11,12	0.89	1 (9%)	15,15,17	1.47	3 (20%)
3	MAN	C	4	3	11,11,12	0.81	0	15,15,17	1.33	2 (13%)
3	MAN	C	5	3	11,11,12	0.70	0	15,15,17	1.63	4 (26%)
3	MAN	C	6	3	11,11,12	0.75	0	15,15,17	1.47	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	1/2/19/22	0/1/1/1
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	0/2/19/22	0/1/1/1
3	MAN	C	5	3	-	0/2/19/22	0/1/1/1
3	MAN	C	6	3	-	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NAG	O5-C1	-2.89	1.39	1.43
3	C	1	NAG	O5-C1	-2.76	1.39	1.43
2	B	2	NAG	O5-C1	-2.53	1.39	1.43
3	C	3	BMA	C2-C3	2.06	1.55	1.52

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	3	BMA	C1-O5-C5	9.79	125.45	112.19
3	C	1	NAG	O5-C5-C6	-5.20	99.06	107.20
2	B	1	NAG	C1-O5-C5	4.18	117.85	112.19
2	B	3	BMA	O6-C6-C5	3.94	124.82	111.29
3	C	5	MAN	C1-O5-C5	3.65	117.14	112.19
3	C	6	MAN	C1-O5-C5	3.56	117.01	112.19
3	C	3	BMA	O3-C3-C2	3.34	116.39	109.99
3	C	6	MAN	O4-C4-C5	3.15	117.12	109.30
3	C	1	NAG	C1-O5-C5	3.11	116.40	112.19
3	C	3	BMA	O2-C2-C3	3.04	116.23	110.14
2	B	3	BMA	O5-C5-C6	-3.03	102.45	107.20
3	C	5	MAN	O3-C3-C2	-2.83	104.57	109.99
3	C	2	NAG	O5-C5-C6	2.71	111.45	107.20
2	B	1	NAG	O7-C7-N2	2.68	126.89	121.95
2	B	1	NAG	C4-C3-C2	-2.65	107.13	111.02
2	B	1	NAG	O5-C5-C6	-2.57	103.17	107.20
3	C	3	BMA	O2-C2-C1	-2.55	103.93	109.15
2	B	3	BMA	C3-C4-C5	2.54	114.77	110.24
2	B	3	BMA	C2-C3-C4	2.49	115.21	110.89
3	C	5	MAN	O2-C2-C3	-2.27	105.60	110.14
2	B	1	NAG	O5-C1-C2	-2.23	107.77	111.29
2	B	3	BMA	C1-C2-C3	-2.22	106.93	109.67
3	C	2	NAG	C2-N2-C7	-2.20	119.77	122.90
3	C	5	MAN	O2-C2-C1	2.08	113.42	109.15
3	C	4	MAN	O2-C2-C1	-2.06	104.93	109.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	C	4	MAN	O3-C3-C2	2.05	113.91	109.99
3	C	2	NAG	O4-C4-C3	-2.02	105.68	110.35

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	2	NAG	C4-C5-C6-O6
2	B	3	BMA	O5-C5-C6-O6
3	C	2	NAG	O5-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

Oligosaccharide Chain B



Oligosaccharide Chain C



5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	NAG	A	560	1	14,14,15	1.03	1 (7%)	17,19,21	1.54	2 (11%)
6	MAN	A	561	1	11,11,12	0.52	0	15,15,17	1.06	1 (6%)
5	NAG	A	548	1	14,14,15	0.75	0	17,19,21	1.39	1 (5%)
7	SO4	A	562	-	4,4,4	0.17	0	6,6,6	0.24	0
5	NAG	A	558	1	14,14,15	0.61	0	17,19,21	1.32	2 (11%)
5	NAG	A	547	1	14,14,15	0.93	1 (7%)	17,19,21	1.14	1 (5%)
5	NAG	A	559	1	14,14,15	0.52	0	17,19,21	1.13	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	560	1	-	0/6/23/26	0/1/1/1
6	MAN	A	561	1	-	0/2/19/22	0/1/1/1
5	NAG	A	548	1	-	0/6/23/26	0/1/1/1
5	NAG	A	558	1	-	0/6/23/26	0/1/1/1
5	NAG	A	547	1	-	0/6/23/26	0/1/1/1
5	NAG	A	559	1	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	547	NAG	O3-C3	-2.07	1.38	1.43
5	A	560	NAG	O5-C5	-2.01	1.39	1.43

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	A	548	NAG	O5-C1-C2	-3.28	106.10	111.29
5	A	560	NAG	C4-C3-C2	-2.99	106.64	111.02
5	A	560	NAG	C1-O5-C5	2.80	115.98	112.19
5	A	559	NAG	C1-C2-N2	-2.59	106.07	110.49
5	A	547	NAG	C1-C2-N2	-2.40	106.39	110.49
5	A	558	NAG	C1-C2-N2	2.29	114.40	110.49
5	A	558	NAG	O5-C5-C4	-2.25	105.36	110.83
6	A	561	MAN	C1-O5-C5	2.14	115.09	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	559	NAG	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	539/580 (92%)	0.03	21 (3%) 39 42	19, 27, 40, 52	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	SER	4.5
1	A	203	ALA	4.5
1	A	2	CYS	4.3
1	A	317	ILE	4.1
1	A	205	THR	4.0
1	A	535	LEU	3.8
1	A	323	VAL	3.3
1	A	202	SER	3.3
1	A	512	VAL	3.2
1	A	404	GLN	3.1
1	A	363	ASN	2.9
1	A	3	ALA	2.9
1	A	321	THR	2.8
1	A	533	THR	2.7
1	A	318	ALA	2.5
1	A	288	TRP	2.3
1	A	24	VAL	2.3
1	A	207	PRO	2.2
1	A	387	PHE	2.2
1	A	204	SER	2.1
1	A	315	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	MAN	C	4	11/12	0.81	0.22	35,44,53,53	0
3	NAG	C	2	14/15	0.82	0.15	37,41,45,49	0
2	BMA	B	3	11/12	0.82	0.20	38,44,47,52	0
3	BMA	C	3	11/12	0.85	0.22	50,53,55,57	0
3	MAN	C	5	11/12	0.93	0.17	34,37,41,45	0
3	MAN	C	6	11/12	0.94	0.16	35,37,39,41	0
3	NAG	C	1	14/15	0.95	0.07	27,32,41,47	0
2	NAG	B	2	14/15	0.95	0.14	29,32,37,39	0
2	NAG	B	1	14/15	0.96	0.07	25,29,30,31	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around Chain C:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	NAG	A	560	14/15	0.84	0.19	40,47,56,56	0
5	NAG	A	559	14/15	0.86	0.25	50,54,55,56	0
5	NAG	A	548	14/15	0.94	0.08	25,30,40,40	0
6	MAN	A	561	11/12	0.94	0.09	29,32,34,34	0
7	SO4	A	562	5/5	0.95	0.23	36,37,41,43	5
5	NAG	A	558	14/15	0.96	0.12	32,36,39,39	0
5	NAG	A	547	14/15	0.98	0.05	22,24,25,25	0
4	CU	A	545	1/1	1.00	0.10	27,27,27,27	1
4	CU	A	546	1/1	1.00	0.07	27,27,27,27	1
4	CU	A	544	1/1	1.00	0.05	25,25,25,25	0

6.5 Other polymers (i)

There are no such residues in this entry.