wwPDB NMR Structure Validation Summary Report (i)

Feb 20, 2022 – 01:19 AM EST

PDB ID	:	1SS6
Title	:	Solution structure of SEP domain from human p47
Authors	:	Soukenik, M.; Leidert, M.; Sievert, V.; Buessow, K.; Leitner, D.; Labudde, D.; Ball, L.J.; Oschkinat, H.
Deposited on	:	2004-03-23

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.26
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	102	28% 25% • 45%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 9 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:14-A:55, A:63-A:76 (56)	0.34	9

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 5 single-model clusters were found.

Cluster number	Models
1	5, 8, 9, 10, 14, 20
2	4, 7, 13
3	2, 3
4	11, 19
5	15, 16
Single-model clusters	1; 6; 12; 17; 18

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1583 atoms, of which 782 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called NSFL1 cofactor p47.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	102	1583	494	782	149	157	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	cloning artifact	UNP Q9UNZ2
A	2	SER	-	cloning artifact	UNP Q9UNZ2

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: NSFL1 cofactor p47

Chain A: 28% 25% • 45%

G1 S2 E3 K4 R5 Q6 H7 S8 S9 Q10 D11 V12 H13 V14 V15 L16 K17 L18 W19 K20 S21 G22 F23 S24 L25 L30 R31 S32 D35 N38 A39 Q40 F41 L42 E43 R47 E53 L54 R55 R56 L57 A58 H59 G60 G61 Q62 V63 N64 L65 R71 D72 E73 D74 F75

V76 K77 P78 K79 G80 A81 F82 K83 A84 F85 T86 G87 E88 G89 Q90 K91 L92 G93 S94 T95 A96 P97 Q98 V99 L100 S101 T102

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 9. Colouring as in section 4.1 above.

- Molecule 1: NSFL1 cofactor p47

Chain A: 26% 26% • 45%

G1 S2 E3 K4 R5 Q6 H7 S8 S9 Q10 D11 V12 H13 V14 V15 L16 K17 L18 W19 K20 S21 G22 F23 S24 L25 D26 L30 R31 S32 D35 N38 F41 L42 E43 R46 R47 V50 E53 L54 R55 R56 L57 A58 H59 G60 G61 Q62 V63 N64 L65 E68 D69

H70 E73 D74 F75 V76 K77 P78 K79 G80 A81 F82 K83 A84 F85 T86 G87 E88 G89 Q90 K91 L92 G93 S94 T95 A96 P97 Q98 V99 L100 S101 T102

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *distance geometry*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Aria/CNS	structure solution	1.2
Aria/CNS	refinement	1.2

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	459	439	438	21±4
All	All	9180	8780	8760	425

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

5 of 139 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:30:LEU:HD12	1:A:75:PHE:HB2	0.89	1.44	3	19
1:A:19:TRP:CD1	1:A:30:LEU:HG	0.84	2.07	17	19
1:A:16:LEU:HD21	1:A:23:PHE:HB2	0.79	1.53	17	12
1:A:41:PHE:CZ	1:A:53:GLU:HG3	0.73	2.18	10	10
1:A:15:VAL:HG13	1:A:17:LYS:HE3	0.73	1.60	14	3

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	56/102 (55%)	49±2 (88±3%)	6±2 (10±3%)	1±1 (2±2%)	10 50
All	All	1120/2040 (55%)	985 (88%)	111 (10%)	24 (2%)	10 50

5 of 7 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	76	VAL	10
1	A	71	ARG	5
1	A	74	ASP	3
1	A	72	ASP	2
1	A	27	ASN	2

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	51/86 (59%)	46±1 (90±3%)	5±1 (10±3%)	10 55
All	All	1020/1720 (59%)	913 (90%)	107 (10%)	10 55

5 of 16 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	20	LYS	18
1	A	40	GLN	13
1	A	25	LEU	13
1	A	16	LEU	9
1	A	30	LEU	8

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided