



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 09:24 PM EDT

PDB ID : 3T1E
Title : The structure of the Newcastle disease virus hemagglutinin-neuraminidase (HN) ectodomain reveals a 4-helix bundle stalk
Authors : Yuan, P.; Swanson, K.; Leser, G.P.; Paterson, R.G.; Lamb, R.A.; Jardetzky, T.S.
Deposited on : 2011-07-21
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	537	 5% 50% 31% 17%
1	B	537	 5% 48% 31% 17%
1	E	537	 93%
1	F	537	 93%

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 6976 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin-neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	447	Total 3204	C 1989	N 549	O 646	S 20	0	0	0
1	B	445	Total 3194	C 1983	N 546	O 646	S 19	0	0	0
1	E	37	Total 289	C 184	N 46	O 58	S 1	0	0	0
1	F	37	Total 289	C 184	N 46	O 58	S 1	0	0	0

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	ALA	-	EXPRESSION TAG	UNP P12554
A	35	MET	-	EXPRESSION TAG	UNP P12554
A	36	ALA	-	EXPRESSION TAG	UNP P12554
A	37	HIS	-	EXPRESSION TAG	UNP P12554
A	38	HIS	-	EXPRESSION TAG	UNP P12554
A	39	HIS	-	EXPRESSION TAG	UNP P12554
A	40	HIS	-	EXPRESSION TAG	UNP P12554
A	41	HIS	-	EXPRESSION TAG	UNP P12554
A	42	HIS	-	EXPRESSION TAG	UNP P12554
A	43	LEU	-	EXPRESSION TAG	UNP P12554
A	44	VAL	-	EXPRESSION TAG	UNP P12554
A	45	PRO	-	EXPRESSION TAG	UNP P12554
A	46	ARG	-	EXPRESSION TAG	UNP P12554
A	47	GLY	-	EXPRESSION TAG	UNP P12554
A	48	SER	-	EXPRESSION TAG	UNP P12554
A	328	ALA	GLY	SEE REMARK 999	UNP P12554
B	34	ALA	-	EXPRESSION TAG	UNP P12554
B	35	MET	-	EXPRESSION TAG	UNP P12554
B	36	ALA	-	EXPRESSION TAG	UNP P12554
B	37	HIS	-	EXPRESSION TAG	UNP P12554
B	38	HIS	-	EXPRESSION TAG	UNP P12554

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	39	HIS	-	EXPRESSION TAG	UNP P12554
B	40	HIS	-	EXPRESSION TAG	UNP P12554
B	41	HIS	-	EXPRESSION TAG	UNP P12554
B	42	HIS	-	EXPRESSION TAG	UNP P12554
B	43	LEU	-	EXPRESSION TAG	UNP P12554
B	44	VAL	-	EXPRESSION TAG	UNP P12554
B	45	PRO	-	EXPRESSION TAG	UNP P12554
B	46	ARG	-	EXPRESSION TAG	UNP P12554
B	47	GLY	-	EXPRESSION TAG	UNP P12554
B	48	SER	-	EXPRESSION TAG	UNP P12554
B	328	ALA	GLY	SEE REMARK 999	UNP P12554
E	34	ALA	-	EXPRESSION TAG	UNP P12554
E	35	MET	-	EXPRESSION TAG	UNP P12554
E	36	ALA	-	EXPRESSION TAG	UNP P12554
E	37	HIS	-	EXPRESSION TAG	UNP P12554
E	38	HIS	-	EXPRESSION TAG	UNP P12554
E	39	HIS	-	EXPRESSION TAG	UNP P12554
E	40	HIS	-	EXPRESSION TAG	UNP P12554
E	41	HIS	-	EXPRESSION TAG	UNP P12554
E	42	HIS	-	EXPRESSION TAG	UNP P12554
E	43	LEU	-	EXPRESSION TAG	UNP P12554
E	44	VAL	-	EXPRESSION TAG	UNP P12554
E	45	PRO	-	EXPRESSION TAG	UNP P12554
E	46	ARG	-	EXPRESSION TAG	UNP P12554
E	47	GLY	-	EXPRESSION TAG	UNP P12554
E	48	SER	-	EXPRESSION TAG	UNP P12554
E	328	ALA	GLY	SEE REMARK 999	UNP P12554
F	34	ALA	-	EXPRESSION TAG	UNP P12554
F	35	MET	-	EXPRESSION TAG	UNP P12554
F	36	ALA	-	EXPRESSION TAG	UNP P12554
F	37	HIS	-	EXPRESSION TAG	UNP P12554
F	38	HIS	-	EXPRESSION TAG	UNP P12554
F	39	HIS	-	EXPRESSION TAG	UNP P12554
F	40	HIS	-	EXPRESSION TAG	UNP P12554
F	41	HIS	-	EXPRESSION TAG	UNP P12554
F	42	HIS	-	EXPRESSION TAG	UNP P12554
F	43	LEU	-	EXPRESSION TAG	UNP P12554
F	44	VAL	-	EXPRESSION TAG	UNP P12554
F	45	PRO	-	EXPRESSION TAG	UNP P12554
F	46	ARG	-	EXPRESSION TAG	UNP P12554
F	47	GLY	-	EXPRESSION TAG	UNP P12554
F	48	SER	-	EXPRESSION TAG	UNP P12554

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	328	ALA	GLY	SEE REMARK 999	UNP P12554

ALA	MET	ALA	HIS	HIS	HIS	HIS	HIS	LEU	VAL	PRO	ARG	GLY	SER	GLU	GLY	ILE	PRO	THR	ASN	GLN	D79	V80	V81	D82	R83	I84	Y85	K86	A95	T99	I102	I103		
PRO	THR	THR	THR	GLY	SER	GLY	CYS	THR	THR	ARG	ARG	ILE	PRO	ALA	GLY	GLY	THR	THR	THR															
ILE	ASN	LEU	ASP	ASP	THR	GLN	ASN	ASN	ARG	ARG	LYS	ILE	SER	PRO	GLY	GLY	LEU	CYS	GLY	THR	TYR	THR	THR	THR										
ASP	VAL	THR	THR	LEU	HIS	PHE	ILE	ARG	ASP	TRP	VAL	ALA	GLY	SER	TYR	PRO	VAL	SER	GLY	THR	PRO	GLY	GLY	THR	THR	THR								
GLU	GLN	ASP	TYR	THR	ILE	ARG	MET	ARG	ALA	LYS	VAL	ALA	GLY	SER	TYR	PRO	LYS	PRO	GLY	VAL	ARG	ARG	PHE	GLY	GLY	THR								
VAL	GLY	THR	THR	SER	HIS	PHE	LEU	TYR	ARG	GLN	ALA	THR	THR	THR	THR																			
THR	GLY	VAL	THR	THR	ASP	THR	PRO	TYR	ARG	GLN	ALA	THR	THR	THR	THR																			
THR	THR	SER	THR	CYS	PHE	ASP	LYS	VAL	VAL	THR	THR	THR	THR																					

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	138.29Å 138.29Å 167.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.89 – 3.30 48.89 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.89-3.30) 100.0 (48.89-3.30)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.63 (at 3.33Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.248 , 0.310 0.244 , 0.309	Depositor DCC
R_{free} test set	1241 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	126.9	Xtrriage
Anisotropy	0.164	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 117.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6976	wwPDB-VP
Average B, all atoms (Å ²)	134.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/3288	0.52	0/4502
1	B	0.29	0/3278	0.51	0/4490
1	E	0.29	0/291	0.46	0/396
1	F	0.28	0/291	0.51	0/396
All	All	0.30	0/7148	0.51	0/9784

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3204	0	2730	165	0
1	B	3194	0	2719	180	0
1	E	289	0	299	19	0
1	F	289	0	299	29	0
All	All	6976	0	6047	377	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

The worst 5 of 377 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:451:SER:H	1:B:466:THR:HG21	1.16	1.06
1:A:224:ARG:HG3	1:A:224:ARG:HH11	1.22	1.04
1:B:152:TYR:HE2	1:B:565:LEU:HB2	1.23	1.02
1:B:471:ASP:HB2	1:B:527:THR:HA	1.49	0.95
1:B:152:TYR:CE2	1:B:565:LEU:HB2	2.03	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/537 (83%)	413 (93%)	31 (7%)	1 (0%)	47	77
1	B	443/537 (82%)	406 (92%)	36 (8%)	1 (0%)	47	77
1	E	35/537 (6%)	29 (83%)	5 (14%)	1 (3%)	4	24
1	F	35/537 (6%)	28 (80%)	7 (20%)	0	100	100
All	All	958/2148 (45%)	876 (91%)	79 (8%)	3 (0%)	41	71

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	301	VAL
1	B	301	VAL
1	E	80	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/463 (68%)	288 (92%)	25 (8%)	12	37
1	B	313/463 (68%)	288 (92%)	25 (8%)	12	37
1	E	34/463 (7%)	28 (82%)	6 (18%)	2	8
1	F	34/463 (7%)	32 (94%)	2 (6%)	19	49
All	All	694/1852 (38%)	636 (92%)	58 (8%)	11	35

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	184	HIS
1	E	115	ASN
1	B	290	LEU
1	E	110	LEU
1	B	565	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	462	ASN
1	B	128	HIS
1	B	271	HIS
1	A	340	ASN
1	A	200	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	447/537 (83%)	0.07	4 (0%) 84 84	83, 111, 163, 215	0
1	B	445/537 (82%)	0.23	28 (6%) 20 20	83, 144, 200, 255	0
1	E	37/537 (6%)	0.02	0 100 100	100, 132, 177, 191	0
1	F	37/537 (6%)	-0.16	0 100 100	103, 133, 203, 212	0
All	All	966/2148 (44%)	0.14	32 (3%) 46 44	83, 129, 190, 255	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	382	GLY	4.5
1	A	485	GLY	3.6
1	B	274	LEU	3.5
1	B	383	GLU	3.3
1	B	345	ASP	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.