Full wwPDB X-ray Structure Validation Report (i)

Sep 3, 2023 – 09:47 AM EDT

PDB ID	:	3T8A
Title	:	Crystal structure of Mycobacterium tuberculosis MenB in complex with substrate analogue, OSB-NCoA
Authors	:	Li, H.-J.; Li, X.; Liu, N.; Zhang, H.; Truglio, J.; Mishra, S.; Kisker, C.; Garcia-Diaz, M.; Tonge, P.
Deposited on	:	2011-08-01
Resolution	:	2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2391 (2.26-2.22)
Clashscore	141614	2539 (2.26-2.22)
Ramachandran outliers	138981	2489 (2.26-2.22)
Sidechain outliers	138945	2490 (2.26-2.22)
RSRZ outliers	127900	2353 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	4% 70% 5% 24%
1	B	334	8% 75% 7% 18%
1	C	334	4% 64% 10% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	S0N	C	600	-	-	-	X

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 6064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1,4-Dihydroxy-2-naphthoyl-CoA synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	253	1929	1217	350	353	9	0	1	1
1	B	274	2089	1325	370	385	9	0	0	0
1	C	250	1863	1183	334	338	8	0	0	1

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP O06414
A	-18	GLY	-	expression tag	UNP O06414
A	-17	SER	-	expression tag	UNP O06414
A	-16	SER	-	expression tag	UNP O06414
A	-15	HIS	-	expression tag	UNP O06414
A	-14	HIS	-	expression tag	UNP O06414
A	-13	HIS	-	expression tag	UNP O06414
A	-12	HIS	-	expression tag	UNP O06414
A	-11	HIS	-	expression tag	UNP O06414
A	-10	HIS	-	expression tag	UNP O06414
A	-9	SER	-	expression tag	UNP O06414
A	-8	SER	-	expression tag	UNP O06414
A	-7	GLY	-	expression tag	UNP O06414
A	-6	LEU	-	expression tag	UNP O06414
A	-5	VAL	-	expression tag	UNP O06414
A	-4	PRO	-	expression tag	UNP O06414
A	-3	ARG	-	expression tag	UNP O06414
A	-2	GLY	-	expression tag	UNP O06414
A	-1	SER	-	expression tag	UNP O06414
A	0	HIS	-	expression tag	UNP O06414
B	-19	MET	-	expression tag	UNP O06414
B	-18	GLY	-	expression tag	UNP O06414
B	-17	SER	-	expression tag	UNP O06414

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	SER	-	expression tag	UNP O06414
B	-15	HIS	-	expression tag	UNP O06414
B	-14	HIS	-	expression tag	UNP O06414
B	-13	HIS	-	expression tag	UNP O06414
B	-12	HIS	-	expression tag	UNP O06414
B	-11	HIS	-	expression tag	UNP O06414
B	-10	HIS	-	expression tag	UNP O06414
B	-9	SER	-	expression tag	UNP O06414
B	-8	SER	-	expression tag	UNP O06414
B	-7	GLY	-	expression tag	UNP O06414
B	-6	LEU	-	expression tag	UNP O06414
B	-5	VAL	-	expression tag	UNP O06414
B	-4	PRO	-	expression tag	UNP O06414
B	-3	ARG	-	expression tag	UNP O06414
B	-2	GLY	-	expression tag	UNP O06414
B	-1	SER	-	expression tag	UNP O06414
B	0	HIS	-	expression tag	UNP O06414
C	-19	MET	-	expression tag	UNP O06414
C	-18	GLY	-	expression tag	UNP O06414
C	-17	SER	-	expression tag	UNP O06414
C	-16	SER	-	expression tag	UNP O06414
C	-15	HIS	-	expression tag	UNP O06414
C	-14	HIS	-	expression tag	UNP O06414
C	-13	HIS	-	expression tag	UNP O06414
C	-12	HIS	-	expression tag	UNP O06414
C	-11	HIS	-	expression tag	UNP O06414
C	-10	HIS	-	expression tag	UNP O06414
C	-9	SER	-	expression tag	UNP O06414
C	-8	SER	-	expression tag	UNP O06414
C	-7	GLY	-	expression tag	UNP O06414
C	-6	LEU	-	expression tag	UNP O06414
C	-5	VAL	-	expression tag	UNP O06414
C	-4	PRO	-	expression tag	UNP O06414
C	-3	ARG	-	expression tag	UNP O06414
C	-2	GLY	-	expression tag	UNP O06414
C	-1	SER	-	expression tag	UNP O06414
C	0	HIS	-	expression tag	UNP O06414

- Molecule 2 is o-succinylbenzoyl-N-coenzyme A (three-letter code: S0N) (formula: $C_{32}H_{45}N_8O_{20}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	40	16	6	15	3	0	0
			Total	C	N	O	P		
2	C	1	36	15	5	13	3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	38	38	38	0	0
			Total	O		
3	B	42	42	42	0	0
			Total	O		
3	C	27	27	27	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 1,4-Dihydroxy-2-naphthoyl-CoA synthase

Chain A:



• Molecule 1: 1,4-Dihydroxy-2-naphthoyl-CoA synthase

Chain B:



• Molecule 1: 1,4-Dihydroxy-2-naphthoyl-CoA synthase

Chain C:




L165
H166
V167
V168
K182
Q183
T184
D187
S190
A198
A201
R202
F213
L272
D290
GLU
ALA
VAL
GLU
GLY
ARG
ASP
ALA
PHE
LEU
GLN
LYS
ARG
PRO
PRO
ASP
TRP
SER
PRO
PHE
PRO
ARG
TYR
PHE

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α, β, γ	87.13Å 87.13Å 414.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.61 – 2.25 46.61 – 2.25	Depositor EDS
% Data completeness (in resolution range)	94.0 (46.61-2.25) 94.1 (46.61-2.25)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.67 (at 2.24Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.194 , 0.237 0.188 , 0.229	Depositor DCC
R_{free} test set	2287 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	30.3	Xtriage
Anisotropy	0.738	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.39 , 52.7	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	6064	wwPDB-VP
Average B, all atoms (Å^2)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: S0N

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.36	0/1975	0.49	0/2673
1	B	0.37	0/2137	0.50	0/2893
1	C	0.33	0/1906	0.48	0/2579
All	All	0.35	0/6018	0.49	0/8145

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1929	0	1833	9	0
1	B	2089	0	1958	15	0
1	C	1863	0	1746	26	0
2	B	40	0	21	2	0
2	C	36	0	19	8	0
3	A	38	0	0	0	0
3	B	42	0	0	0	0
3	C	27	0	0	0	0
All	All	6064	0	5577	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:107:GLN:H	2:C:600:S0N:C25	2.08	0.67
1:B:268:LEU:HG	1:C:198:ALA:HB1	1.79	0.65
1:B:268:LEU:HD11	1:C:198:ALA:O	1.99	0.62
1:C:92:PRO:HG3	1:C:99:TRP:CZ2	2.36	0.61
1:C:66:VAL:HG11	1:C:135:HIS:HB3	1.82	0.60
1:A:255:PRO:HD2	1:B:291:GLU:OE1	2.05	0.57
1:A:51:ALA:HA	1:A:88:THR:O	2.07	0.54
1:C:159:ALA:HA	1:C:182:LYS:O	2.08	0.54
1:B:260:MET:SD	1:B:281:GLU:HB3	2.48	0.53
1:C:58:ARG:HG3	2:C:600:S0N:H14	1.89	0.53
1:B:107:GLN:H	2:B:500:S0N:C25	2.23	0.51
1:A:159:ALA:HA	1:A:182:LYS:O	2.11	0.51
1:B:58:ARG:HG2	2:B:500:S0N:H14	1.93	0.50
1:B:184:THR:O	1:B:187:ASP:HB2	2.12	0.49
1:C:62:ARG:HB2	1:C:63:PRO:HD2	1.94	0.49
1:A:260:MET:SD	1:A:281:GLU:HB3	2.52	0.48
1:B:51:ALA:HA	1:B:88:THR:O	2.14	0.48
1:B:78:MET:O	1:B:80:PRO:HD3	2.14	0.47
1:B:256:GLN:O	1:B:260:MET:HG2	2.15	0.47
1:C:184:THR:O	1:C:187:ASP:HB2	2.15	0.46
1:C:16:SER:C	1:C:18:ASN:H	2.19	0.46
1:C:20:PHE:CE2	1:C:22:ALA:HA	2.51	0.46
1:C:36:ASP:HB3	1:C:65:THR:HG23	1.97	0.45
1:B:159:ALA:HA	1:B:182:LYS:O	2.16	0.45
1:C:165:LEU:HD23	1:C:165:LEU:HA	1.83	0.45
1:C:165:LEU:O	1:C:168:VAL:HG12	2.16	0.45
1:C:58:ARG:HD2	2:C:600:S0N:H04	1.99	0.44
1:C:106:ASP:OD1	2:C:600:S0N:H25	2.18	0.44
1:A:183:GLN:HG2	1:A:213:PHE:CE2	2.53	0.44
1:A:291:GLU:OE1	1:B:255:PRO:HD2	2.18	0.44
1:C:39:TYR:CG	1:C:72:VAL:HG13	2.53	0.44
1:C:54:ARG:N	1:C:55:PRO:HD3	2.33	0.43
1:C:106:ASP:HA	2:C:600:S0N:N24	2.34	0.43
1:C:146:MET:HA	1:C:147:PRO:HD3	1.83	0.43
1:A:36:ASP:HB3	1:A:65:THR:HG23	2.01	0.43
1:C:25:TRP:HB3	1:C:40:HIS:HB3	2.00	0.43
1:C:107:GLN:H	2:C:600:S0N:H25	1.84	0.43
1:C:107:GLN:N	2:C:600:S0N:H25	2.35	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:SER:HA	1:A:80:PRO:HD3	1.87	0.42
1:B:147:PRO:HB2	1:B:259:ARG:HD2	2.02	0.42
1:C:183:GLN:HG2	1:C:213:PHE:CE2	2.54	0.41
1:B:308:SER:N	1:B:309:PRO:CD	2.83	0.41
1:C:51:ALA:HA	1:C:88:THR:O	2.21	0.41
1:A:184:THR:O	1:A:187:ASP:HB2	2.21	0.40
1:C:107:GLN:N	2:C:600:S0N:C25	2.81	0.40
1:B:25:TRP:HB3	1:B:40:HIS:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/334 (75%)	245 (98%)	5 (2%)	0	100	100
1	B	268/334 (80%)	262 (98%)	6 (2%)	0	100	100
1	C	246/334 (74%)	237 (96%)	9 (4%)	0	100	100
All	All	764/1002 (76%)	744 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/264 (72%)	187 (99%)	2 (1%)	73	80
1	B	201/264 (76%)	199 (99%)	2 (1%)	76	82
1	C	178/264 (67%)	175 (98%)	3 (2%)	60	68
All	All	568/792 (72%)	561 (99%)	7 (1%)	71	78

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	VAL
1	A	166	HIS
1	B	15	LEU
1	B	166	HIS
1	C	57	VAL
1	C	147	PRO
1	C	166	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	107	GLN
1	C	135	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	S0N	B	500	-	35,42,66	2.18	9 (25%)	45,66,97	1.60	7 (15%)
2	S0N	C	600	-	33,38,66	1.98	6 (18%)	40,60,97	1.45	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	S0N	B	500	-	-	8/33/54/80	0/3/3/4
2	S0N	C	600	-	-	7/24/44/80	0/3/3/4

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	S0N	C25-N26	6.62	1.42	1.32
2	C	600	S0N	C25-N26	6.62	1.42	1.32
2	B	500	S0N	C38-N40	5.76	1.47	1.32
2	C	600	S0N	O16-C17	5.18	1.48	1.41
2	B	500	S0N	O16-C17	5.16	1.48	1.41
2	B	500	S0N	C22-N23	3.48	1.46	1.34
2	C	600	S0N	C22-N23	3.47	1.46	1.34
2	C	600	S0N	C19-N20	3.39	1.40	1.34
2	B	500	S0N	C19-N20	3.09	1.40	1.34
2	B	500	S0N	P32-O35	2.67	1.59	1.50
2	B	500	S0N	O39-C38	-2.67	1.18	1.23
2	C	600	S0N	O29-C28	2.21	1.48	1.43
2	B	500	S0N	P10-O11	2.18	1.58	1.50
2	B	500	S0N	O29-C28	2.10	1.47	1.43
2	C	600	S0N	P32-O31	2.03	1.63	1.59

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	500	S0N	N26-C25-N24	-5.63	119.88	128.68
2	C	600	S0N	N26-C25-N24	-5.42	120.20	128.68
2	B	500	S0N	C01-C02-C36	4.28	116.23	108.82
2	B	500	S0N	C36-C38-N40	4.11	120.56	117.18
2	C	600	S0N	P10-O09-P06	-3.78	119.86	132.83
2	B	500	S0N	P10-O09-P06	-3.29	121.52	132.83
2	C	600	S0N	C30-C28-C17	2.81	106.12	99.89
2	B	500	S0N	O05-C04-C02	-2.39	106.70	110.55
2	B	500	S0N	O37-C36-C02	-2.24	104.99	110.25
2	B	500	S0N	O39-C38-N40	-2.06	119.42	123.00
2	C	600	S0N	O13-C14-C15	2.05	116.06	108.99

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	600	S0N	C14-O13-P10-O11
2	C	600	S0N	C30-O31-P32-O33
2	B	500	S0N	C01-C02-C04-O05
2	C	600	S0N	O13-C14-C15-O16
2	B	500	S0N	C03-C02-C04-O05
2	C	600	S0N	O13-C14-C15-C30
2	C	600	S0N	C01-C02-C04-O05
2	B	500	S0N	C04-O05-P06-O08
2	B	500	S0N	C36-C02-C04-O05
2	B	500	S0N	C02-C36-C38-O39
2	C	600	S0N	C03-C02-C04-O05
2	B	500	S0N	C30-O31-P32-O35
2	B	500	S0N	C04-O05-P06-O09
2	B	500	S0N	O37-C36-C38-N40
2	C	600	S0N	C36-C02-C04-O05

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	500	S0N	2	0
2	C	600	S0N	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand S0N B 500



Bond lengths



Bond angles



Torsions



Rings

Ligand S0N C 600



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	253/334 (75%)	0.02	13 (5%)	28	27	23, 34, 56, 74	0
1	B	274/334 (82%)	0.17	27 (9%)	7	6	22, 32, 63, 78	1 (0%)
1	C	250/334 (74%)	0.14	13 (5%)	27	26	22, 41, 63, 85	0
All	All	777/1002 (77%)	0.11	53 (6%)	17	17	22, 35, 63, 85	1 (0%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	310	PHE	5.1
1	B	307	TRP	4.5
1	B	309	PRO	4.2
1	B	304	PRO	4.1
1	B	306	ASP	4.1
1	B	300	LEU	4.0
1	B	299	PHE	3.7
1	A	290	ASP	3.4
1	B	267	LEU	3.4
1	B	297	ASP	3.2
1	B	272	LEU	3.1
1	A	293	VAL	3.0
1	A	287	TYR	3.0
1	A	271	GLY	2.9
1	C	94	PRO	2.9
1	C	97	GLY	2.8
1	B	292	ALA	2.8
1	A	272	LEU	2.8
1	B	290	ASP	2.8
1	B	294	GLU	2.7
1	A	267	LEU	2.7
1	C	272	LEU	2.7
1	B	293	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	295	GLY	2.6
1	A	190	SER	2.6
1	A	187	ASP	2.6
1	C	45	ASP	2.5
1	B	296	ARG	2.5
1	C	201	ALA	2.5
1	C	44	ASP	2.4
1	B	271	GLY	2.4
1	C	190	SER	2.4
1	B	201	ALA	2.4
1	B	305	PRO	2.4
1	C	202	ARG	2.3
1	B	15	LEU	2.3
1	C	29	ASP	2.3
1	C	91	GLY	2.3
1	A	292	ALA	2.3
1	B	291	GLU	2.2
1	B	295	GLY	2.2
1	B	288	MET	2.2
1	C	290	ASP	2.2
1	A	273	VAL	2.2
1	B	273	VAL	2.1
1	B	200	LEU	2.1
1	A	201	ALA	2.1
1	C	43	VAL	2.1
1	B	274	GLY	2.1
1	B	43	VAL	2.1
1	C	96	ASP	2.0
1	A	134	LEU	2.0
1	B	202	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	S0N	C	600	36/63	0.76	0.44	52,72,83,84	27
2	S0N	B	500	40/63	0.92	0.16	41,52,64,67	22

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around S0N C 600:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around S0N B 500:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.