



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 28, 2020 – 08:44 pm BST

PDB ID : 1TXK  
Title : Crystal structure of Escherichia coli OpgG  
Authors : Hanouille, X.; Rollet, E.; Clantin, B.; Landrieu, I.; Odberg-Ferragut, C.; Lip-pens, G.; Bohin, J.P.; Villeret, V.  
Deposited on : 2004-07-05  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	498	
1	B	498	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucans biosynthesis protein G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	490	3922	2495	673	744	10	0	0	0
1	B	487	3891	2477	664	739	1 10	0	0	0

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	MET	-	INITIATING METHIONINE	UNP P33136
A	54	MSE	MET	MODIFIED RESIDUE	UNP P33136
A	87	MSE	MET	MODIFIED RESIDUE	UNP P33136
A	151	MSE	MET	MODIFIED RESIDUE	UNP P33136
A	223	MSE	MET	MODIFIED RESIDUE	UNP P33136
A	253	MSE	MET	MODIFIED RESIDUE	UNP P33136
A	304	MSE	MET	MODIFIED RESIDUE	UNP P33136
A	379	MSE	MET	MODIFIED RESIDUE	UNP P33136
A	436	MSE	MET	MODIFIED RESIDUE	UNP P33136
A	475	MSE	MET	MODIFIED RESIDUE	UNP P33136
A	488	MSE	MET	MODIFIED RESIDUE	UNP P33136
A	512	VAL	-	EXPRESSION TAG	UNP P33136
A	513	GLU	-	EXPRESSION TAG	UNP P33136
A	514	HIS	-	EXPRESSION TAG	UNP P33136
A	515	HIS	-	EXPRESSION TAG	UNP P33136
A	516	HIS	-	EXPRESSION TAG	UNP P33136
A	517	HIS	-	EXPRESSION TAG	UNP P33136
A	518	HIS	-	EXPRESSION TAG	UNP P33136
A	519	HIS	-	EXPRESSION TAG	UNP P33136
B	22	MET	-	INITIATING METHIONINE	UNP P33136
B	54	MSE	MET	MODIFIED RESIDUE	UNP P33136
B	87	MSE	MET	MODIFIED RESIDUE	UNP P33136
B	151	MSE	MET	MODIFIED RESIDUE	UNP P33136
B	223	MSE	MET	MODIFIED RESIDUE	UNP P33136
B	253	MSE	MET	MODIFIED RESIDUE	UNP P33136

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	304	MSE	MET	MODIFIED RESIDUE	UNP P33136
B	379	MSE	MET	MODIFIED RESIDUE	UNP P33136
B	436	MSE	MET	MODIFIED RESIDUE	UNP P33136
B	475	MSE	MET	MODIFIED RESIDUE	UNP P33136
B	488	MSE	MET	MODIFIED RESIDUE	UNP P33136
B	512	VAL	-	EXPRESSION TAG	UNP P33136
B	513	GLU	-	EXPRESSION TAG	UNP P33136
B	514	HIS	-	EXPRESSION TAG	UNP P33136
B	515	HIS	-	EXPRESSION TAG	UNP P33136
B	516	HIS	-	EXPRESSION TAG	UNP P33136
B	517	HIS	-	EXPRESSION TAG	UNP P33136
B	518	HIS	-	EXPRESSION TAG	UNP P33136
B	519	HIS	-	EXPRESSION TAG	UNP P33136

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Na 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	243	Total O 243 243	0	0
3	B	234	Total O 234 234	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.86Å 88.12Å 215.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.99 – 2.50 20.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.0 (19.99-2.50) 99.6 (20.00-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.18 (at 2.50Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.192 , 0.229 0.206 , 0.239	Depositor DCC
$R_{free}$ test set	7856 reflections (9.76%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.3	Xtrriage
Anisotropy	0.459	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 51.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8291	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/4017	0.67	4/5438 (0.1%)
1	B	0.37	0/3983	0.64	0/5391
All	All	0.38	0/8000	0.65	4/10829 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	492	LEU	O-C-N	6.71	133.44	122.70
1	A	492	LEU	C-N-CA	-6.08	106.51	121.70
1	A	492	LEU	CA-C-N	-5.67	104.73	117.20
1	A	348	VAL	N-CA-C	-5.29	96.72	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3922	0	3814	123	0
1	B	3891	0	3796	130	0
2	A	1	0	0	0	0
3	A	243	0	0	4	0
3	B	234	0	0	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8291	0	7610	248	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 248 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:253:MSE:HE1	1:B:255:LEU:HB2	1.24	1.18
1:A:253:MSE:HE1	1:A:255:LEU:HB2	1.21	1.08
1:B:409:THR:HG23	1:B:410:GLY:H	1.11	1.06
1:B:253:MSE:HE3	1:B:358:GLU:HA	1.36	1.03
1:A:253:MSE:HE3	1:A:358:GLU:HA	1.36	1.02

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	486/498 (98%)	464 (96%)	19 (4%)	3 (1%)	25	43
1	B	483/498 (97%)	457 (95%)	22 (5%)	4 (1%)	19	35
All	All	969/996 (97%)	921 (95%)	41 (4%)	7 (1%)	22	39

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	119	GLN
1	B	409	THR
1	A	48	PRO
1	A	119	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	410	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	425/423 (100%)	406 (96%)	19 (4%)	27	51
1	B	422/423 (100%)	403 (96%)	19 (4%)	27	51
All	All	847/846 (100%)	809 (96%)	38 (4%)	27	51

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	488	MSE
1	B	55	LYS
1	B	293	ASN
1	B	22	MET
1	B	65	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	31	GLN
1	B	96	ASN
1	B	414	GLN
1	B	63	GLN
1	B	65	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	480/498 (96%)	0.13	25 (5%) 27 29	19, 32, 63, 82	0
1	B	477/498 (95%)	0.09	23 (4%) 30 32	20, 33, 57, 78	0
All	All	957/996 (96%)	0.11	48 (5%) 28 30	19, 33, 60, 82	0

The worst 5 of 48 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	515	HIS	8.7
1	B	119	GLN	7.0
1	A	514	HIS	6.9
1	A	420	GLN	6.1
1	B	123	ASP	5.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NA	A	600	1/1	0.88	0.19	38,38,38,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.