Full wwPDB X-ray Structure Validation Report (i)

Sep 24, 2023 – 06:52 AM EDT

PDB ID	:	5UIK
Title	:	X-ray structure of the FdtF formyltransferase from salmonella enteric O60 in complex with TDP-Fuc3N and folinic acid
Authors	:	Woodford, C.R.; Thoden, J.B.; Holden, H.M.
Deposited on	:	2017-01-14
Resolution	:	2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	405	86% 11% • •
1	B	405	89% 8% • •

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 6957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Formyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	398	3231	2097	513	609	12	0	0	0
1	B	398	3240	2103	514	611	12	0	1	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	GLY	-	expression tag	UNP U3GK13
A	-5	GLY	-	expression tag	UNP U3GK13
A	-4	GLY	-	expression tag	UNP U3GK13
A	-3	GLY	-	expression tag	UNP U3GK13
A	-2	GLY	-	expression tag	UNP U3GK13
A	-1	GLY	-	expression tag	UNP U3GK13
A	0	HIS	-	expression tag	UNP U3GK13
A	395	ALA	GLU	engineered mutation	UNP U3GK13
B	-6	GLY	-	expression tag	UNP U3GK13
B	-5	GLY	-	expression tag	UNP U3GK13
B	-4	GLY	-	expression tag	UNP U3GK13
B	-3	GLY	-	expression tag	UNP U3GK13
B	-2	GLY	-	expression tag	UNP U3GK13
B	-1	GLY	-	expression tag	UNP U3GK13
B	0	HIS	-	expression tag	UNP U3GK13
B	395	ALA	GLU	engineered mutation	UNP U3GK13

- Molecule 2 is (3R,4S,5R,6R)-4-amino-3,5-dihydroxy-6-methyloxan-2-yl][hydroxy-[[(2R,3S,5R)-3-hydroxy-5-(5-methyl-2,4-dioxopyrimidin-1-yl)oxolan-2-yl]methoxy]phosphoryl] hydrogen phosphate (three-letter code: T3F) (formula: $C_{16}H_{27}N_3O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	35	16	3	14	2	0	0
			Total	C	N	O	P		
2	A	1	35	16	3	14	2	0	0
			Total	C	N	O	P		
2	B	1	34	16	3	13	2	0	0
			Total	C	N	O	P		
2	B	1	35	16	3	14	2	0	0

- Molecule 3 is N-{[4-({[(6R)-2-amino-5-formyl-4-oxo-1,4,5,6,7,8-hexahydropteridin-6-yl]methyl}amino)phenyl]carbonyl}-L-glutamic acid (three-letter code: FON) (formula: $C_{20}H_{23}N_7O_7$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	25	15	7	3	0	0
3	B	1	25	15	7	3	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Na		
4	A	1	1	1	0	0
4	B	2	2	2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	132	132	132	0	0
5	B	162	162	162	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Formyltransferase

Chain A:



• Molecule 1: Formyltransferase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	66.24Å 69.20Å 200.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.09 – 2.20 48.09 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.3 (48.09-2.20) 97.3 (48.09-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$< I/\sigma(I) >$ [1]	1.69 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0124	Depositor
R, R_{free}	0.206 , 0.271 0.210 , 0.270	Depositor DCC
R_{free} test set	2366 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å^2)	25.0	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 35.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.47$, $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	0.030 for k,h,-l	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	6957	wwPDB-VP
Average B, all atoms (Å^2)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FON, T3F, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.64	0/3297	0.83	5/4448 (0.1%)
1	B	0.66	0/3309	0.80	2/4462 (0.0%)
All	All	0.65	0/6606	0.82	7/8910 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	44	ARG	NE-CZ-NH2	-7.60	116.50	120.30
1	A	44	ARG	NE-CZ-NH1	7.16	123.88	120.30
1	A	93	ILE	CG1-CB-CG2	-6.70	96.66	111.40
1	A	214	ARG	NE-CZ-NH2	6.06	123.33	120.30
1	B	17	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	A	240	ARG	NE-CZ-NH2	-5.17	117.71	120.30
1	B	240	ARG	NE-CZ-NH2	5.00	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3231	0	3262	19	0
1	B	3240	0	3278	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	70	0	50	0	0
2	B	69	0	50	0	0
3	A	25	0	15	1	0
3	B	25	0	15	1	0
4	A	1	0	0	0	0
4	B	2	0	0	0	0
5	A	132	0	0	0	0
5	B	162	0	0	0	0
All	All	6957	0	6670	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:276:LEU:HD22	1:A:310:VAL:HG21	1.81	0.63
1:A:240:ARG:NH2	1:B:179:GLU:OE1	2.37	0.57
1:A:213:LEU:HD11	1:A:267:LEU:CD2	2.35	0.56
1:A:19:LEU:HD21	1:A:167:ILE:HD13	1.88	0.55
1:B:227:ASP:OD2	1:B:251:ARG:NH1	2.39	0.55
1:A:348:MET:HE1	1:A:357:LEU:HD22	1.88	0.54
1:B:1:MET:HB2	1:B:26:ILE:HD12	1.90	0.52
1:A:237:LEU:HD21	1:A:266:VAL:CG2	2.42	0.49
1:B:235:ASP:OD2	1:B:268:TYR:OH	2.14	0.49
1:B:80:VAL:H	3:B:402:FON:HN1	1.61	0.49
1:A:69:ALA:O	1:A:87:HIS:CE1	2.67	0.48
1:A:213:LEU:HD11	1:A:267:LEU:HD23	1.95	0.47
1:A:42:PHE:CD2	1:A:220:GLU:HG3	2.50	0.47
1:A:80:VAL:H	3:A:402:FON:HN1	1.63	0.47
1:B:7:GLY:HA3	1:B:74:LEU:O	2.15	0.47
1:A:93:ILE:CD1	1:A:122:LEU:HD13	2.46	0.45
1:B:213:LEU:HD11	1:B:267:LEU:HD22	1.99	0.45
1:B:9:ASN:HB3	1:B:43:GLN:OE1	2.18	0.43
1:A:261:ILE:HG13	1:B:187:TYR:CD2	2.54	0.43
1:A:319:LEU:HD22	1:A:323:LEU:HG	2.00	0.43
1:B:345:LYS:NZ	1:B:346:ASP:OD2	2.52	0.43
1:A:190:LYS:H	1:A:190:LYS:CE	2.32	0.42
1:A:229:VAL:HG11	1:A:269:LYS:HG2	2.02	0.42
1:A:357:LEU:HD23	1:A:396:PHE:CE2	2.55	0.42
1:B:129:ILE:N	1:B:129:ILE:HD12	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:237:LEU:HD21	1:B:266:VAL:CG2	2.49	0.41
1:B:70:ILE:HG22	1:B:89:GLU:HB3	2.01	0.41
1:B:389:LEU:HD22	1:B:393:MET:HG3	2.03	0.41
1:A:312:ALA:HA	1:A:320:ILE:HD11	2.03	0.41
1:A:241:SER:HB2	1:A:249:LEU:HD11	2.03	0.40
1:A:44:ARG:HD3	1:A:372:TYR:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/405 (98%)	368 (93%)	27 (7%)	1 (0%)	41	46
1	B	397/405 (98%)	385 (97%)	11 (3%)	1 (0%)	41	46
All	All	793/810 (98%)	753 (95%)	38 (5%)	2 (0%)	41	46

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	316	ASN
1	B	24	LYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/359 (99%)	339 (95%)	18 (5%)	24	30
1	B	359/359 (100%)	346 (96%)	13 (4%)	35	45
All	All	716/718 (100%)	685 (96%)	31 (4%)	29	36

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67	ASP
1	A	86	ASN
1	A	93	ILE
1	A	138	LYS
1	A	190	LYS
1	A	191	LYS
1	A	218	PHE
1	A	235	ASP
1	A	244	LYS
1	A	254	LYS
1	A	273	LYS
1	A	291	ILE
1	A	308	VAL
1	A	319	LEU
1	A	346	ASP
1	A	374	ASP
1	A	388	ASN
1	A	389	LEU
1	B	24	LYS
1	B	137	GLN
1	B	218	PHE
1	B	240	ARG
1	B	244	LYS
1	B	290	LEU
1	B	292	ARG
1	B	294	LYS
1	B	330	ILE
1	B	345	LYS
1	B	349	LEU
1	B	389	LEU
1	B	394	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	A	92	ASN
1	A	127	HIS
1	A	143	GLN
1	A	329	ASN
1	A	388	ASN
1	B	92	ASN
1	B	143	GLN
1	B	388	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	T3F	A	403	-	34,37,37	1.69	5 (14%)	52,57,57	1.45	7 (13%)
2	T3F	A	401	-	34,37,37	1.65	3 (8%)	52,57,57	2.01	14 (26%)
3	FON	B	402	-	25,27,36	1.90	7 (28%)	24,38,50	2.75	11 (45%)
3	FON	A	402	-	25,27,36	1.73	3 (12%)	24,38,50	1.96	7 (29%)
2	T3F	B	401	-	32,36,37	2.17	4 (12%)	47,55,57	1.77	6 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	T3F	B	403	-	34,37,37	1.51	4 (11%)	52,57,57	1.91	13 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	T3F	A	403	-	-	4/21/53/53	0/3/3/3
2	T3F	A	401	-	-	8/21/53/53	0/3/3/3
3	FON	B	402	-	-	4/11/24/37	0/2/3/3
3	FON	A	402	-	-	0/11/24/37	0/2/3/3
2	T3F	B	401	-	-	1/20/53/53	0/3/3/3
2	T3F	B	403	-	-	0/21/53/53	0/3/3/3

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	T3F	C2-N3	-7.77	1.37	1.46
2	A	401	T3F	O4-C4	7.12	1.37	1.23
2	A	403	T3F	O4-C4	6.54	1.36	1.23
2	B	401	T3F	O4-C4	6.41	1.35	1.23
2	B	403	T3F	O4-C4	5.87	1.34	1.23
3	B	402	FON	O4-C4	5.60	1.38	1.24
3	A	402	FON	O4-C4	4.72	1.36	1.24
2	B	401	T3F	C4-N3	4.48	1.39	1.33
3	A	402	FON	CP1-N5	-4.36	1.29	1.35
3	B	402	FON	CP1-N5	-4.08	1.30	1.35
2	A	403	T3F	C4-C5	-3.51	1.39	1.44
2	B	403	T3F	C4-C5	-3.44	1.39	1.44
2	A	401	T3F	C4-C5	-3.42	1.39	1.44
2	A	403	T3F	C6-C5	3.11	1.39	1.34
2	B	401	T3F	C4-C5	-3.09	1.39	1.44
2	A	403	T3F	C2-N1	-2.90	1.33	1.38
3	B	402	FON	C7-C6	2.70	1.55	1.52
2	B	403	T3F	C6-C5	2.65	1.38	1.34
3	B	402	FON	C11-C	2.64	1.54	1.50
2	A	403	T3F	C4-N3	-2.59	1.34	1.38
2	A	401	T3F	C6-C5	2.56	1.38	1.34
3	A	402	FON	C7-N8	2.37	1.48	1.44
2	B	403	T3F	C4-N3	-2.26	1.34	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	FON	C13-C12	2.21	1.42	1.38
3	B	402	FON	C13-C14	2.02	1.42	1.39
3	B	402	FON	C12-C11	2.02	1.42	1.39

All (58) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	402	FON	C8A-C4A-C4	8.14	120.80	114.44
2	B	401	T3F	C5-C4-N3	6.43	120.80	115.31
2	B	401	T3F	O4-C4-C5	-6.09	117.84	124.90
2	A	401	T3F	C4-N3-C2	-5.99	119.60	127.35
2	A	401	T3F	N3-C2-N1	5.94	122.77	114.89
2	B	403	T3F	C5-C4-N3	5.87	120.32	115.31
3	B	402	FON	C11-C-N	5.19	123.97	117.75
2	B	403	T3F	C4-N3-C2	-5.03	120.84	127.35
2	A	401	T3F	C5-C4-N3	5.02	119.59	115.31
2	A	403	T3F	C5-C4-N3	4.19	118.89	115.31
3	A	402	FON	C8A-C4A-C4	4.18	117.71	114.44
2	B	403	T3F	O4-C4-C5	-3.75	120.56	124.90
3	B	402	FON	C2-N3-C4	3.73	121.85	115.93
3	B	402	FON	O-C-N	-3.70	117.32	122.58
2	B	403	T3F	N3-C2-N1	3.51	119.55	114.89
2	A	403	T3F	C4-N3-C2	-3.51	122.81	127.35
3	A	402	FON	N1-C2-N3	-3.49	119.95	125.42
2	A	401	T3F	O4-C4-C5	-3.48	120.87	124.90
2	A	403	T3F	N3-C2-N1	3.47	119.49	114.89
2	B	403	T3F	C5M-C5-C4	3.37	122.48	118.77
3	A	402	FON	C2-N3-C4	3.23	121.06	115.93
2	B	403	T3F	O4Q-C4Q-C3Q	-3.23	104.42	110.22
2	A	401	T3F	C5Q-C4Q-C3Q	-3.16	105.72	110.51
3	B	402	FON	C4A-C4-N3	-3.15	116.67	123.14
3	A	402	FON	O-C-N	-3.14	118.12	122.58
2	A	403	T3F	C5M-C5-C4	3.11	122.19	118.77
2	B	401	T3F	PA-O3A-PB	-3.10	122.19	132.83
2	B	401	T3F	C5M-C5-C4	3.05	122.13	118.77
2	A	401	T3F	C6-N1-C2	-2.99	118.27	121.30
2	B	403	T3F	C5-C6-N1	-2.97	120.28	123.34
2	B	403	T3F	C5M-C5-C6	-2.91	118.97	122.85
3	A	402	FON	C2-N1-C8A	2.74	120.67	114.54
2	A	401	T3F	O2-C2-N3	-2.66	116.55	121.50
2	B	401	T3F	C2'-C1'-N1	-2.66	112.36	115.61
3	A	402	FON	C11-C-N	2.60	120.88	117.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	401	T3F	O5Q-C5Q-C6Q	2.60	112.31	106.70
2	A	401	T3F	PA-O3A-PB	-2.59	123.95	132.83
2	B	403	T3F	O1A-PA-O2A	2.53	124.77	112.24
3	B	402	FON	C13-C14-N10	2.53	126.21	120.97
2	B	403	T3F	C5Q-C4Q-C3Q	2.50	114.31	110.51
2	A	403	T3F	O3A-PB-O3B	-2.46	97.53	102.48
2	A	401	T3F	O5Q-C1Q-O3B	-2.41	108.21	111.36
2	A	401	T3F	O1A-PA-O2A	2.37	123.96	112.24
3	B	402	FON	N1-C2-N3	-2.37	121.70	125.42
2	A	403	T3F	O5Q-C5Q-C6Q	2.36	111.80	106.70
2	A	401	T3F	O4'-C1'-N1	2.34	112.05	107.86
3	A	402	FON	NA2-C2-N1	2.28	120.80	117.25
2	B	403	T3F	C1'-N1-C2	2.28	122.13	117.64
2	B	403	T3F	O4'-C1'-N1	2.27	111.92	107.86
2	B	403	T3F	O5Q-C5Q-C4Q	2.23	113.52	109.52
3	B	402	FON	C2-N1-C8A	2.23	119.53	114.54
2	B	401	T3F	C2-N1-C6	2.17	123.64	117.58
2	A	401	T3F	C5-C6-N1	-2.17	121.11	123.34
3	B	402	FON	C4A-N5-C6	-2.09	115.60	119.31
2	A	403	T3F	C5-C6-N1	-2.06	121.22	123.34
3	B	402	FON	C15-C14-N10	-2.05	116.72	120.97
3	B	402	FON	NA2-C2-N1	2.03	120.41	117.25
2	A	401	T3F	O2B-PB-O1B	2.02	122.23	112.24

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	T3F	C5'-O5'-PA-O1A
2	B	401	T3F	O4'-C1'-N1-C2
3	B	402	FON	N-C-C11-C16
3	B	402	FON	O-C-C11-C12
3	B	402	FON	N-C-C11-C12
3	B	402	FON	O-C-C11-C16
2	A	403	T3F	PB-O3A-PA-O5'
2	A	401	T3F	C5'-O5'-PA-O3A
2	A	403	T3F	PA-O3A-PB-O2B
2	A	401	T3F	O4'-C1'-N1-C6
2	A	403	T3F	PA-O3A-PB-O1B
2	A	401	T3F	O4'-C1'-N1-C2
2	A	401	T3F	C2'-C1'-N1-C6
2	A	403	T3F	C5'-O5'-PA-O3A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	401	T3F	C2'-C1'-N1-C2
2	A	401	T3F	PB-O3A-PA-O2A
2	A	401	T3F	C5'-O5'-PA-O2A

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	FON	1	0
3	A	402	FON	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand T3F A 403



Ligand T3F A 401



Ligand FON B 402



Ligand FON A 402



Ligand T3F B 401



Ligand T3F B 403



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	398/405 (98%)	-0.26	1 (0%) 94 93	15, 30, 53, 70	0
1	B	398/405 (98%)	-0.47	0 100 100	15, 23, 42, 64	0
All	All	796/810 (98%)	-0.36	1 (0%) 95 95	15, 26, 49, 70	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	347	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	NA	B	405	1/1	0.85	0.12	38,38,38,38	0
3	FON	B	402	25/34	0.90	0.15	22,27,49,51	0
3	FON	A	402	25/34	0.90	0.12	27,34,44,46	0
2	T3F	B	401	34/35	0.96	0.12	19,23,25,28	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	T3F	A	401	35/35	0.96	0.12	29,35,42,43	0
2	T3F	B	403	35/35	0.97	0.10	19,23,30,32	0
4	NA	A	404	1/1	0.97	0.07	25,25,25,25	0
2	T3F	A	403	35/35	0.97	0.11	23,26,35,38	0
4	NA	B	404	1/1	0.98	0.07	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FON B 402:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FON A 402:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around T3F B 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around T3F A 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around T3F B 403:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around T3F A 403:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.