



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 19, 2024 – 04:03 PM EDT

PDB ID : 8VK8
Title : Structure of UbV.d2.1 in complex with Ube2d2
Authors : Middleton, A.J.
Deposited on : 2024-01-08
Resolution : 2.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.002 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.38.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2511 (3.00-3.00)
Clashscore	180529	2866 (3.00-3.00)
Ramachandran outliers	177936	2778 (3.00-3.00)
Sidechain outliers	177891	2781 (3.00-3.00)
RSRZ outliers	164620	2523 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	152	<div> <div>5%</div> <div> <div></div> <div>85%</div> <div>13%</div> <div>..</div> </div> </div>
1	D	152	<div> <div>5%</div> <div> <div></div> <div>82%</div> <div>15%</div> <div>.</div> </div> </div>
1	G	152	<div> <div>7%</div> <div> <div></div> <div>82%</div> <div>17%</div> <div>.</div> </div> </div>
2	B	94	<div> <div>2%</div> <div> <div></div> <div>73%</div> <div>9%</div> <div>18%</div> </div> </div>
2	C	94	<div> <div>4%</div> <div> <div></div> <div>48%</div> <div>24%</div> <div>24%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	94	
2	F	94	
2	H	94	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6499 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin-conjugating enzyme E2 D2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	151	Total	C	N	O	S	0	0	0
			1203	771	206	221	5			
1	D	147	Total	C	N	O	S	0	0	0
			1178	755	202	216	5			
1	G	150	Total	C	N	O	S	0	0	0
			1194	766	205	219	4			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP P62837
A	-3	PRO	-	expression tag	UNP P62837
A	-2	LEU	-	expression tag	UNP P62837
A	-1	GLY	-	expression tag	UNP P62837
A	0	SER	-	expression tag	UNP P62837
A	21	SER	CYS	engineered mutation	UNP P62837
A	107	SER	CYS	engineered mutation	UNP P62837
A	111	SER	CYS	engineered mutation	UNP P62837
D	-4	GLY	-	expression tag	UNP P62837
D	-3	PRO	-	expression tag	UNP P62837
D	-2	LEU	-	expression tag	UNP P62837
D	-1	GLY	-	expression tag	UNP P62837
D	0	SER	-	expression tag	UNP P62837
D	21	SER	CYS	engineered mutation	UNP P62837
D	107	SER	CYS	engineered mutation	UNP P62837
D	111	SER	CYS	engineered mutation	UNP P62837
G	-4	GLY	-	expression tag	UNP P62837
G	-3	PRO	-	expression tag	UNP P62837
G	-2	LEU	-	expression tag	UNP P62837
G	-1	GLY	-	expression tag	UNP P62837
G	0	SER	-	expression tag	UNP P62837
G	21	SER	CYS	engineered mutation	UNP P62837
G	107	SER	CYS	engineered mutation	UNP P62837

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	111	SER	CYS	engineered mutation	UNP P62837



- Molecule 2 is a protein called Ubiquitin variant D2.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	77	Total	C	N	O	S	0	0	0
			615	389	104	121	1			
2	C	71	Total	C	N	O	S	0	0	0
			567	361	94	111	1			
2	E	76	Total	C	N	O	S	0	0	0
			607	385	103	118	1			
2	F	69	Total	C	N	O		0	0	0
			547	349	90	108				
2	H	74	Total	C	N	O	S	0	0	0
			588	375	98	114	1			



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ubiquitin-conjugating enzyme E2 D2



- Molecule 1: Ubiquitin-conjugating enzyme E2 D2




- Molecule 1: Ubiquitin-conjugating enzyme E2 D2





- Molecule 2: Ubiquitin variant D2.1



- Molecule 2: Ubiquitin variant D2.1



Chain E:



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	129.77Å 129.77Å 123.11Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.50 – 2.99 41.50 – 2.99	Depositor EDS
% Data completeness (in resolution range)	99.8 (41.50-2.99) 99.8 (41.50-2.99)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 3.01Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.247 , 0.295 0.247 , 0.296	Depositor DCC
R_{free} test set	1145 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	93.0	Xtriage
Anisotropy	0.079	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 70.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.018 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6499	wwPDB-VP
Average B, all atoms (Å ²)	107.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/1240	0.48	0/1689
1	D	0.25	0/1214	0.45	0/1654
1	G	0.24	0/1231	0.48	0/1679
2	B	0.24	0/625	0.49	0/847
2	C	0.23	0/576	0.50	0/781
2	E	0.24	0/617	0.49	0/836
2	F	0.25	0/556	0.52	0/756
2	H	0.25	0/598	0.48	0/811
All	All	0.25	0/6657	0.48	0/9053

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1203	0	1191	14	0
1	D	1178	0	1165	14	0
1	G	1194	0	1182	17	0
2	B	615	0	623	5	0
2	C	567	0	574	17	0
2	E	607	0	619	17	0
2	F	547	0	554	9	0
2	H	588	0	602	18	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6499	0	6510	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 101 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:30:MET:HG3	2:H:4:PHE:HD1	1.51	0.76
2:H:1:MET:HB3	2:H:17:VAL:O	1.93	0.68
1:D:45:TYR:HA	1:D:142:THR:HG21	1.75	0.67
1:D:28:ASP:HB3	2:E:45:PHE:HE1	1.60	0.66
2:E:22:THR:HA	2:E:55:THR:HA	1.79	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/152 (98%)	145 (97%)	4 (3%)	0	100	100
1	D	145/152 (95%)	143 (99%)	2 (1%)	0	100	100
1	G	148/152 (97%)	146 (99%)	2 (1%)	0	100	100
2	B	75/94 (80%)	75 (100%)	0	0	100	100
2	C	67/94 (71%)	67 (100%)	0	0	100	100
2	E	74/94 (79%)	73 (99%)	1 (1%)	0	100	100
2	F	67/94 (71%)	62 (92%)	5 (8%)	0	100	100
2	H	72/94 (77%)	72 (100%)	0	0	100	100
All	All	797/926 (86%)	783 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/134 (100%)	131 (98%)	3 (2%)	47	76
1	D	131/134 (98%)	129 (98%)	2 (2%)	60	83
1	G	133/134 (99%)	131 (98%)	2 (2%)	60	83
2	B	70/80 (88%)	69 (99%)	1 (1%)	62	83
2	C	65/80 (81%)	61 (94%)	4 (6%)	15	45
2	E	69/80 (86%)	68 (99%)	1 (1%)	62	83
2	F	62/80 (78%)	60 (97%)	2 (3%)	34	67
2	H	67/80 (84%)	66 (98%)	1 (2%)	60	83
All	All	731/802 (91%)	715 (98%)	16 (2%)	47	76

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	144	LYS
1	G	130	ASP
1	D	15	ARG
2	F	54	ARG
2	C	58	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	151/152 (99%)	0.32	7 (4%) 38 22	51, 75, 140, 177	0
1	D	147/152 (96%)	0.64	8 (5%) 32 18	68, 98, 139, 189	0
1	G	150/152 (98%)	0.67	10 (6%) 25 14	85, 115, 152, 173	0
2	B	77/94 (81%)	-0.01	2 (2%) 57 35	50, 66, 103, 149	0
2	C	71/94 (75%)	0.69	4 (5%) 31 18	64, 114, 147, 162	0
2	E	76/94 (80%)	0.98	11 (14%) 7 4	96, 120, 150, 172	0
2	F	69/94 (73%)	1.13	11 (15%) 6 4	116, 152, 201, 232	0
2	H	74/94 (78%)	0.54	3 (4%) 42 24	87, 112, 134, 152	0
All	All	815/926 (88%)	0.59	56 (6%) 24 14	50, 106, 157, 232	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	74	TYR	4.0
1	A	18	PRO	3.7
2	B	73	LEU	3.6
1	A	90	ARG	3.6
2	B	74	ARG	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.