wwPDB X-ray Structure Validation Summary Report (i)

Sep 25, 2023 – 09:43 AM EDT

PDB ID	:	5VVC
Title	:	Structure of human endothelial nitric oxide synthase heme domain in complex with 4-(2-(((2-Amino-4-methylquinolin-7-yl)methyl)amino)ethyl)-2-methylbenzonitrile
Authors	:	Chreifi, G.; Li, H.; Poulos, T.L.
Deposited on	:	2017-05-19
Resolution	:	2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	3%; 67%; 23%; • ; 8%
1	B	440	77%; 14%; • ; 8%
1	C	440	%; 69%; 20%; • ; 9%
1	D	440	77%; 15%; 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	BTB	B	505	-	-	X	-
5	BTB	C	506	-	-	X	-

2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 13874 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, endothelial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	404	3237	2062	570	589	16	0	2	0
1	B	404	3241	2063	572	589	17	0	3	0
1	C	401	3200	2038	563	583	16	0	0	0
1	D	405	3249	2069	573	590	17	0	3	0

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0
2	B	1	43	34	1	4	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	C	1	43	34	1	4	4	0	0
			Total	C	Fe	N	O		
2	D	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	17	9	5	3	0	0
			Total	C	N	O		
3	B	1	17	9	5	3	0	0
			Total	C	N	O		
3	C	1	17	9	5	3	0	0
			Total	C	N	O		
3	D	1	17	9	5	3	0	0

- Molecule 4 is 4-(2-{[(2-amino-4-methylquinolin-7-yl)methyl]amino}ethyl)-2-methylbenzonitrile (three-letter code: 9OJ) (formula: $C_{21}H_{22}N_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	N		
4	A	1	25	21	4	0	0
			Total	C	N		
4	B	1	25	21	4	0	0
			Total	C	N		
4	C	1	25	21	4	0	0
			Total	C	N		
4	D	1	25	21	4	0	0

- Molecule 5 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: BTB) (formula: $C_8H_{19}NO_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
			Total	C	N	O		
5	B	1	14	8	1	5	0	0
			Total	C	N	O		
5	B	1	14	8	1	5	0	0
			Total	C	N	O		
5	B	1	14	8	1	5	0	0
			Total	C	N	O		
5	C	1	14	8	1	5	0	0
			Total	C	N	O		
5	C	1	14	8	1	5	0	0
			Total	C	N	O		
5	C	1	14	8	1	5	0	0
			Total	C	N	O		
5	D	1	14	8	1	5	0	0
			Total	C	N	O		
5	D	1	14	8	1	5	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
6	A	1	1	1	0	0
			Total	Zn		
6	C	1	1	1	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 6 3 3	0	0
7	C	1	Total C O 6 3 3	0	0

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total Cl 1 1	0	0
8	B	1	Total Cl 1 1	0	0
8	C	1	Total Cl 1 1	0	0
8	D	1	Total Cl 1 1	0	0

- Molecule 9 is GADOLINIUM ATOM (three-letter code: GD) (formula: Gd).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total Gd 1 1	0	0
9	B	1	Total Gd 1 1	0	0
9	C	1	Total Gd 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Gd		
9	D	1	1	1	0	0

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
10	A	89	89	89	0	0
			Total	O		
10	B	111	111	111	0	0
			Total	O		
10	C	95	95	95	0	0
			Total	O		
10	D	136	136	136	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nitric oxide synthase, endothelial

Chain A:



• Molecule 1: Nitric oxide synthase, endothelial

Chain B:



• Molecule 1: Nitric oxide synthase, endothelial

Chain C:



- Molecule 1: Nitric oxide synthase, endothelial

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	59.33Å 152.62Å 108.74Å 90.00° 90.57° 90.00°	Depositor
Resolution (Å)	88.56 – 2.40 88.56 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (88.56-2.40) 99.9 (88.56-2.40)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	0.18	Depositor
$< I/\sigma(I) >$ [1]	1.22 (at 2.40Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.206 , 0.272 0.198 , 0.267	Depositor DCC
R_{free} test set	3765 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å^2)	42.9	Xtriage
Anisotropy	0.988	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 37.0	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.109 for h,-k,-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	13874	wwPDB-VP
Average B, all atoms (Å^2)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.12% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GD, ZN, HEM, 9OJ, GOL, BTB, CL, H4B

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.37	0/3335	0.55	0/4543
1	B	0.42	0/3339	0.55	0/4548
1	C	0.39	0/3292	0.55	0/4487
1	D	0.45	0/3347	0.58	1/4559 (0.0%)
All	All	0.41	0/13313	0.56	1/18137 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	336	VAL	CB-CA-C	-5.08	101.76	111.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3237	0	3146	69	0
1	B	3241	0	3152	41	0
1	C	3200	0	3098	64	0
1	D	3249	0	3163	38	0
2	A	43	0	30	4	0
2	B	43	0	30	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	43	0	30	3	0
2	D	43	0	30	3	0
3	A	17	0	15	1	0
3	B	17	0	15	0	0
3	C	17	0	15	1	0
3	D	17	0	15	1	0
4	A	25	0	0	2	0
4	B	25	0	0	0	0
4	C	25	0	0	1	0
4	D	25	0	0	0	0
5	A	42	0	56	10	0
5	B	42	0	54	12	0
5	C	42	0	57	12	0
5	D	28	0	36	6	0
6	A	1	0	0	0	0
6	C	1	0	0	0	0
7	A	6	0	8	1	0
7	C	6	0	8	0	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	1	0
9	A	1	0	0	0	0
9	B	1	0	0	0	0
9	C	1	0	0	0	0
9	D	1	0	0	0	0
10	A	89	0	0	4	0
10	B	111	0	0	4	0
10	C	95	0	0	10	0
10	D	136	0	0	2	0
All	All	13874	0	12958	234	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 234 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:ILE:HG13	1:A:140:ARG:HG3	1.66	0.77
1:A:298:GLU:HG3	1:A:299:PRO:HD2	1.67	0.74
1:C:156:GLU:OE2	1:C:164:GLN:N	2.22	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:147:GLU:HA	1:C:150:LEU:HD12	1.73	0.70
1:A:298:GLU:OE1	5:A:506:BTB:O4	2.10	0.69

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/440 (91%)	371 (92%)	28 (7%)	3 (1%)	22	32
1	B	403/440 (92%)	389 (96%)	14 (4%)	0	100	100
1	C	397/440 (90%)	371 (94%)	24 (6%)	2 (0%)	29	41
1	D	404/440 (92%)	390 (96%)	13 (3%)	1 (0%)	47	62
All	All	1606/1760 (91%)	1521 (95%)	79 (5%)	6 (0%)	34	48

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	90	GLN
1	A	283	ASN
1	C	89	GLN
1	A	106	PRO
1	C	154	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/373 (92%)	326 (94%)	19 (6%)	21	35
1	B	346/373 (93%)	334 (96%)	12 (4%)	36	55
1	C	340/373 (91%)	320 (94%)	20 (6%)	19	32
1	D	347/373 (93%)	340 (98%)	7 (2%)	55	74
All	All	1378/1492 (92%)	1320 (96%)	58 (4%)	31	47

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	326	LEU
1	D	124	LEU
1	C	102	SER
1	D	109	LEU
1	C	391	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	257	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 35 ligands modelled in this entry, 10 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	9OJ	C	503	-	27,27,27	2.33	1 (3%)	35,37,37	1.50	7 (20%)
5	BTB	C	506	-	13,13,13	0.58	0	7,16,16	0.88	0
5	BTB	D	504	9	13,13,13	0.43	0	7,16,16	0.48	0
2	HEM	D	501	1	41,50,50	1.94	7 (17%)	45,82,82	2.12	13 (28%)
5	BTB	B	508	9	13,13,13	0.43	0	7,16,16	0.45	0
2	HEM	B	501	1	41,50,50	1.88	8 (19%)	45,82,82	2.08	16 (35%)
5	BTB	C	505	-	13,13,13	0.35	0	7,16,16	0.38	0
3	H4B	A	502	-	16,18,18	0.93	0	11,26,26	2.71	6 (54%)
2	HEM	C	501	1	41,50,50	2.03	9 (21%)	45,82,82	1.92	10 (22%)
5	BTB	B	504	9	13,13,13	0.49	0	7,16,16	0.36	0
3	H4B	D	502	-	16,18,18	0.78	0	11,26,26	2.72	6 (54%)
7	GOL	A	508	-	5,5,5	0.37	0	5,5,5	0.28	0
3	H4B	B	502	-	16,18,18	0.99	1 (6%)	11,26,26	2.86	6 (54%)
3	H4B	C	502	-	16,18,18	0.93	0	11,26,26	2.63	6 (54%)
4	9OJ	D	503	-	27,27,27	2.14	1 (3%)	35,37,37	1.23	4 (11%)
5	BTB	C	504	-	13,13,13	0.49	0	7,16,16	0.73	0
5	BTB	A	504	9	13,13,13	0.40	0	7,16,16	0.47	0
7	GOL	C	508	-	5,5,5	0.39	0	5,5,5	0.37	0
4	9OJ	A	503	-	27,27,27	2.09	1 (3%)	35,37,37	1.21	4 (11%)
5	BTB	A	506	-	13,13,13	0.35	0	7,16,16	0.48	0
4	9OJ	B	503	-	27,27,27	2.14	2 (7%)	35,37,37	1.41	7 (20%)
5	BTB	A	505	-	13,13,13	0.38	0	7,16,16	0.55	0
5	BTB	B	505	-	13,13,13	0.44	0	7,16,16	0.46	0
2	HEM	A	501	1	41,50,50	1.83	6 (14%)	45,82,82	1.62	9 (20%)
5	BTB	D	505	-	13,13,13	0.51	0	7,16,16	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	9OJ	C	503	-	-	1/9/9/9	0/3/3/3
5	BTB	C	506	-	-	11/21/21/21	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BTB	D	504	9	-	2/21/21/21	-
2	HEM	D	501	1	-	1/12/54/54	-
5	BTB	B	508	9	-	5/21/21/21	-
2	HEM	B	501	1	-	1/12/54/54	-
5	BTB	C	505	-	-	6/21/21/21	-
3	H4B	A	502	-	-	1/8/17/17	0/2/2/2
2	HEM	C	501	1	-	2/12/54/54	-
5	BTB	B	504	9	-	3/21/21/21	-
3	H4B	D	502	-	-	0/8/17/17	0/2/2/2
7	GOL	A	508	-	-	4/4/4/4	-
3	H4B	B	502	-	-	3/8/17/17	0/2/2/2
3	H4B	C	502	-	-	0/8/17/17	0/2/2/2
4	9OJ	D	503	-	-	1/9/9/9	0/3/3/3
5	BTB	C	504	-	-	5/21/21/21	-
5	BTB	A	504	9	-	6/21/21/21	-
7	GOL	C	508	-	-	2/4/4/4	-
4	9OJ	A	503	-	-	3/9/9/9	0/3/3/3
5	BTB	A	506	-	-	7/21/21/21	-
4	9OJ	B	503	-	-	1/9/9/9	0/3/3/3
5	BTB	A	505	-	-	7/21/21/21	-
5	BTB	B	505	-	-	9/21/21/21	-
2	HEM	A	501	1	-	3/12/54/54	-
5	BTB	D	505	-	-	11/21/21/21	-

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	503	9OJ	C23-C24	-11.12	1.27	1.44
4	D	503	9OJ	C23-C24	-10.20	1.29	1.44
4	B	503	9OJ	C23-C24	-9.94	1.29	1.44
4	A	503	9OJ	C23-C24	-9.76	1.29	1.44
2	D	501	HEM	C3D-C2D	7.75	1.53	1.36

The worst 5 of 94 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	501	HEM	C4D-ND-C1D	6.63	111.93	105.07
2	D	501	HEM	C4D-ND-C1D	6.40	111.68	105.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	D	502	H4B	C8A-C4A-C4	6.37	120.23	114.57
2	C	501	HEM	C4D-ND-C1D	6.37	111.65	105.07
2	D	501	HEM	CBA-CAA-C2A	-5.85	102.63	112.62

There are no chirality outliers.

5 of 95 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	HEM	C1A-C2A-CAA-CBA
2	A	501	HEM	C3A-C2A-CAA-CBA
5	A	504	BTB	O1-C1-C2-C3
5	A	504	BTB	O1-C1-C2-C4
5	A	504	BTB	O1-C1-C2-N

There are no ring outliers.

20 monomers are involved in 57 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	503	9OJ	1	0
5	C	506	BTB	8	0
5	D	504	BTB	4	0
2	D	501	HEM	3	0
5	B	508	BTB	5	0
2	B	501	HEM	1	0
5	C	505	BTB	2	0
3	A	502	H4B	1	0
2	C	501	HEM	3	0
3	D	502	H4B	1	0
7	A	508	GOL	1	0
3	C	502	H4B	1	0
5	C	504	BTB	2	0
5	A	504	BTB	3	0
4	A	503	9OJ	2	0
5	A	506	BTB	3	0
5	A	505	BTB	4	0
5	B	505	BTB	7	0
2	A	501	HEM	4	0
5	D	505	BTB	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 9OJ C 503



Ligand HEM D 501



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM B 501



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM C 501



Bond lengths



Bond angles



Torsions



Rings

Ligand 9OJ D 503



Ligand 9OJ A 503



Ligand 9OJ B 503



Bond lengths



Bond angles



Torsions



Rings

Ligand HEM A 501



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	404/440 (91%)	-0.03	13 (3%)	47	46	33, 63, 110, 141	0
1	B	404/440 (91%)	-0.33	1 (0%)	95	94	27, 47, 79, 122	0
1	C	401/440 (91%)	-0.18	3 (0%)	87	86	31, 57, 99, 129	0
1	D	405/440 (92%)	-0.32	1 (0%)	95	94	27, 46, 75, 118	0
All	All	1614/1760 (91%)	-0.22	18 (1%)	80	79	27, 52, 97, 141	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	119	ALA	5.0
1	A	144	GLN	4.8
1	A	480	TRP	3.8
1	A	163	TYR	3.1
1	A	109	LEU	2.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
5	BTB	C	506	14/14	0.83	0.15	83,88,90,92	0
5	BTB	C	505	14/14	0.87	0.17	100,106,109,110	0
5	BTB	A	506	14/14	0.87	0.16	114,120,123,123	0
5	BTB	D	504	14/14	0.89	0.12	50,69,74,75	0
7	GOL	A	508	6/6	0.89	0.17	74,81,86,87	0
7	GOL	C	508	6/6	0.90	0.22	73,76,78,79	0
5	BTB	B	505	14/14	0.91	0.22	63,68,77,81	0
5	BTB	B	504	14/14	0.91	0.14	32,58,67,70	0
5	BTB	B	508	14/14	0.93	0.18	47,76,83,83	0
5	BTB	A	505	14/14	0.93	0.14	81,90,93,94	0
5	BTB	C	504	14/14	0.94	0.21	49,66,73,74	0
5	BTB	D	505	14/14	0.94	0.20	67,79,81,83	0
9	GD	A	510	1/1	0.94	0.11	137,137,137,137	0
4	9OJ	A	503	25/25	0.95	0.17	39,51,63,68	0
4	9OJ	D	503	25/25	0.95	0.16	31,48,60,62	0
5	BTB	A	504	14/14	0.95	0.17	89,93,95,95	0
3	H4B	A	502	17/17	0.95	0.12	47,55,58,60	0
4	9OJ	C	503	25/25	0.96	0.14	47,54,58,61	0
4	9OJ	B	503	25/25	0.96	0.17	36,44,53,60	0
8	CL	A	509	1/1	0.97	0.12	64,64,64,64	0
3	H4B	C	502	17/17	0.97	0.11	44,54,59,59	0
2	HEM	D	501	43/43	0.98	0.16	30,39,48,52	0
2	HEM	A	501	43/43	0.98	0.17	39,52,60,62	0
3	H4B	B	502	17/17	0.98	0.11	28,41,44,45	0
2	HEM	B	501	43/43	0.98	0.14	31,44,53,59	0
3	H4B	D	502	17/17	0.98	0.12	30,37,40,43	0
8	CL	B	506	1/1	0.98	0.24	67,67,67,67	0
8	CL	C	509	1/1	0.98	0.09	67,67,67,67	0
2	HEM	C	501	43/43	0.98	0.13	31,47,59,63	0
8	CL	D	507	1/1	0.99	0.04	53,53,53,53	0
6	ZN	C	507	1/1	0.99	0.14	48,48,48,48	0
9	GD	B	507	1/1	0.99	0.20	55,55,55,55	0
9	GD	C	510	1/1	0.99	0.15	119,119,119,119	0
6	ZN	A	507	1/1	1.00	0.13	51,51,51,51	0
9	GD	D	506	1/1	1.00	0.19	53,53,53,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 9OJ A 503:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 9OJ D 503:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 9OJ C 503:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around 9OJ B 503:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around HEM D 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around HEM A 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around HEM B 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM C 501:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.