wwPDB NMR Structure Validation Summary Report (i)

Feb 22, 2022 – 09:20 AM EST

PDB ID	:	1WEM
Title	:	Solution structure of PHD domain in death inducer-obliterator 1(DIO-1)
Authors	:	He, F.; Muto, Y.; Inoue, M.; Kigawa, T.; Shirouzu, M.; Terada, T.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on	:	2004-05-25

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.26
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	76	47% 22% • 28%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:16-A:70 (55)	0.28	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 7, 8, 9, 10, 12, 14, 17, 18, 19, 20
2	6, 11, 16
3	13, 15

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1101 atoms, of which 520 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Death associated transcription factor 1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	76	1099	348	520	105	116	10	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	cloning artifact	UNP Q8C9B9
A	2	SER	-	cloning artifact	UNP Q8C9B9
A	3	SER	-	cloning artifact	UNP Q8C9B9
A	4	GLY	-	cloning artifact	UNP Q8C9B9
A	5	SER	-	cloning artifact	UNP Q8C9B9
A	6	SER	-	cloning artifact	UNP Q8C9B9
A	7	GLY	-	cloning artifact	UNP Q8C9B9
A	71	SER	-	cloning artifact	UNP Q8C9B9
A	72	GLY	-	cloning artifact	UNP Q8C9B9
A	73	PRO	-	cloning artifact	UNP Q8C9B9
A	74	SER	-	cloning artifact	UNP Q8C9B9
A	75	SER	-	cloning artifact	UNP Q8C9B9
A	76	GLY	-	cloning artifact	UNP Q8C9B9

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	
			Total	Zn
2	A	2	2	2

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Death associated transcription factor 1

Chain A: 47% 22% • 28%

G1 S2 S3 G4 S5 S6 G7 E8 C9 E10 V11 Y12 D13 P14 N15 R22 N27 R28 F29 M30 I31 E38 W39 F40 H41 G42 G46 I47 G52 R53 L54 L55 E56 R57 Y62 I63 C64 I69 L70 S71 G72 P73 S74 S75 G76

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 10. Colouring as in section 4.1 above.

- Molecule 1: Death associated transcription factor 1

Chain A: 41% 25% 7% 28%

G1 S2 S3 G4 S5 S6 G7 E8 C9 E10 V11 Y12 D13 P14 N15 A16 L17 R22 H25 N26 N27 R28 F29 M30 I31 E38 W39 F40 H41 G42 D43 C44 V45 G46 I47 L54 L55 E56 R57 D61 Y62 I63 C64 P65 N66 C67 T68 I69 L70 S71 G72 P73 S74 S75 G76

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy, target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.0.17
CYANA	refinement	2.0.17

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	444	411	410	19±4
All	All	8920	8220	8200	377

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

5 of 115 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:52:GLY:HA2	1:A:55:LEU:HD23	0.78	1.52	9	4
1:A:55:LEU:C	1:A:55:LEU:HD12	0.76	2.00	7	4
1:A:52:GLY:HA2	1:A:55:LEU:HD12	0.73	1.60	12	7
1:A:55:LEU:HD13	1:A:62:TYR:HB3	0.69	1.63	4	3
1:A:51:ARG:HH11	1:A:55:LEU:HD11	0.69	1.46	1	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	55/76 (72%)	38±2 (70±3%)	15±2 (27±3%)	2±1 (4±2%)	5 32
All	All	1100/1520 (72%)	765 (70%)	292 (27%)	43 (4%)	5 32

5 of 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	55	LEU	13
1	A	28	ARG	11
1	A	47	ILE	7
1	A	33	CYS	3
1	A	69	ILE	3

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	49/65 (75%)	44±1 (90±3%)	5±1 (10±3%)	12 58
All	All	980/1300 (75%)	886 (90%)	94 (10%)	12 58

5 of 18 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	31	ILE	20
1	A	64	CYS	20
1	A	22	ARG	9
1	A	57	ARG	6
1	A	35	ARG	5

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided