wwPDB EM Validation Summary Report (i)

Oct 5, 2026 – 12:49 PM JST

PDB ID	:	9X4L / pdb_00009x4l
EMDB ID	:	EMD-66552
Title	:	Cdc45 and Sld3 Loading on MCM Double Hexamer
Authors	:	Yu, D.; Huo, Y.; Dang, S.; Zhai, Y.
Deposited on	:	2025-10-11
Resolution	:	3.50 Å(reported)
Based on initial model	:	.

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev133
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4-5-2 with Phenix2.2
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13950 (3.00 - 4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	2	868	63% 9% 28%
1	B	868	63% 9% 28%
2	3	971	59% 6% 35%
2	C	971	59% 6% 35%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	4	933	62% 9% 28%
3	D	933	61% 11% 28%
4	5	775	74% 11% 15%
4	E	775	76% 8% 15%
5	6	1017	55% 8% 36%
5	F	1017	54% 9% 37%
6	7	845	74% 8% 17%
6	G	845	73% 9% 18%
7	J	650	5% 67% 21% 12%
8	K	668	29% 8% 63%
8	X	668	98%

2 Entry composition (i)

There are 12 unique types of molecules in this entry. The entry contains 69053 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA replication licensing factor MCM2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	2	628	4970	3133	883	935	19	0	0
1	B	628	4970	3133	883	935	19	0	0

- Molecule 2 is a protein called DNA replication licensing factor MCM3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	3	636	4983	3140	888	942	13	0	0
2	C	636	4978	3137	888	940	13	0	0

- Molecule 3 is a protein called DNA replication licensing factor MCM4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	4	669	5316	3334	917	1035	30	0	0
3	D	670	5323	3339	918	1036	30	0	0

- Molecule 4 is a protein called Minichromosome maintenance protein 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	5	658	5145	3228	882	1010	25	0	0
4	E	657	5137	3222	881	1009	25	0	0

- Molecule 5 is a protein called DNA replication licensing factor MCM6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	6	650	5143	3236	898	983	26	0	0
5	F	638	5052	3184	884	959	25	0	0

- Molecule 6 is a protein called DNA replication licensing factor MCM7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	7	699	5505	3463	952	1059	31	0	0
6	G	694	5467	3440	947	1049	31	0	0

- Molecule 7 is a protein called Cell division control protein 45.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	J	571	4635	2956	780	885	14	0	0

- Molecule 8 is a protein called DNA replication regulator SLD3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	K	245	2010	1301	333	369	7	0	0
8	X	13	105	64	21	19	1	0	0

- Molecule 9 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).

ATP



Mol	Chain	Residues	Atoms					AltConf
9	2	1	Total 31	C 10	N 5	O 13	P 3	0
9	4	1	Total 31	C 10	N 5	O 13	P 3	0
9	7	1	Total 31	C 10	N 5	O 13	P 3	0
9	B	1	Total 31	C 10	N 5	O 13	P 3	0
9	D	1	Total 31	C 10	N 5	O 13	P 3	0
9	G	1	Total 31	C 10	N 5	O 13	P 3	0

- Molecule 10 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
10	2	1	Total 1	Mg 1	0
10	3	1	Total 1	Mg 1	0
10	4	1	Total 1	Mg 1	0
10	5	1	Total 1	Mg 1	0
10	7	1	Total 1	Mg 1	0
10	B	1	Total 1	Mg 1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
			Total	Mg	
10	C	1	1	1	0
10	D	1	1	1	0
10	E	1	1	1	0
10	G	1	1	1	0

- Molecule 11 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
			Total	Zn	
11	2	1	1	1	0
11	4	1	1	1	0
11	5	1	1	1	0
11	6	1	1	1	0
11	7	1	1	1	0
11	B	1	1	1	0
11	D	1	1	1	0
11	E	1	1	1	0
11	F	1	1	1	0
11	G	1	1	1	0

- Molecule 12 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					AltConf
12	3	1	Total	C	N	O	P	0
			27	10	5	10	2	
12	5	1	Total	C	N	O	P	0
			27	10	5	10	2	
12	C	1	Total	C	N	O	P	0
			27	10	5	10	2	
12	E	1	Total	C	N	O	P	0
			27	10	5	10	2	

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA replication licensing factor MCM2

Chain 2: 63% 9% 28%



• Molecule 1: DNA replication licensing factor MCM2

Chain B: 63% 9% 28%





• Molecule 2: DNA replication licensing factor MCM3

Chain 3:



• Molecule 2: DNA replication licensing factor MCM3

Chain C:



• Molecule 3: DNA replication licensing factor MCM4

Chain 4:



• Molecule 3: DNA replication licensing factor MCM4

Chain D:



• Molecule 4: Minichromosome maintenance protein 5

Chain 5:





• Molecule 4: Minichromosome maintenance protein 5

Chain E:



• Molecule 5: DNA replication licensing factor MCM6

Chain 6:





• Molecule 5: DNA replication licensing factor MCM6

Chain F:



• Molecule 6: DNA replication licensing factor MCM7

Chain 7:



• Molecule 6: DNA replication licensing factor MCM7

Chain G:



• Molecule 7: Cell division control protein 45

Chain J:





- Molecule 8: DNA replication regulator SLD3

Chain K:



- Molecule 8: DNA replication regulator SLD3

Chain X: 98%



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	34886	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 BIOCONTINUUM (6k x 4k)	Depositor
Maximum map value	0.835	Depositor
Minimum map value	-0.287	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.033	Depositor
Recommended contour level	0.15	Depositor
Map size (Å)	423.99997, 423.99997, 423.99997	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.06, 1.06, 1.06	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, ZN, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	2	0.14	0/5057	0.36	1/6834 (0.0%)
1	B	0.14	0/5057	0.37	0/6834
2	3	0.15	0/5069	0.33	0/6875
2	C	0.15	0/5064	0.32	0/6868
3	4	0.15	0/5392	0.34	1/7289 (0.0%)
3	D	0.14	0/5400	0.35	0/7300
4	5	0.15	0/5219	0.34	0/7058
4	E	0.15	0/5211	0.35	0/7047
5	6	0.16	0/5223	0.39	4/7046 (0.1%)
5	F	0.14	0/5132	0.33	0/6921
6	7	0.16	0/5590	0.34	0/7554
6	G	0.16	0/5550	0.34	0/7496
7	J	0.13	0/4721	0.38	2/6390 (0.0%)
8	K	0.13	0/2047	0.42	0/2755
8	X	0.20	0/106	0.46	0/141
All	All	0.15	0/69838	0.35	8/94408 (0.0%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
5	6	333	CYS	CB-CA-C	5.72	117.62	109.38
3	4	421	ASP	CB-CA-C	-5.67	110.06	116.63
5	6	332	PHE	CA-C-N	-5.62	113.81	121.61
5	6	332	PHE	C-N-CA	-5.62	113.81	121.61
7	J	598	LYS	CA-C-N	5.51	130.42	121.83

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	2	4970	0	5026	51	0
1	B	4970	0	5026	60	0
2	3	4983	0	5046	36	0
2	C	4978	0	5041	36	0
3	4	5316	0	5365	57	0
3	D	5323	0	5373	64	0
4	5	5145	0	5199	59	0
4	E	5137	0	5188	41	0
5	6	5143	0	5168	59	0
5	F	5052	0	5095	68	0
6	7	5505	0	5565	43	0
6	G	5467	0	5531	57	0
7	J	4635	0	4622	97	0
8	K	2010	0	2081	36	0
8	X	105	0	98	1	0
9	2	31	0	12	1	0
9	4	31	0	12	2	0
9	7	31	0	12	0	0
9	B	31	0	12	0	0
9	D	31	0	12	0	0
9	G	31	0	12	0	0
10	2	1	0	0	0	0
10	3	1	0	0	0	0
10	4	1	0	0	0	0
10	5	1	0	0	0	0
10	7	1	0	0	0	0
10	B	1	0	0	0	0
10	C	1	0	0	0	0
10	D	1	0	0	0	0
10	E	1	0	0	0	0
10	G	1	0	0	0	0
11	2	1	0	0	0	0
11	4	1	0	0	0	0
11	5	1	0	0	0	0
11	6	1	0	0	0	0
11	7	1	0	0	0	0
11	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	D	1	0	0	0	0
11	E	1	0	0	0	0
11	F	1	0	0	0	0
11	G	1	0	0	0	0
12	3	27	0	12	1	0
12	5	27	0	12	4	0
12	C	27	0	12	0	0
12	E	27	0	12	0	0
All	All	69053	0	69544	713	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 713 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:6:311:CYS:HB3	5:6:314:CYS:HB2	1.63	0.81
5:F:379:VAL:HG23	5:F:454:PHE:HB3	1.62	0.80
5:6:644:MET:HE3	5:6:648:ASP:HB2	1.63	0.79
5:6:314:CYS:SG	5:F:315:ARG:NH1	2.56	0.79
3:D:347:PHE:HB3	3:D:382:MET:HE2	1.64	0.79

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	2	622/868 (72%)	607 (98%)	15 (2%)	0	100	100
1	B	622/868 (72%)	598 (96%)	24 (4%)	0	100	100
2	3	626/971 (64%)	612 (98%)	14 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	626/971 (64%)	617 (99%)	9 (1%)	0	100	100
3	4	663/933 (71%)	646 (97%)	17 (3%)	0	100	100
3	D	664/933 (71%)	648 (98%)	16 (2%)	0	100	100
4	5	650/775 (84%)	638 (98%)	12 (2%)	0	100	100
4	E	649/775 (84%)	638 (98%)	11 (2%)	0	100	100
5	6	640/1017 (63%)	622 (97%)	18 (3%)	0	100	100
5	F	628/1017 (62%)	612 (98%)	16 (2%)	0	100	100
6	7	693/845 (82%)	684 (99%)	9 (1%)	0	100	100
6	G	686/845 (81%)	679 (99%)	7 (1%)	0	100	100
7	J	563/650 (87%)	550 (98%)	13 (2%)	0	100	100
8	K	241/668 (36%)	231 (96%)	10 (4%)	0	100	100
8	X	11/668 (2%)	10 (91%)	1 (9%)	0	100	100
All	All	8584/12804 (67%)	8392 (98%)	192 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	2	552/770 (72%)	550 (100%)	2 (0%)	84	81
1	B	552/770 (72%)	548 (99%)	4 (1%)	76	78
2	3	550/835 (66%)	545 (99%)	5 (1%)	70	76
2	C	549/835 (66%)	543 (99%)	6 (1%)	65	74
3	4	605/848 (71%)	600 (99%)	5 (1%)	73	77
3	D	606/848 (72%)	600 (99%)	6 (1%)	68	75
4	5	588/688 (86%)	584 (99%)	4 (1%)	76	78
4	E	587/688 (85%)	586 (100%)	1 (0%)	87	85
5	6	568/886 (64%)	563 (99%)	5 (1%)	70	76

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
5	F	559/886 (63%)	557 (100%)	2 (0%)	84	81
6	7	618/753 (82%)	615 (100%)	3 (0%)	81	80
6	G	613/753 (81%)	611 (100%)	2 (0%)	86	83
7	J	517/586 (88%)	515 (100%)	2 (0%)	84	81
8	K	229/628 (36%)	227 (99%)	2 (1%)	70	76
8	X	12/628 (2%)	12 (100%)	0	100	100
All	All	7705/11402 (68%)	7656 (99%)	49 (1%)	76	79

5 of 49 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	220	THR
3	D	356	MET
2	C	384	MET
2	C	533	ILE
3	D	427	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
6	7	107	GLN
6	G	90	ASN
1	B	665	GLN
5	F	833	GLN
7	J	101	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 30 ligands modelled in this entry, 20 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
12	ADP	C	2001	10	27,29,29	1.34	4 (14%)	42,45,45	1.96	10 (23%)
9	ATP	7	2001	10	29,33,33	0.31	0	44,52,52	0.50	0
12	ADP	3	2001	10	27,29,29	1.36	4 (14%)	42,45,45	1.95	10 (23%)
12	ADP	E	2001	10	27,29,29	1.36	4 (14%)	42,45,45	1.95	11 (26%)
9	ATP	D	2001	10	29,33,33	0.32	0	44,52,52	0.52	0
9	ATP	4	2001	10	29,33,33	0.29	0	44,52,52	0.46	1 (2%)
12	ADP	5	2001	10	27,29,29	1.33	4 (14%)	42,45,45	2.00	10 (23%)
9	ATP	G	2001	10	29,33,33	0.32	0	44,52,52	0.51	0
9	ATP	2	2001	10	29,33,33	0.29	0	44,52,52	0.48	1 (2%)
9	ATP	B	2001	10	29,33,33	0.30	0	44,52,52	0.47	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	ADP	C	2001	10	-	3/16/32/32	0/3/3/3
9	ATP	7	2001	10	-	4/22/38/38	0/3/3/3
12	ADP	3	2001	10	-	5/16/32/32	0/3/3/3
12	ADP	E	2001	10	-	6/16/32/32	0/3/3/3
9	ATP	D	2001	10	-	1/22/38/38	0/3/3/3
9	ATP	4	2001	10	-	3/22/38/38	0/3/3/3
12	ADP	5	2001	10	-	3/16/32/32	0/3/3/3
9	ATP	G	2001	10	-	3/22/38/38	0/3/3/3
9	ATP	2	2001	10	-	9/22/38/38	0/3/3/3
9	ATP	B	2001	10	-	5/22/38/38	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	3	2001	ADP	C5-C4	4.54	1.47	1.39
12	E	2001	ADP	C5-C4	4.46	1.47	1.39
12	C	2001	ADP	C5-C4	4.43	1.47	1.39
12	5	2001	ADP	C5-C4	4.42	1.47	1.39
12	E	2001	ADP	C5-C6	2.69	1.48	1.41

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
12	5	2001	ADP	C5-C4-N3	-6.28	118.56	126.75
12	E	2001	ADP	C5-C4-N3	-6.20	118.67	126.75
12	3	2001	ADP	C5-C4-N3	-6.15	118.73	126.75
12	C	2001	ADP	C5-C4-N3	-6.11	118.78	126.75
12	5	2001	ADP	N3-C4-N9	5.00	135.31	127.08

There are no chirality outliers.

5 of 42 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	2	2001	ATP	C5'-O5'-PA-O1A
9	2	2001	ATP	C5'-O5'-PA-O2A
9	B	2001	ATP	C5'-O5'-PA-O1A
9	B	2001	ATP	C5'-O5'-PA-O2A
9	B	2001	ATP	C5'-O5'-PA-O3A

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	3	2001	ADP	1	0
9	4	2001	ATP	2	0
12	5	2001	ADP	4	0
9	2	2001	ATP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand ADP C 2001



Ligand ATP 7 2001



Ligand ADP 3 2001



Ligand ADP E 2001



Ligand ATP D 2001



Ligand ATP 4 2001



Ligand ADP 5 2001



Ligand ATP G 2001



Ligand ATP 2 2001





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-66552. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map


X Index: 200


Y Index: 200


Z Index: 200

6.2.2 Raw map


X Index: 200


Y Index: 200


Z Index: 200

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 200



Y Index: 192



Z Index: 230

6.3.2 Raw map



X Index: 0



Y Index: 0



Z Index: 0

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map


X


Y


Z

6.4.2 Raw map


X


Y


Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map

X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.15. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 717 nm^3; this corresponds to an approximate mass of 648 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.286 Å^{-1}

8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.286 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	3.50	3.96	3.54
Unmasked-calculated*	6.68	8.98	7.09

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.68 differs from the reported value 3.5 by more than 10 %

9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-66552 and PDB model 9X4L. Per-residue inclusion information can be found in section 3 on page 9.

9.1 Map-model overlay (i)



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.15 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.15).

9.4 Atom inclusion (i)



At the recommended contour level, 97% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.15) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8530	0.4010
2	0.8390	0.3960
3	0.8870	0.4390
4	0.8700	0.4250
5	0.8430	0.4130
6	0.8310	0.3950
7	0.8850	0.4370
B	0.8430	0.3900
C	0.8800	0.4370
D	0.8660	0.4190
E	0.8520	0.4110
F	0.8260	0.3890
G	0.8830	0.4240
J	0.7860	0.2800
K	0.8170	0.2400
X	0.6020	0.3910