Full wwPDB X-ray Structure Validation Report (i)

Oct 26, 2026 – 10:39 PM JST

PDB ID	:	9XCL / pdb_00009xcl
Title	:	Crystal structure of 3-deoxy-8-phosphooctulonate synthase from Acinetobacter baumannii
Authors	:	Cheng, Q.; Chen, S.
Deposited on	:	2025-10-26
Resolution	:	1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	92% 7% ••
1	B	285	91% 8% •
1	C	285	11% 80% 15% • •
1	D	285	9% 87% 8% • 5%

2 Entry composition


There are 3 unique types of molecules in this entry. The entry contains 9561 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-dehydro-3-deoxyphosphooctonate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	0	0
			2204	1409	381	402	12			
1	B	283	Total	C	N	O	S	0	0	0
			2204	1409	381	402	12			
1	C	274	Total	C	N	O	S	0	0	0
			2149	1379	369	389	12			
1	D	272	Total	C	N	O	S	0	0	0
			2133	1371	364	386	12			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O_4S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	O	S		0	0
			5	4	1			
2	A	1	Total	O	S		0	0
			5	4	1			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	226	Total O 226 226	0	0
3	B	209	Total O 209 209	0	0
3	C	192	Total O 192 192	0	0
3	D	214	Total O 214 214	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase

Chain A:




• Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase

Chain B:




• Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase

Chain C:



• Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	81.58Å 89.12Å 92.68Å 90.00° 114.98° 90.00°	Depositor
Resolution (Å)	33.70 – 1.71 33.70 – 1.71	Depositor EDS
% Data completeness (in resolution range)	99.8 (33.70-1.71) 99.8 (33.70-1.71)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.74 (at 1.71Å)	Xtriage
Refinement program	REFMAC 5.8.0430	Depositor
R, R_{free}	0.177 , 0.209 0.189 , 0.217	Depositor DCC
R_{free} test set	6547 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å^2)	16.5	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.39 , 35.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.51$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	0.015 for h,-k,-h-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	9561	wwPDB-VP
Average B, all atoms (Å^2)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.66% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.68	0/2250	1.08	4/3038 (0.1%)
1	B	0.67	0/2250	1.07	5/3038 (0.2%)
1	C	0.66	0/2193	1.14	7/2960 (0.2%)
1	D	0.65	0/2177	1.09	4/2941 (0.1%)
All	All	0.66	0/8870	1.10	20/11977 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	1
All	All	0	5

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	99	ASP	CA-CB-CG	8.08	120.68	112.60
1	C	239	PHE	CB-CA-C	-7.34	97.02	109.72
1	A	99	ASP	CA-CB-CG	7.34	119.94	112.60
1	C	193	GLU	CB-CG-CD	6.87	124.27	112.60
1	B	99	ASP	CA-CB-CG	6.81	119.41	112.60
1	A	208	THR	OG1-CB-CG2	-6.60	96.10	109.30
1	D	99	ASP	CA-CB-CG	6.21	118.81	112.60
1	D	239	PHE	CB-CA-C	6.18	120.41	109.72
1	C	285	HIS	CA-CB-CG	6.03	119.83	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	208	THR	OG1-CB-CG2	-5.96	97.37	109.30
1	B	45	TYR	N-CA-CB	5.59	118.12	110.01
1	C	248	LYS	CB-CA-C	-5.58	101.12	110.56
1	A	214	ASP	CA-CB-CG	5.44	118.04	112.60
1	D	239	PHE	CA-CB-CG	5.36	119.16	113.80
1	D	36	ASP	CA-CB-CG	5.35	117.95	112.60
1	A	143	LYS	CG-CD-CE	5.34	123.58	111.30
1	C	125	GLN	N-CA-CB	5.12	117.67	110.24
1	B	201	ASP	CA-CB-CG	5.12	117.72	112.60
1	B	207	GLN	N-CA-CB	5.11	117.55	110.04
1	C	258	ARG	N-CA-CB	-5.04	102.56	109.97

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	124	ARG	Sidechain
1	A	228	ARG	Sidechain
1	B	124	ARG	Sidechain
1	B	51	ARG	Sidechain
1	C	220	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2204	0	2231	11	0
1	B	2204	0	2231	11	0
1	C	2149	0	2182	23	0
1	D	2133	0	2166	17	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
3	A	226	0	0	2	0
3	B	209	0	0	0	0
3	C	192	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	214	0	0	1	0
All	All	9561	0	8810	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:207:GLN:H	1:D:222:GLN:HE22	1.37	0.73
1:C:207:GLN:H	1:C:222:GLN:HE22	1.47	0.62
1:A:8:GLU:HG3	3:A:434:HOH:O	1.99	0.61
1:C:206:LEU:HD12	1:C:255:SER:OG	2.01	0.60
1:D:45:TYR:CD1	1:D:262:LEU:HD21	2.37	0.59
1:B:143:LYS:CE	1:B:171:GLU:OE1	2.52	0.58
1:A:222:GLN:HE21	1:B:189:ASP:HB2	1.67	0.57
1:B:143:LYS:HE3	1:B:171:GLU:OE1	2.06	0.56
1:C:226:LEU:HG	1:D:186:LEU:HD21	1.86	0.56
1:C:258:ARG:HB3	1:C:261:GLN:OE1	2.05	0.55
1:B:6:PRO:HB2	1:B:196:VAL:HG12	1.88	0.55
1:C:19:ASN:O	1:C:277:LYS:HE2	2.07	0.54
1:C:201:ASP:HA	1:C:239:PHE:HB2	1.90	0.53
1:C:258:ARG:HB3	1:C:261:GLN:CD	2.34	0.52
1:C:44:THR:HG21	1:C:259:LEU:HG	1.92	0.52
1:C:26:PHE:HD2	1:C:239:PHE:CE1	2.28	0.51
1:D:45:TYR:CE1	1:D:262:LEU:HD21	2.46	0.50
1:A:133:ALA:HA	1:A:139:ILE:HD11	1.93	0.50
1:C:185:MET:HG3	1:D:185:MET:HE3	1.93	0.50
1:D:44:THR:HG21	1:D:259:LEU:HG	1.94	0.49
1:D:249:ALA:HB3	1:D:253:GLY:CA	2.42	0.49
1:A:203:THR:HG21	1:A:241:GLU:HB2	1.94	0.49
1:A:222:GLN:NE2	1:B:189:ASP:HB2	2.28	0.48
1:D:187:GLY:HA3	3:D:423:HOH:O	2.13	0.48
1:D:59:LYS:HD3	1:D:59:LYS:C	2.39	0.47
1:C:17:MET:HG2	1:C:24:VAL:HG21	1.97	0.47
1:D:17:MET:HG2	1:D:24:VAL:HG21	1.97	0.46
1:C:6:PRO:HB2	1:C:196:VAL:HG12	1.97	0.45
1:A:280:LYS:HA	1:A:280:LYS:HE3	1.97	0.45
1:C:253:GLY:N	1:C:254:PRO:HD2	2.31	0.45
1:A:199:PHE:CE2	1:A:237:GLY:HA3	2.52	0.45
1:C:59:LYS:HD3	1:C:59:LYS:C	2.42	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:19:ASN:O	1:B:277:LYS:HE2	2.17	0.45
1:B:188:PHE:O	1:B:192:LYS:HG3	2.17	0.44
1:A:120:ALA:HB1	1:A:144:ALA:HA	2.00	0.43
1:D:249:ALA:HB3	1:D:253:GLY:HA3	2.00	0.43
1:D:258:ARG:HB3	1:D:261:GLN:NE2	2.34	0.43
1:C:45:TYR:CD1	1:C:262:LEU:HD21	2.54	0.43
1:D:201:ASP:HA	1:D:239:PHE:HB3	2.01	0.42
1:C:185:MET:HE3	1:D:185:MET:HG3	2.01	0.42
1:A:253:GLY:N	1:A:254:PRO:CD	2.82	0.42
1:C:263:GLU:HB3	1:C:264:PRO:HD3	2.02	0.42
1:D:207:GLN:H	1:D:222:GLN:NE2	2.11	0.42
1:D:245:ASP:C	1:D:247:ASP:H	2.27	0.42
1:A:187:GLY:HA3	3:A:437:HOH:O	2.20	0.42
1:C:130:GLU:O	1:C:134:LYS:HG2	2.19	0.42
1:B:199:PHE:CE2	1:B:237:GLY:HA3	2.55	0.42
1:C:244:PRO:C	1:C:246:PRO:HD3	2.45	0.41
1:A:185:MET:HG3	1:B:185:MET:HE3	2.02	0.41
1:C:207:GLN:H	1:C:222:GLN:NE2	2.14	0.41
1:D:6:PRO:HB2	1:D:196:VAL:HG12	2.01	0.41
1:B:120:ALA:HB1	1:B:144:ALA:HA	2.03	0.41
1:C:30:ASN:OD1	1:C:64:LYS:HD2	2.21	0.41
1:C:117:GLN:OE1	1:C:239:PHE:CZ	2.74	0.41
1:B:59:LYS:C	1:B:59:LYS:HD3	2.46	0.41
1:C:226:LEU:HD23	1:C:226:LEU:HA	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/285 (99%)	277 (99%)	4 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	281/285 (99%)	277 (99%)	4 (1%)	0	100	100
1	C	270/285 (95%)	261 (97%)	9 (3%)	0	100	100
1	D	268/285 (94%)	260 (97%)	8 (3%)	0	100	100
All	All	1100/1140 (96%)	1075 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/238 (99%)	234 (99%)	2 (1%)	73	61
1	B	236/238 (99%)	233 (99%)	3 (1%)	61	43
1	C	231/238 (97%)	223 (96%)	8 (4%)	32	9
1	D	230/238 (97%)	228 (99%)	2 (1%)	70	57
All	All	933/952 (98%)	918 (98%)	15 (2%)	55	34

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	143	LYS
1	A	280	LYS
1	B	5	LYS
1	B	99	ASP
1	B	283	ASP
1	C	5	LYS
1	C	69	SER
1	C	130	GLU
1	C	208	THR
1	C	245	ASP
1	C	250	LYS
1	C	252	ASP
1	C	280	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	220	ARG
1	D	247	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	GLN
1	A	136	GLN
1	A	154	HIS
1	A	222	GLN
1	B	136	GLN
1	C	117	GLN
1	C	140	ASN
1	C	150	HIS
1	C	207	GLN
1	C	222	GLN
1	C	285	HIS
1	D	3	GLN
1	D	16	GLN
1	D	91	HIS
1	D	93	ASN
1	D	136	GLN
1	D	207	GLN
1	D	222	GLN
1	D	261	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	SO4	D	301	-	4,4,4	0.50	0	6,6,6	0.32	0
2	SO4	A	301	-	4,4,4	0.28	0	6,6,6	0.31	0
2	SO4	B	301	-	4,4,4	0.40	0	6,6,6	0.21	0
2	SO4	B	302	-	4,4,4	0.48	0	6,6,6	0.16	0
2	SO4	A	302	-	4,4,4	0.25	0	6,6,6	0.23	0
2	SO4	C	301	-	4,4,4	0.27	0	6,6,6	0.22	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand SO4 D 301




Bond lengths



Bond angles




Torsions




Rings


Ligand SO4 A 301




Bond lengths



Bond angles




Torsions




Rings


Ligand SO4 B 301




Bond lengths



Bond angles




Torsions




Rings


Ligand SO4 B 302




Bond lengths



Bond angles




Torsions




Rings


Ligand SO4 A 302




Bond lengths



Bond angles




Torsions



Rings

Ligand SO4 C 301



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	283/285 (99%)	-0.22	4 (1%) 73 79	10, 14, 28, 96	0
1	B	283/285 (99%)	-0.09	4 (1%) 73 79	10, 16, 30, 65	0
1	C	274/285 (96%)	0.27	31 (11%) 10 14	10, 16, 50, 81	0
1	D	272/285 (95%)	0.27	25 (9%) 14 20	10, 16, 46, 79	0
All	All	1112/1140 (97%)	0.05	64 (5%) 29 35	10, 16, 38, 96	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	249	ALA	6.5
1	D	253	GLY	6.1
1	C	218	GLY	5.5
1	D	254	PRO	5.3
1	C	208	THR	4.9
1	C	252	ASP	4.7
1	C	253	GLY	4.5
1	C	251	CYS	4.4
1	D	247	ASP	4.4
1	A	4	LEU	4.3
1	D	208	THR	4.3
1	C	254	PRO	4.3
1	B	4	LEU	4.2
1	C	250	LYS	4.1
1	C	285	HIS	4.0
1	D	248	LYS	4.0
1	C	239	PHE	4.0
1	D	252	ASP	3.9
1	C	219	ARG	3.8
1	C	249	ALA	3.8
1	D	246	PRO	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	68	SER	3.8
1	D	251	CYS	3.7
1	C	247	ASP	3.7
1	C	246	PRO	3.7
1	D	245	ASP	3.5
1	C	248	LYS	3.5
1	C	245	ASP	3.5
1	D	250	LYS	3.4
1	D	259	LEU	3.4
1	D	255	SER	3.3
1	C	68	SER	3.3
1	A	3	GLN	3.1
1	C	220	ARG	3.0
1	B	3	GLN	2.9
1	D	239	PHE	2.9
1	C	259	LEU	2.9
1	D	220	ARG	2.8
1	D	283	ASP	2.8
1	C	72	SER	2.7
1	C	280	LYS	2.7
1	D	221	ALA	2.7
1	C	283	ASP	2.7
1	D	72	SER	2.7
1	C	70	LEU	2.7
1	B	285	HIS	2.6
1	C	71	HIS	2.6
1	D	256	ALA	2.6
1	C	69	SER	2.6
1	C	284	THR	2.4
1	C	4	LEU	2.4
1	C	255	SER	2.4
1	A	285	HIS	2.4
1	C	281	LYS	2.3
1	D	285	HIS	2.3
1	B	5	LYS	2.3
1	C	256	ALA	2.3
1	D	70	LEU	2.2
1	D	257	LEU	2.2
1	C	14	ASP	2.1
1	D	3	GLN	2.1
1	A	5	LYS	2.1
1	D	244	PRO	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	221	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	SO4	A	302	5/5	0.99	0.04	13,14,16,16	0
2	SO4	B	301	5/5	0.99	0.04	13,14,15,17	0
2	SO4	B	302	5/5	0.99	0.04	15,15,16,19	0
2	SO4	C	301	5/5	0.99	0.05	15,16,17,18	0
2	SO4	D	301	5/5	0.99	0.04	16,16,18,18	0
2	SO4	A	301	5/5	1.00	0.03	13,13,14,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around SO4 A 302:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SO4 B 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SO4 B 302:


$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SO4 C 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around SO4 D 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)








Electron density around SO4 A 301:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.