



## wwPDB EM Validation Summary Report ⓘ

Jun 11, 2025 – 02:35 AM JST

PDB ID : 8XFP / pdb\_00008xfp  
EMDB ID : EMD-38307  
Title : the pentamerA complex of LGR4-RSPO2-ZNRF3(delta RING)  
Authors : Geng, Y.; Wang, L.  
Deposited on : 2023-12-14  
Resolution : 3.21 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev118  
MolProbity : 4-5-2 with Phenix2.0rc1  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.43.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	951	
2	E	560	
3	C	936	
3	H	936	
4	G	101	
4	J	101	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10742 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leucine-rich repeat-containing G-protein coupled receptor 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	734	Total	C	N	O	S	0	0
			5733	3725	943	1037	28		

- Molecule 2 is a protein called nanobody Nb52.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	E	114	Total	C	N	O	S	0	0
			876	550	149	170	7		

- Molecule 3 is a protein called E3 ubiquitin-protein ligase ZNRF3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	181	Total	C	N	O	S	0	0
			1400	896	232	266	6		
3	H	152	Total	C	N	O	S	0	0
			1171	736	202	229	4		

- Molecule 4 is a protein called R-spondin-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	G	100	Total	C	N	O	S	0	0
			781	479	144	140	18		
4	J	100	Total	C	N	O	S	0	0
			781	479	144	140	18		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	130	ALA	ASP	conflict	UNP Q8BFU0
J	130	ALA	ASP	conflict	UNP Q8BFU0



- Molecule 3: E3 ubiquitin-protein ligase ZNRF3


Tyr	Thr	Ser	Asp	T214		Met
His	Pro	Asn	Cys			Arg
Gly	Ala	Leu	Ala	L223	T75	Arg
His	Tyr	Ser	ILE		T76	Arg
Arg	ILE	Arg	Cys			Ser
Ser	Arg	Gly	Leu	V228	G83	Gly
Val	Ser	Arg	GLU	L231		Gly
Cys	Tyr	Arg	Lys		E95	Arg
Ser	Pro	Gln	Tyr			Pro
Pro	Pro	Arg	ILE	L240	I98	Gly
Leu	Leu	Val	Asp		V99	Ala
His	Thr	Leu	Gly	Lys	Q100	Thr
Leu	Leu	Leu	GLU	Lys		Arg
Ala	His	Val	GLU	Gln		Gly
Asp	Asp	Pro	GLU	Lys		Arg
Cys	His	Val	Leu	Arg	N109	Arg
Pro	Ser	His	Arg	Arg	N110	Arg
Pro	Pro	His	Ser	Ser	D111	Arg
Gly	Leu	Tyr	Val	Gln	E112	Arg
Ser	Ala	Pro	ILE	Asn	E113	Arg
Asp	Ala	Gly	Pro	Asn	D114	Arg
Ser	His	Arg	Cys	Met	L115	Leu
Ser	Arg	Val	Thr	Arg	Y116	Arg
Ser	Arg	His	His	Arg		Arg
Ser	Gly	Arg	Arg	Leu	G119	Pro
Ser	Leu	Thr	Phe	Ala		Arg
Ser	Leu	Asn	His	Val	K125	Gly
Gly	His	Ala	Arg	Gln		Leu
Gln	Arg	ILE	Lys	Ala	L131	Arg
Cys	Ala	Pro	Cys	Leu		Cys
Pro	Tyr	Ala	Val	GLU	G141	Ser
Pro	Ser	Tyr	Asp	Lys		Ser
Cys	Pro	Pro	Asp	Met	K144	Arg
Ser	Ala	Thr	Trp	GLU		Pro
Ser	His	Arg	Leu	Thr	V147	Pro
Ser	Pro	Thr	Leu	Arg	Q148	Pro
Asp	Phe	Ser	Gln	Lys	R149	Pro
Ser	Arg	Met	His	Phe		Pro
Leu	Arg	Ser	His	Asn	D157	Leu
Val	Pro	Ser	Thr	Ser		Pro
Asp	Lys	His	Cys	Lys	L176	Leu
Thr	Ser	Gly	Pro	Ser		Leu
Thr	Leu	Asn	His	Lys	V183	Gly
Val	Gly	Pro	Cys	Gly		Leu
Val	Arg	Val	Arg	Arg	M192	Leu
Ser	Ser	Thr	His	Arg		Leu
Asn	Phe	Leu	His	GLU	V195	Leu
Pro	Ser	Leu	ILE	Gly	N196	Ala
Gln	Ser	Leu	ILE	Ser	K197	Ala
Gly	Lys	Thr	Ser	Cys	Q198	Gly
Val	Ala	Met	GLN	GLY		Ala
Tyr	Ala	Asp	GLN	Cys		Ala
His	Cys	Arg	Lys	Leu	R202	Arg
Gly	Phe	His	Gly	Leu	A203	Gly
Ser	Ser	Gly	Asn	Asp	R204	Ala
Cys	Ser	GLY	Pro	Thr		Ala
Thr	Tyr	Gln	Ser	Leu	H207	Arg
Phe	GLU	Ser	Ala	Ser		Ala
Arg	Thr	Leu	Val	Ser	Arg	Arg
Ser	Met	Tyr	Cys	Ser	Pro	Arg
Ser	Tyr	Ser	Val	Ser		Arg
Phe	Phe	Ser	Val	Ser		Arg
Leu	Gln	Pro	GLU	Thr	Q212	Val
Cys	His	Gln	Thr	Thr		Val




ARG  
ASP  
LEU  
SER  
ALA  
GLN  
ARG  
TLE  
PRO  
TLE  
TLE  
CYS  
PRO  
PRO  
GLU  
ASP  
VAL  
ASP  
CYS  
ASP  
LEU  
GLY  
LEU  
PRO  
ASN  
SER  
ASP  
CYS  
GLN  
GLY  
THR  
HIS  
SER  
LEU  
GLY  
SER  
TRP  
GLY  
GLY  
THR  
ARG  
GLY  
PRO  
ASP  
THR  
ARG  
PRO  
HIS  
SER  
ALA  
GLY  
LEU  
GLY  
ALA  
THR  
SER  
THR  
ARG  
GLY  
GLU  
ARG  
ALA  
CYS

GLN  
ALA  
ARG  
ALA  
LEU  
ARG  
LEU  
ARG  
PRO  
GLY  
CYS  
PRO  
PRO  
GLU  
GLU  
GLU  
ALA  
GLY  
ALA  
VAL  
ARG  
ALA  
ASN  
PHE  
PRO  
ASP  
SER  
ALA  
LEU  
GLN  
ASP  
THR  
GLN  
GLY  
SER  
SER  
THR  
THR  
ALA  
THR  
THR  
GLU  
ALA  
ALA  
GLY  
PRO  
ARG  
SER  
HIS  
SER  
ALA  
ASP  
SER  
SER  
SER  
PRO  
GLY  
ALA

● Molecule 4: R-spondin-2



● Molecule 4: R-spondin-2



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	73170	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TECNAI F30	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	1.944	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.393	Depositor
Minimum map value	-0.228	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.008	Depositor
Recommended contour level	0.04	Depositor
Map size ( $\text{\AA}$ )	385.56, 385.56, 385.56	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.071, 1.071, 1.071	Depositor



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.10	0/5869	0.27	0/7981
2	E	0.10	0/894	0.26	0/1212
3	C	0.10	0/1425	0.26	0/1934
3	H	0.10	0/1191	0.24	0/1615
4	G	0.08	0/798	0.25	0/1062
4	J	0.10	0/798	0.27	0/1062
All	All	0.10	0/10975	0.27	0/14866

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5733	0	5787	79	0
2	E	876	0	841	18	0
3	C	1400	0	1405	24	0
3	H	1171	0	1167	18	0
4	G	781	0	711	14	0
4	J	781	0	710	17	0
All	All	10742	0	10621	158	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 158 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:518:TYR:O	2:E:543:GLY:HA2	1.87	0.74
2:E:515:THR:HA	2:E:546:VAL:O	1.87	0.73
1:A:329:THR:OG1	1:A:353:ASN:OD1	2.10	0.69
1:A:161:ASP:OD1	1:A:162:ASP:N	2.25	0.69
4:G:43:CYS:SG	4:G:44:LEU:N	2.65	0.68

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	726/951 (76%)	679 (94%)	47 (6%)	0	100	100
2	E	110/560 (20%)	105 (96%)	5 (4%)	0	100	100
3	C	177/936 (19%)	170 (96%)	6 (3%)	1 (1%)	22	56
3	H	150/936 (16%)	146 (97%)	4 (3%)	0	100	100
4	G	96/101 (95%)	89 (93%)	7 (7%)	0	100	100
4	J	96/101 (95%)	87 (91%)	9 (9%)	0	100	100
All	All	1355/3585 (38%)	1276 (94%)	78 (6%)	1 (0%)	50	79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	75	THR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	646/827 (78%)	633 (98%)	13 (2%)	50	73
2	E	93/452 (21%)	93 (100%)	0	100	100
3	C	153/773 (20%)	150 (98%)	3 (2%)	50	73
3	H	127/773 (16%)	120 (94%)	7 (6%)	18	49
4	G	85/88 (97%)	84 (99%)	1 (1%)	67	83
4	J	85/88 (97%)	84 (99%)	1 (1%)	67	83
All	All	1189/3001 (40%)	1164 (98%)	25 (2%)	49	72

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	98	ILE
3	H	65	LEU
4	J	136	ASP
4	G	119	LEU
3	H	85	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
3	H	100	GLN
3	H	207	HIS
4	J	91	ASN
3	H	196	ASN
1	A	354	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-38307. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 180



Y Index: 180



Z Index: 180


### 6.2.2 Raw map



X Index: 180



Y Index: 180



Z Index: 180

The images above show central slices of the map in three orthogonal directions.




## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 192




Y Index: 205




Z Index: 144


### 6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 359


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.04. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 295 nm<sup>3</sup>; this corresponds to an approximate mass of 267 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.312 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of  $0.312 \text{ \AA}^{-1}$

## 8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.21	-	-
Author-provided FSC curve	3.21	3.61	3.32
Unmasked-calculated*	4.36	9.29	6.39

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.36 differs from the reported value 3.21 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-38307 and PDB model 8XFP. Per-residue inclusion information can be found in [section 3](#) on [page 4](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.04 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.04).



## 9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 92% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.04) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div>0.9200</div>	<div><div></div>0.4050</div>
A	<div><div></div>0.9170</div>	<div><div></div>0.3980</div>
C	<div><div></div>0.9150</div>	<div><div></div>0.4040</div>
E	<div><div></div>0.9450</div>	<div><div></div>0.4180</div>
G	<div><div></div>0.8750</div>	<div><div></div>0.2880</div>
H	<div><div></div>0.9210</div>	<div><div></div>0.4650</div>
J	<div><div></div>0.9620</div>	<div><div></div>0.4740</div>

1.0

0.0

<0.0