wwPDB X-ray Structure Validation Summary Report (i)

Feb 5, 2024 – 12:21 PM EST

PDB ID	:	1XX7
Title	:	Conserved hypothetical protein from Pyrococcus furiosus Pfu-403030-001
Authors	:	Chen, L.; Tempel, W.; Habel, J.; Zhou, W.; Nguyen, D.; Chang, S.-H.; Lee, D.; Kelley, L.-L.C.; Dillard, B.D.; Liu, Z.-J.; Bridger, S.; Eneh, J.C.; Hopkins, R.C.; Jenney Jr, F.E.; Lee, H.-S.; Li, T.; Poole II, F.L.; Shah, C.; Sugar, F.J.; Adams, M.W.W.; Arendall III, W.B.; Richardson, J.S.; Richardson, D.C.; Rose, J.P.; Wang, B.-C.; Southeast Collaboratory for Structural Genomics (SECSG)
Deposited on	:	2004-11-04
Resolution	:	2.26 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	184	% 86% 7% 7%
1	B	184	3% 86% 7% • 7%
1	C	184	3% 87% 6% • 7%
1	D	184	83% 5% 12%

Continued on next page...

Validation Pipeline (wwPDB-VP) : 2.36

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	184	% 82% 8% 10%
1	F	184	% 85% 5% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	UNX	A	2001	-	-	-	X
3	UNX	A	2002	-	-	-	X
3	UNX	A	2003	-	-	-	X
3	UNX	A	2006	-	-	-	X
3	UNX	A	2019	-	-	-	X
3	UNX	B	2004	-	-	-	X
3	UNX	B	2005	-	-	-	X
3	UNX	B	2007	-	-	-	X
3	UNX	B	2009	-	-	-	X
3	UNX	C	2008	-	-	-	X
3	UNX	D	2011	-	-	-	X
3	UNX	D	2012	-	-	-	X
3	UNX	E	2013	-	-	-	X
3	UNX	E	2014	-	-	-	X
3	UNX	E	2018	-	-	-	X
3	UNX	F	2016	-	-	-	X
3	UNX	F	2017	-	-	-	X

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 7822 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called oxetanocin-like protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	172	1329	867	210	250	2	0	0	0
1	B	172	1304	847	211	244	2	0	0	0
1	C	172	1304	851	204	247	2	0	0	0
1	D	161	1231	803	193	233	2	0	0	0
1	E	166	1283	839	202	240	2	0	0	0
1	F	165	1277	835	200	240	2	0	0	0

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	ALA	-	expression tag	UNP Q8U3R1
A	-6	HIS	-	expression tag	UNP Q8U3R1
A	-5	HIS	-	expression tag	UNP Q8U3R1
A	-4	HIS	-	expression tag	UNP Q8U3R1
A	-3	HIS	-	expression tag	UNP Q8U3R1
A	-2	HIS	-	expression tag	UNP Q8U3R1
A	-1	HIS	-	expression tag	UNP Q8U3R1
A	0	GLY	-	expression tag	UNP Q8U3R1
A	1	SER	-	expression tag	UNP Q8U3R1
B	-7	ALA	-	expression tag	UNP Q8U3R1
B	-6	HIS	-	expression tag	UNP Q8U3R1
B	-5	HIS	-	expression tag	UNP Q8U3R1
B	-4	HIS	-	expression tag	UNP Q8U3R1
B	-3	HIS	-	expression tag	UNP Q8U3R1
B	-2	HIS	-	expression tag	UNP Q8U3R1
B	-1	HIS	-	expression tag	UNP Q8U3R1
B	0	GLY	-	expression tag	UNP Q8U3R1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	SER	-	expression tag	UNP Q8U3R1
C	-7	ALA	-	expression tag	UNP Q8U3R1
C	-6	HIS	-	expression tag	UNP Q8U3R1
C	-5	HIS	-	expression tag	UNP Q8U3R1
C	-4	HIS	-	expression tag	UNP Q8U3R1
C	-3	HIS	-	expression tag	UNP Q8U3R1
C	-2	HIS	-	expression tag	UNP Q8U3R1
C	-1	HIS	-	expression tag	UNP Q8U3R1
C	0	GLY	-	expression tag	UNP Q8U3R1
C	1	SER	-	expression tag	UNP Q8U3R1
D	-7	ALA	-	expression tag	UNP Q8U3R1
D	-6	HIS	-	expression tag	UNP Q8U3R1
D	-5	HIS	-	expression tag	UNP Q8U3R1
D	-4	HIS	-	expression tag	UNP Q8U3R1
D	-3	HIS	-	expression tag	UNP Q8U3R1
D	-2	HIS	-	expression tag	UNP Q8U3R1
D	-1	HIS	-	expression tag	UNP Q8U3R1
D	0	GLY	-	expression tag	UNP Q8U3R1
D	1	SER	-	expression tag	UNP Q8U3R1
E	-7	ALA	-	expression tag	UNP Q8U3R1
E	-6	HIS	-	expression tag	UNP Q8U3R1
E	-5	HIS	-	expression tag	UNP Q8U3R1
E	-4	HIS	-	expression tag	UNP Q8U3R1
E	-3	HIS	-	expression tag	UNP Q8U3R1
E	-2	HIS	-	expression tag	UNP Q8U3R1
E	-1	HIS	-	expression tag	UNP Q8U3R1
E	0	GLY	-	expression tag	UNP Q8U3R1
E	1	SER	-	expression tag	UNP Q8U3R1
F	-7	ALA	-	expression tag	UNP Q8U3R1
F	-6	HIS	-	expression tag	UNP Q8U3R1
F	-5	HIS	-	expression tag	UNP Q8U3R1
F	-4	HIS	-	expression tag	UNP Q8U3R1
F	-3	HIS	-	expression tag	UNP Q8U3R1
F	-2	HIS	-	expression tag	UNP Q8U3R1
F	-1	HIS	-	expression tag	UNP Q8U3R1
F	0	GLY	-	expression tag	UNP Q8U3R1
F	1	SER	-	expression tag	UNP Q8U3R1

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ni 1 1	0	0
2	B	1	Total Ni 1 1	0	0
2	C	1	Total Ni 1 1	0	0
2	D	1	Total Ni 1 1	0	0
2	E	1	Total Ni 1 1	0	0
2	F	1	Total Ni 1 1	0	0

- Molecule 3 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	5	Total X 5 5	0	0
3	B	4	Total X 4 4	0	0
3	C	1	Total X 1 1	0	0
3	D	4	Total X 4 4	0	0
3	E	3	Total X 3 3	0	0
3	F	2	Total X 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	13	Total O 13 13	0	0
4	B	11	Total O 11 11	0	0
4	C	10	Total O 10 10	0	0
4	D	13	Total O 13 13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	E	11	11	11	0	0
			Total	O		
4	F	11	11	11	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: oxetanocin-like protein

Chain A:



- Molecule 1: oxetanocin-like protein

Chain B:



- Molecule 1: oxetanocin-like protein

Chain C:



- Molecule 1: oxetanocin-like protein

Chain D:



- Molecule 1: oxetanocin-like protein

Chain E:



- Molecule 1: oxetanocin-like protein

Chain F:


%
85%
5%
10%
ALA
HIS
HIS
HIS
HIS
HIS
HIS
GLY
S1
R13
K59
L69
E108
K111
A112
L113
Y133
W147
N148
A149
L150
GLU
ASP
LEU
GLU
LYS
LEU
GLU
I158
S159
L172
LYS
ASP
ASP
HIS

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	98.07Å 111.33Å 125.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.33 – 2.26 29.99 – 2.26	Depositor EDS
% Data completeness (in resolution range)	99.4 (83.33-2.26) 99.4 (29.99-2.26)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.55 (at 2.26Å)	Xtriage
Refinement program	REFMAC refmac_5.2.0005	Depositor
R, R_{free}	0.200 , 0.246 0.208 , 0.253	Depositor DCC
R_{free} test set	2015 reflections (3.11%)	wwPDB-VP
Wilson B-factor (Å^2)	27.3	Xtriage
Anisotropy	0.132	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 49.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	7822	wwPDB-VP
Average B, all atoms (Å^2)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.88	2/1349 (0.1%)	0.73	0/1831
1	B	0.81	0/1323	0.70	1/1793 (0.1%)
1	C	0.85	0/1324	0.76	4/1800 (0.2%)
1	D	0.86	0/1248	0.74	0/1691
1	E	0.85	0/1300	0.72	2/1759 (0.1%)
1	F	0.84	0/1296	0.74	0/1757
All	All	0.85	2/7840 (0.0%)	0.73	7/10631 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	46	GLU	CD-OE2	6.27	1.32	1.25
1	A	46	GLU	CG-CD	5.88	1.60	1.51

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	160	ARG	NE-CZ-NH2	-6.19	117.21	120.30
1	E	171	ARG	NE-CZ-NH2	-6.01	117.29	120.30
1	B	13	ARG	NE-CZ-NH1	5.72	123.16	120.30
1	C	32	ASP	CB-CG-OD1	5.71	123.44	118.30
1	C	160	ARG	NE-CZ-NH1	5.62	123.11	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1329	0	1325	8	0
1	B	1304	0	1291	8	0
1	C	1304	0	1281	8	0
1	D	1231	0	1222	11	0
1	E	1283	0	1301	9	0
1	F	1277	0	1279	5	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	5	0	0	0	0
3	B	4	0	0	0	0
3	C	1	0	0	0	0
3	D	4	0	0	0	0
3	E	3	0	0	0	0
3	F	2	0	0	0	0
4	A	13	0	0	0	0
4	B	11	0	0	0	0
4	C	10	0	0	0	0
4	D	13	0	0	0	0
4	E	11	0	0	0	0
4	F	11	0	0	0	0
All	All	7822	0	7699	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 41 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:122:ILE:CD1	1:D:158:ILE:CG2	2.69	0.70
1:D:122:ILE:CD1	1:D:158:ILE:HG23	2.27	0.65
1:C:59:LYS:HE2	1:C:108:GLU:OE2	2.02	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:122:ILE:HD11	1:D:158:ILE:CG2	2.32	0.59
1:A:20:LEU:HD21	1:D:13:ARG:O	2.04	0.57

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	170/184 (92%)	169 (99%)	1 (1%)	0	100	100
1	B	170/184 (92%)	164 (96%)	5 (3%)	1 (1%)	25	25
1	C	170/184 (92%)	165 (97%)	3 (2%)	2 (1%)	13	9
1	D	157/184 (85%)	154 (98%)	3 (2%)	0	100	100
1	E	162/184 (88%)	159 (98%)	3 (2%)	0	100	100
1	F	161/184 (88%)	158 (98%)	3 (2%)	0	100	100
All	All	990/1104 (90%)	969 (98%)	18 (2%)	3 (0%)	41	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	101	GLU
1	B	51	LYS
1	C	171	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	129/159 (81%)	129 (100%)	0	100	100
1	B	124/159 (78%)	122 (98%)	2 (2%)	62	73
1	C	124/159 (78%)	123 (99%)	1 (1%)	81	88
1	D	119/159 (75%)	119 (100%)	0	100	100
1	E	127/159 (80%)	126 (99%)	1 (1%)	81	88
1	F	125/159 (79%)	124 (99%)	1 (1%)	81	88
All	All	748/954 (78%)	743 (99%)	5 (1%)	84	90

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1	SER
1	B	159	SER
1	C	101	GLU
1	E	115	LEU
1	F	159	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 25 ligands modelled in this entry, 6 are monoatomic and 19 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	172/184 (93%)	-0.28	1 (0%)	89	89	13, 28, 51, 59	0
1	B	172/184 (93%)	-0.10	5 (2%)	51	55	15, 28, 68, 93	0
1	C	172/184 (93%)	-0.00	6 (3%)	44	46	15, 33, 62, 85	0
1	D	161/184 (87%)	-0.33	0	100	100	15, 28, 48, 65	0
1	E	166/184 (90%)	-0.30	1 (0%)	89	89	15, 30, 51, 61	0
1	F	165/184 (89%)	-0.22	1 (0%)	89	89	14, 27, 53, 66	0
All	All	1008/1104 (91%)	-0.21	14 (1%)	75	77	13, 29, 57, 93	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	169	VAL	4.4
1	C	152	ASP	4.2
1	C	113	LEU	3.9
1	C	151	GLU	3.4
1	F	148	ASN	3.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	UNX	F	2016	1/1	0.24	0.81	2,2,2,2	1
3	UNX	A	2002	1/1	0.34	2.74	2,2,2,2	1
3	UNX	B	2007	1/1	0.38	1.30	2,2,2,2	1
3	UNX	B	2009	1/1	0.52	1.64	2,2,2,2	1
3	UNX	B	2005	1/1	0.55	1.03	2,2,2,2	1
3	UNX	F	2017	1/1	0.56	2.13	2,2,2,2	1
3	UNX	E	2013	1/1	0.57	1.09	2,2,2,2	1
3	UNX	D	2011	1/1	0.63	1.57	2,2,2,2	1
3	UNX	E	2014	1/1	0.65	1.57	2,2,2,2	1
3	UNX	A	2006	1/1	0.70	2.01	2,2,2,2	1
3	UNX	A	2003	1/1	0.71	1.38	2,2,2,2	1
3	UNX	E	2018	1/1	0.71	1.53	2,2,2,2	1
3	UNX	A	2001	1/1	0.72	1.85	2,2,2,2	1
3	UNX	B	2004	1/1	0.73	1.42	2,2,2,2	1
3	UNX	C	2008	1/1	0.74	1.16	2,2,2,2	1
3	UNX	D	2012	1/1	0.76	2.39	2,2,2,2	1
3	UNX	A	2019	1/1	0.78	0.57	2,2,2,2	1
3	UNX	D	2010	1/1	0.80	0.89	2,2,2,2	1
3	UNX	D	2015	1/1	0.88	1.18	2,2,2,2	1
2	NI	F	201	1/1	0.96	0.09	33,33,33,33	0
2	NI	C	201	1/1	0.97	0.11	35,35,35,35	0
2	NI	D	201	1/1	0.97	0.10	30,30,30,30	0
2	NI	E	201	1/1	0.98	0.09	34,34,34,34	0
2	NI	A	201	1/1	0.98	0.08	32,32,32,32	0
2	NI	B	201	1/1	0.98	0.12	34,34,34,34	0

6.5 Other polymers (i)

There are no such residues in this entry.