Full wwPDB X-ray Structure Validation Report (i)

Jun 24, 2024 – 08:09 AM EDT

PDB ID	:	6YBG
Title	:	Structure of Mcl-1 in complex with compound 2g
Authors	:	Dokurno, P.; Surgenor, A.E.; Murray, J.B.
Deposited on	:	2020-03-17
Resolution	:	2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	10% 71% 14% 16%
1	B	170	8% 66% 16% • 15%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2450 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Induced myeloid leukemia cell differentiation protein Mcl-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	143	1155	726	212	214	3	0	0	0
			Total	C	N	O	S			
1	B	144	1133	713	203	214	3	0	0	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	158	MET	-	initiating methionine	UNP Q07820
A	159	HIS	-	expression tag	UNP Q07820
A	160	HIS	-	expression tag	UNP Q07820
A	161	HIS	-	expression tag	UNP Q07820
A	162	HIS	-	expression tag	UNP Q07820
A	163	HIS	-	expression tag	UNP Q07820
A	164	HIS	-	expression tag	UNP Q07820
A	165	LEU	-	expression tag	UNP Q07820
A	166	VAL	-	expression tag	UNP Q07820
A	167	PRO	-	expression tag	UNP Q07820
A	168	ARG	-	expression tag	UNP Q07820
A	169	GLY	-	expression tag	UNP Q07820
A	170	SER	-	expression tag	UNP Q07820
B	158	MET	-	initiating methionine	UNP Q07820
B	159	HIS	-	expression tag	UNP Q07820
B	160	HIS	-	expression tag	UNP Q07820
B	161	HIS	-	expression tag	UNP Q07820
B	162	HIS	-	expression tag	UNP Q07820
B	163	HIS	-	expression tag	UNP Q07820
B	164	HIS	-	expression tag	UNP Q07820
B	165	LEU	-	expression tag	UNP Q07820
B	166	VAL	-	expression tag	UNP Q07820
B	167	PRO	-	expression tag	UNP Q07820
B	168	ARG	-	expression tag	UNP Q07820
B	169	GLY	-	expression tag	UNP Q07820

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	170	SER	-	expression tag	UNP Q07820

- Molecule 2 is (2 {R})-2-[5-[3-chloranyl-2-methyl-4-[2-(4-methylpiperazin-1-yl)ethoxy]phenyl]-6-(3-chlorophenyl)thieno[2,3-d]pyrimidin-4-yl]oxy-3-(2-methoxyphenyl)propanoic acid (three-letter code: OJT) (formula: $C_{36}H_{36}Cl_2N_4O_5S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	Cl	N	O	S		
2	A	1	48	36	2	4	5	1	0	0
2	B	1	48	36	2	4	5	1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
3	A	1	1	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	27	27	27	0	0
4	B	38	38	38	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Induced myeloid leukemia cell differentiation protein Mcl-1

Chain A:



• Molecule 1: Induced myeloid leukemia cell differentiation protein Mcl-1

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	126.75Å 79.06Å 42.29Å 90.00° 98.86° 90.00°	Depositor
Resolution (Å)	20.00 – 2.10 19.94 – 2.10	Depositor EDS
% Data completeness (in resolution range)	78.9 (20.00-2.10) 79.0 (19.94-2.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.28 (at 2.09Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.180 , 0.235 0.187 , 0.233	Depositor DCC
R_{free} test set	885 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å^2)	40.9	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 49.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	2450	wwPDB-VP
Average B, all atoms (Å^2)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.01% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: OJT, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.71	0/1173	0.80	0/1576
1	B	0.75	0/1151	0.85	1/1555 (0.1%)
All	All	0.73	0/2324	0.82	1/3131 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	233	ARG	NE-CZ-NH2	-5.51	117.55	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1155	0	1158	7	0
1	B	1133	0	1098	20	0
2	A	48	0	0	0	0
2	B	48	0	0	0	0
3	A	1	0	0	0	0
4	A	27	0	0	0	0
4	B	38	0	0	6	0
All	All	2450	0	2256	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:172:ASP:OD2	1:B:303:ARG:NH2	2.10	0.83
1:B:179:LEU:O	1:B:183:SER:HB2	1.86	0.76
1:B:248:ARG:HG3	4:B:530:HOH:O	1.97	0.64
1:B:235:LEU:HD13	1:B:246:LEU:HD11	1.82	0.62
1:B:235:LEU:HD13	1:B:246:LEU:CD1	2.34	0.58
1:B:322:GLU:OE1	4:B:501:HOH:O	2.18	0.55
1:B:216:VAL:HG12	1:B:265:VAL:HG11	1.88	0.55
1:B:251:ILE:HG12	1:B:297:VAL:CG2	2.36	0.54
1:B:236:ASP:O	4:B:502:HOH:O	2.19	0.52
1:A:258:VAL:O	1:A:263:ARG:HD3	2.11	0.50
1:B:240:GLU:HG3	1:B:286:CYS:SG	2.55	0.46
1:A:259:THR:OG1	1:A:302:LYS:HD2	2.16	0.46
1:B:187:ARG:O	1:B:191:THR:HG23	2.16	0.45
1:A:251:ILE:HG13	1:A:297:VAL:HG22	1.97	0.45
1:B:305:TRP:CH2	1:B:309:GLN:HG3	2.52	0.45
1:A:204:ALA:O	1:A:208:LYS:HG2	2.17	0.45
1:B:312:TRP:O	1:B:316:VAL:HG23	2.16	0.44
1:B:302:LYS:NZ	4:B:510:HOH:O	2.50	0.44
1:B:293:SER:O	1:B:297:VAL:HG23	2.18	0.43
1:A:232:LEU:HD21	1:A:277:HIS:CG	2.54	0.42
1:A:212:THR:O	1:A:216:VAL:HG23	2.20	0.42
1:B:247:SER:O	1:B:250:MET:HB2	2.20	0.42
1:B:322:GLU:HB2	4:B:505:HOH:O	2.19	0.41
1:B:243:VAL:O	1:B:246:LEU:HB3	2.20	0.41
1:B:277:HIS:ND1	4:B:504:HOH:O	2.37	0.41
1:B:305:TRP:CZ3	1:B:309:GLN:HG3	2.56	0.41
1:A:261:TRP:CH2	1:A:315:PHE:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/170 (82%)	135 (97%)	4 (3%)	0	100	100
1	B	140/170 (82%)	132 (94%)	8 (6%)	0	100	100
All	All	279/340 (82%)	267 (96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	125/148 (84%)	116 (93%)	9 (7%)	14	11
1	B	119/148 (80%)	112 (94%)	7 (6%)	19	17
All	All	244/296 (82%)	228 (93%)	16 (7%)	16	14

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	173	GLU
1	A	194	LYS
1	A	234	LYS
1	A	248	ARG
1	A	300	ARG
1	A	303	ARG
1	A	309	GLN
1	A	317	GLU
1	A	320	HIS
1	B	183	SER
1	B	247	SER
1	B	248	ARG
1	B	296	ASP
1	B	303	ARG
1	B	304	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	310	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	177	GLN
1	B	282	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	OJT	B	401	-	49,53,53	0.85	3 (6%)	58,75,75	0.98	5 (8%)
2	OJT	A	401	-	49,53,53	0.80	3 (6%)	58,75,75	1.00	4 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OJT	B	401	-	-	2/28/38/38	0/6/6/6
2	OJT	A	401	-	-	3/28/38/38	0/6/6/6

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	OJT	C12-C13	-2.89	1.37	1.42
2	B	401	OJT	C12-C13	-2.58	1.38	1.42
2	B	401	OJT	C22-C14	-2.51	1.47	1.50
2	A	401	OJT	C22-C14	-2.25	1.47	1.50
2	B	401	OJT	O4-C36	-2.14	1.23	1.30
2	A	401	OJT	O4-C36	-2.04	1.24	1.30

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	401	OJT	C14-C13-C12	3.52	110.34	107.54
2	B	401	OJT	C3-C2-C1	-3.23	110.35	114.13
2	A	401	OJT	C3-C2-C1	-3.21	110.36	114.13
2	A	401	OJT	N2-C11-N1	-2.95	124.67	128.67
2	A	401	OJT	C15-C14-C13	2.71	113.34	109.16
2	B	401	OJT	C15-C14-C13	2.57	113.13	109.16
2	B	401	OJT	N2-C11-N1	-2.35	125.48	128.67
2	B	401	OJT	C14-C13-C12	2.03	109.16	107.54
2	B	401	OJT	C13-C12-N2	-2.02	121.60	124.89

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	OJT	C13-C14-C22-C35
2	B	401	OJT	C13-C14-C22-C35
2	A	401	OJT	C15-C14-C22-C35
2	B	401	OJT	C15-C14-C22-C35
2	A	401	OJT	C27-C28-N3-C29

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand OJT B 401



Ligand OJT A 401



Bond lengths



Bond angles



Torsions



Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	143/170 (84%)	0.52	17 (11%)	4	5	30, 46, 79, 104	7 (4%)
1	B	144/170 (84%)	0.48	14 (9%)	7	10	29, 49, 79, 99	6 (4%)
All	All	287/340 (84%)	0.50	31 (10%)	5	7	29, 48, 80, 104	13 (4%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	193	ALA	6.2
1	A	195	ASP	5.2
1	A	193	ALA	5.1
1	B	300	ARG	4.4
1	A	268	ILE	3.9
1	A	202	SER	3.8
1	A	194	LYS	3.8
1	A	204	ALA	3.5
1	B	195	ASP	3.5
1	B	194	LYS	3.4
1	B	312	TRP	3.4
1	A	203	GLY	3.3
1	B	172	ASP	3.3
1	B	267	LEU	3.3
1	B	304	ASP	3.2
1	B	204	ALA	3.2
1	B	268	ILE	3.2
1	A	320	HIS	2.9
1	A	267	LEU	2.8
1	B	310	ARG	2.7
1	A	272	ALA	2.6
1	B	250	MET	2.4
1	A	300	ARG	2.4
1	A	294	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	182	ILE	2.3
1	A	271	GLY	2.1
1	B	246	LEU	2.1
1	A	274	VAL	2.1
1	B	264	ILE	2.0
1	A	215	ARG	2.0
1	A	256	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	CL	A	402	1/1	0.91	0.24	86,86,86,86	0
2	OJT	B	401	48/48	0.95	0.11	29,39,73,81	0
2	OJT	A	401	48/48	0.95	0.12	31,40,89,101	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OJT B 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OJT A 401:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.