



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 23, 2025 – 12:10 PM EDT

PDB ID : 9YUP / pdb\_00009yup  
Title : Crystal structure of PprA S-F-S tetramer from *Deinococcus radiodurans*  
Authors : Szabla, R.; Junop, M.S.  
Deposited on : 2025-10-22  
Resolution : 2.07 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.46


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3436 (2.08-2.04)
Clashscore	180529	3661 (2.08-2.04)
Ramachandran outliers	177936	3649 (2.08-2.04)
Sidechain outliers	177891	3649 (2.08-2.04)
RSRZ outliers	164620	3436 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	<div> <div>5%</div> <div>91%</div> <div>5%</div> </div>
1	B	307	<div> <div>14%</div> <div>85%</div> <div>10%</div> </div>
1	C	307	<div> <div>8%</div> <div>83%</div> <div>6%</div> <div>11%</div> </div>
1	D	307	<div> <div>51%</div> <div>76%</div> <div>11%</div> <div>12%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16960 atoms, of which 8305 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA repair protein PprA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	296	Total	C	H	N	O	Se	2220	0	0
			4472	1403	2216	414	435	4			
1	B	275	Total	C	H	N	O	Se	2125	0	0
			4120	1292	2040	384	400	4			
1	C	273	Total	C	H	N	O	Se	2036	0	0
			4104	1286	2034	383	397	4			
1	D	269	Total	C	H	N	O	Se	2154	0	0
			4039	1268	2000	376	391	4			

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-31	MSE	-	initiating methionine	UNP O32504
A	-30	ARG	-	expression tag	UNP O32504
A	-29	SER	-	expression tag	UNP O32504
A	-28	GLY	-	expression tag	UNP O32504
A	-27	SER	-	expression tag	UNP O32504
A	-26	HIS	-	expression tag	UNP O32504
A	-25	HIS	-	expression tag	UNP O32504
A	-24	HIS	-	expression tag	UNP O32504
A	-23	HIS	-	expression tag	UNP O32504
A	-22	HIS	-	expression tag	UNP O32504
A	-21	HIS	-	expression tag	UNP O32504
A	-20	ARG	-	expression tag	UNP O32504
A	-19	SER	-	expression tag	UNP O32504
A	-18	ASP	-	expression tag	UNP O32504
A	-17	ILE	-	expression tag	UNP O32504
A	-16	THR	-	expression tag	UNP O32504
A	-15	SER	-	expression tag	UNP O32504
A	-14	LEU	-	expression tag	UNP O32504
A	-13	TYR	-	expression tag	UNP O32504
A	-12	LYS	-	expression tag	UNP O32504
A	-11	LYS	-	expression tag	UNP O32504

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	ALA	-	expression tag	UNP O32504
A	-9	GLY	-	expression tag	UNP O32504
A	-8	LEU	-	expression tag	UNP O32504
A	-7	GLU	-	expression tag	UNP O32504
A	-6	ASN	-	expression tag	UNP O32504
A	-5	LEU	-	expression tag	UNP O32504
A	-4	TYR	-	expression tag	UNP O32504
A	-3	PHE	-	expression tag	UNP O32504
A	-2	GLN	-	expression tag	UNP O32504
A	-1	GLY	-	expression tag	UNP O32504
A	180	LYS	ASP	engineered mutation	UNP O32504
A	184	LYS	ASP	engineered mutation	UNP O32504
B	-31	MSE	-	initiating methionine	UNP O32504
B	-30	ARG	-	expression tag	UNP O32504
B	-29	SER	-	expression tag	UNP O32504
B	-28	GLY	-	expression tag	UNP O32504
B	-27	SER	-	expression tag	UNP O32504
B	-26	HIS	-	expression tag	UNP O32504
B	-25	HIS	-	expression tag	UNP O32504
B	-24	HIS	-	expression tag	UNP O32504
B	-23	HIS	-	expression tag	UNP O32504
B	-22	HIS	-	expression tag	UNP O32504
B	-21	HIS	-	expression tag	UNP O32504
B	-20	ARG	-	expression tag	UNP O32504
B	-19	SER	-	expression tag	UNP O32504
B	-18	ASP	-	expression tag	UNP O32504
B	-17	ILE	-	expression tag	UNP O32504
B	-16	THR	-	expression tag	UNP O32504
B	-15	SER	-	expression tag	UNP O32504
B	-14	LEU	-	expression tag	UNP O32504
B	-13	TYR	-	expression tag	UNP O32504
B	-12	LYS	-	expression tag	UNP O32504
B	-11	LYS	-	expression tag	UNP O32504
B	-10	ALA	-	expression tag	UNP O32504
B	-9	GLY	-	expression tag	UNP O32504
B	-8	LEU	-	expression tag	UNP O32504
B	-7	GLU	-	expression tag	UNP O32504
B	-6	ASN	-	expression tag	UNP O32504
B	-5	LEU	-	expression tag	UNP O32504
B	-4	TYR	-	expression tag	UNP O32504
B	-3	PHE	-	expression tag	UNP O32504
B	-2	GLN	-	expression tag	UNP O32504

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP O32504
B	180	LYS	ASP	engineered mutation	UNP O32504
B	184	LYS	ASP	engineered mutation	UNP O32504
C	-22	MSE	-	initiating methionine	UNP O32504
C	-21	ARG	-	expression tag	UNP O32504
C	-20	SER	-	expression tag	UNP O32504
C	-19	GLY	-	expression tag	UNP O32504
C	-18	SER	-	expression tag	UNP O32504
C	-17	HIS	-	expression tag	UNP O32504
C	-16	HIS	-	expression tag	UNP O32504
C	-15	HIS	-	expression tag	UNP O32504
C	-14	HIS	-	expression tag	UNP O32504
C	-13	HIS	-	expression tag	UNP O32504
C	-12	HIS	-	expression tag	UNP O32504
C	-11	ARG	-	expression tag	UNP O32504
C	-10	SER	-	expression tag	UNP O32504
C	-9	ASP	-	expression tag	UNP O32504
C	-8	ILE	-	expression tag	UNP O32504
C	-7	THR	-	expression tag	UNP O32504
C	-6	SER	-	expression tag	UNP O32504
C	-5	LEU	-	expression tag	UNP O32504
C	-4	TYR	-	expression tag	UNP O32504
C	-3	LYS	-	expression tag	UNP O32504
C	-2	LYS	-	expression tag	UNP O32504
C	-1	ALA	-	expression tag	UNP O32504
C	0	GLY	-	expression tag	UNP O32504
C	1	LEU	-	expression tag	UNP O32504
C	2	GLU	-	expression tag	UNP O32504
C	3	ASN	-	expression tag	UNP O32504
C	4	LEU	-	expression tag	UNP O32504
C	5	TYR	-	expression tag	UNP O32504
C	6	PHE	-	expression tag	UNP O32504
C	7	GLN	-	expression tag	UNP O32504
C	8	GLY	-	expression tag	UNP O32504
C	180	LYS	ASP	engineered mutation	UNP O32504
C	184	LYS	ASP	engineered mutation	UNP O32504
D	-22	MSE	-	initiating methionine	UNP O32504
D	-21	ARG	-	expression tag	UNP O32504
D	-20	SER	-	expression tag	UNP O32504
D	-19	GLY	-	expression tag	UNP O32504
D	-18	SER	-	expression tag	UNP O32504
D	-17	HIS	-	expression tag	UNP O32504

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	HIS	-	expression tag	UNP O32504
D	-15	HIS	-	expression tag	UNP O32504
D	-14	HIS	-	expression tag	UNP O32504
D	-13	HIS	-	expression tag	UNP O32504
D	-12	HIS	-	expression tag	UNP O32504
D	-11	ARG	-	expression tag	UNP O32504
D	-10	SER	-	expression tag	UNP O32504
D	-9	ASP	-	expression tag	UNP O32504
D	-8	ILE	-	expression tag	UNP O32504
D	-7	THR	-	expression tag	UNP O32504
D	-6	SER	-	expression tag	UNP O32504
D	-5	LEU	-	expression tag	UNP O32504
D	-4	TYR	-	expression tag	UNP O32504
D	-3	LYS	-	expression tag	UNP O32504
D	-2	LYS	-	expression tag	UNP O32504
D	-1	ALA	-	expression tag	UNP O32504
D	0	GLY	-	expression tag	UNP O32504
D	1	LEU	-	expression tag	UNP O32504
D	2	GLU	-	expression tag	UNP O32504
D	3	ASN	-	expression tag	UNP O32504
D	4	LEU	-	expression tag	UNP O32504
D	5	TYR	-	expression tag	UNP O32504
D	6	PHE	-	expression tag	UNP O32504
D	7	GLN	-	expression tag	UNP O32504
D	8	GLY	-	expression tag	UNP O32504
D	180	LYS	ASP	engineered mutation	UNP O32504
D	184	LYS	ASP	engineered mutation	UNP O32504

- Molecule 2 is CITRATE ANION (CCD ID: FLC) (formula:  $C_6H_5O_7$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	5	0
			18	6	5	7		
2	C	1	Total	C	H	O	5	0
			18	6	5	7		
2	C	1	Total	C	H	O	5	0
			18	6	5	7		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	103	Total	O	0	0
			103	103		
3	B	38	Total	O	0	0
			38	38		
3	C	25	Total	O	0	0
			25	25		
3	D	5	Total	O	0	0
			5	5		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: DNA repair protein PprA





- Molecule 1: DNA repair protein PprA




- Molecule 1: DNA repair protein PprA



- Molecule 1: DNA repair protein PprA







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.68Å 123.22Å 139.29Å 90.00° 93.69° 90.00°	Depositor
Resolution (Å)	41.65 – 2.07 41.65 – 2.07	Depositor EDS
% Data completeness (in resolution range)	51.5 (41.65-2.07) 51.8 (41.65-2.07)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.92 (at 2.06Å)	Xtriage
Refinement program	PHENIX 2.0_5824	Depositor
R, $R_{free}$	0.211 , 0.236 0.212 , 0.237	Depositor DCC
$R_{free}$ test set	2489 reflections (2.44%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.5	Xtriage
Anisotropy	0.076	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 53.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	16960	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.29	0/2290	0.51	0/3087
1	B	0.29	0/2111	0.50	0/2849
1	C	0.22	0/2101	0.41	0/2833
1	D	0.37	0/2070	0.57	0/2794
All	All	0.30	0/8572	0.50	0/11563

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	196	ARG	Sidechain
1	D	70	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2256	2216	2218	11	0
1	B	2080	2040	2043	10	0
1	C	2070	2034	2034	13	0
1	D	2039	2000	2000	24	0
2	A	13	5	5	1	0
2	C	26	10	10	0	0
3	A	103	0	0	1	0
3	B	38	0	0	0	0
3	C	25	0	0	2	0
3	D	5	0	0	0	0
All	All	8655	8305	8310	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 56 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:27:ASP:OD1	1:D:29:GLN:HG2	1.80	0.81
1:A:186:ILE:HD12	1:A:207:MSE:HE3	1.74	0.69
1:D:13:ILE:HG12	1:D:76:ASP:HA	1.77	0.66
1:A:183:TRP:HA	1:A:207:MSE:HE2	1.78	0.65
1:D:21:MSE:SE	1:D:54:ALA:HB2	2.47	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	294/307 (96%)	288 (98%)	5 (2%)	1 (0%)	37 31
1	B	273/307 (89%)	266 (97%)	6 (2%)	1 (0%)	30 23

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	271/307 (88%)	264 (97%)	7 (3%)	0	100	100
1	D	267/307 (87%)	260 (97%)	7 (3%)	0	100	100
All	All	1105/1228 (90%)	1078 (98%)	25 (2%)	2 (0%)	44	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	214	ALA
1	A	11	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/222 (98%)	216 (100%)	1 (0%)	86	88
1	B	198/222 (89%)	198 (100%)	0	100	100
1	C	197/222 (89%)	197 (100%)	0	100	100
1	D	194/222 (87%)	192 (99%)	2 (1%)	73	73
All	All	806/888 (91%)	803 (100%)	3 (0%)	89	90

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	D	43	LEU
1	D	76	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	9	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FLC	C	301	-	12,12,12	1.45	1 (8%)	17,17,17	1.67	6 (35%)
2	FLC	A	301	-	12,12,12	1.45	1 (8%)	17,17,17	1.81	5 (29%)
2	FLC	C	302	-	12,12,12	1.34	1 (8%)	17,17,17	1.51	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FLC	C	301	-	-	5/16/16/16	-
2	FLC	A	301	-	-	3/16/16/16	-
2	FLC	C	302	-	-	0/16/16/16	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	302	FLC	CB-CBC	2.77	1.56	1.53
2	A	301	FLC	CB-CBC	2.65	1.56	1.53
2	C	301	FLC	CB-CBC	2.40	1.56	1.53

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	301	FLC	OB1-CBC-CB	-3.53	115.25	122.09
2	C	302	FLC	OB1-CBC-CB	-3.24	115.83	122.09
2	A	301	FLC	CA-CB-CBC	-3.15	103.06	110.03
2	A	301	FLC	OB1-CBC-CB	-3.14	116.01	122.09
2	A	301	FLC	CB-CG-CGC	2.92	121.90	113.92

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	FLC	CAC-CA-CB-OHB
2	C	301	FLC	CA-CB-CBC-OB1
2	C	301	FLC	CA-CB-CBC-OB2
2	C	301	FLC	OHB-CB-CBC-OB1
2	C	301	FLC	OHB-CB-CBC-OB2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	FLC	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	292/307 (95%)	0.02	14 (4%) 36 38	7, 16, 39, 55	1 (0%)
1	B	259/307 (84%)	0.79	42 (16%) 5 5	9, 25, 56, 76	1 (0%)
1	C	269/307 (87%)	0.71	24 (8%) 17 18	13, 25, 44, 63	1 (0%)
1	D	252/307 (82%)	2.52	158 (62%) 0 0	18, 51, 75, 110	18 (7%)
All	All	1072/1228 (87%)	0.97	238 (22%) 3 2	7, 26, 64, 110	21 (1%)

The worst 5 of 238 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	233	VAL	6.0
1	B	212	LEU	6.0
1	D	211	LEU	5.2
1	D	194	PHE	5.2
1	D	226	LEU	5.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FLC	A	301	13/13	0.84	0.18	15,81,114,125	5
2	FLC	C	301	13/13	0.84	0.22	15,86,116,124	5
2	FLC	C	302	13/13	0.88	0.19	15,90,115,122	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.