wwPDB X-ray Structure Validation Summary Report (i)

Aug 16, 2023 – 07:31 AM EDT

PDB ID	:	1ZBL
Title	:	Bacillus halodurans RNase H catalytic domain mutant D192N in complex with 12-mer RNA/DNA hybrid
Authors	:	Nowotny, M.; Gaidamakov, S.A.; Crouch, R.J.; Yang, W.
Deposited on	:	2005-04-08
Resolution	:	2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)
RNA backbone	3102	1032 (2.60-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1-C	12	100% 67% 33%
1	2-C	12	100% 67% 33%
2	1-D	12	92% 33% 67%
2	2-D	12	92% 42% 58%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	1-A	139	2% 76% 19%
3	1-B	139	74% 21%
3	2-A	139	2% 78% 18%
3	2-B	139	74% 21%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 5582 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(*GP*AP*CP*AP*CP*CP*UP*GP*AP*UP*UP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	1-C	12	249	113	43	82	11	0	0	0
1	2-C	12	249	113	43	82	11	0	0	0

- Molecule 2 is a DNA chain called 5'-D(*GP*AP*AP*TP*CP*AP*GP*GP*TP*GP*TP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	1-D	12	246	118	47	70	11	0	0	0
2	2-D	12	246	118	47	70	11	0	0	0

- Molecule 3 is a protein called ribonuclease H-related protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	1-A	133	1086	694	181	210	1	0	0	0
3	2-A	133	1086	694	181	210	1	0	0	0
3	1-B	133	1086	694	181	210	1	8	0	0
3	2-B	133	1086	694	181	210	1	8	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	55	GLY	-	cloning artifact	GB 10173478
A	56	SER	-	cloning artifact	GB 10173478
A	57	HIS	-	cloning artifact	GB 10173478

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	58	MET	-	cloning artifact	GB 10173478
A	192	ASN	ASP	engineered mutation	GB 10173478
B	55	GLY	-	cloning artifact	GB 10173478
B	56	SER	-	cloning artifact	GB 10173478
B	57	HIS	-	cloning artifact	GB 10173478
B	58	MET	-	cloning artifact	GB 10173478
B	192	ASN	ASP	engineered mutation	GB 10173478

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	1-A	2	2	2	0	0
4	2-A	2	2	2	0	0
4	1-B	2	2	2	0	0
4	2-B	2	2	2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	1-C	17	17	17	0	0
5	2-C	7	7	7	0	0
5	1-D	11	11	11	0	0
5	2-D	8	8	8	0	0
5	1-A	47	47	47	0	0
5	2-A	50	50	50	0	0
5	1-B	45	45	45	0	0
5	2-B	55	55	55	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 5'-R(*GP*AP*CP*AP*CP*CP*UP*GP*AP*UP*UP*C)-3'

Chain 1-C:



• Molecule 1: 5'-R(*GP*AP*CP*AP*CP*CP*UP*GP*AP*UP*UP*C)-3'

Chain 2-C:



• Molecule 2: 5'-D(*GP*AP*AP*TP*CP*AP*GP*GP*TP*GP*TP*C)-3'

Chain 1-D:





• Molecule 2: 5'-D(*GP*AP*AP*TP*CP*AP*GP*GP*TP*GP*TP*C)-3'

Chain 2-D:





• Molecule 3: ribonuclease H-related protein

Chain 1-A:



• Molecule 3: ribonuclease H-related protein

Chain 1-B:



- Molecule 3: ribonuclease H-related protein

Chain 2-A:



- Molecule 3: ribonuclease H-related protein

Chain 2-B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	36.90Å 44.98Å 62.28Å 83.63° 89.90° 65.69°	Depositor
Resolution (Å)	30.00 – 2.20 33.62 – 2.01	Depositor EDS
% Data completeness (in resolution range)	82.7 (30.00-2.20) 67.7 (33.62-2.01)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	0.84 (at 2.01Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.217 , 0.253 0.248 , 0.271	Depositor DCC
R_{free} test set	1785 reflections (9.87%)	wwPDB-VP
Wilson B-factor (Å^2)	28.5	Xtriage
Anisotropy	0.304	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.51 , 30.1	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.476 for h,h-k,-l	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	5582	wwPDB-VP
Average B, all atoms (Å^2)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.65% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	1-C	0.26	0/277	0.64	0/429
1	2-C	0.28	0/277	0.65	0/429
2	1-D	0.29	0/276	0.62	0/425
2	2-D	0.30	0/276	0.61	0/425
3	1-A	0.43	0/1111	0.63	0/1505
3	1-B	0.42	0/1111	0.62	0/1505
3	2-A	0.43	0/1111	0.63	0/1505
3	2-B	0.42	0/1111	0.62	0/1505
All	All	0.40	0/5550	0.63	0/7728

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1-C	249	0	131	2	0
1	2-C	249	0	131	2	0
2	1-D	246	0	137	7	0
2	2-D	246	0	137	4	0
3	1-A	1086	0	1074	21	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	1-B	1086	0	1074	21	0
3	2-A	1086	0	1074	19	0
3	2-B	1086	0	1074	21	0
4	1-A	2	0	0	0	0
4	1-B	2	0	0	0	0
4	2-A	2	0	0	0	0
4	2-B	2	0	0	0	0
5	1-A	47	0	0	0	0
5	1-B	45	0	0	1	0
5	1-C	17	0	0	0	0
5	1-D	11	0	0	2	0
5	2-A	50	0	0	0	0
5	2-B	55	0	0	1	0
5	2-C	7	0	0	0	0
5	2-D	8	0	0	1	0
All	All	5582	0	4832	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 95 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:128:PRO:HB3	3:B:177:PRO:HG2	1.64	0.79
3:B:128:PRO:HB3	3:B:177:PRO:HG2	1.64	0.79
3:A:118:ARG:HG2	3:A:168:TRP:CE2	2.21	0.76
3:A:118:ARG:HG2	3:A:168:TRP:CE2	2.21	0.76
3:A:101:PRO:O	3:A:102:ILE:HD13	1.87	0.74

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	1-A	131/139 (94%)	127 (97%)	4 (3%)	0	100	100
3	1-B	131/139 (94%)	125 (95%)	6 (5%)	0	100	100
3	2-A	131/139 (94%)	127 (97%)	4 (3%)	0	100	100
3	2-B	131/139 (94%)	125 (95%)	6 (5%)	0	100	100
All	All	524/556 (94%)	504 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	1-A	118/122 (97%)	117 (99%)	1 (1%)	81	90
3	1-B	118/122 (97%)	115 (98%)	3 (2%)	47	60
3	2-A	118/122 (97%)	117 (99%)	1 (1%)	81	90
3	2-B	118/122 (97%)	115 (98%)	3 (2%)	47	60
All	All	472/488 (97%)	464 (98%)	8 (2%)	60	74

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	2-B	190	LYS
3	2-B	117	LEU
3	2-A	81	VAL
3	1-B	190	LYS
3	2-B	81	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
3	1-A	115	HIS
3	1-B	115	HIS
3	2-A	115	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	2-B	115	HIS

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	1-C	11/12 (91%)	0	0
1	2-C	11/12 (91%)	0	0
All	All	22/24 (91%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	1-C	12/12 (100%)	3.29	12 (100%)	0	0	31, 39, 52, 57	12 (100%)
1	2-C	12/12 (100%)	3.29	12 (100%)	0	0	31, 39, 52, 57	12 (100%)
2	1-D	12/12 (100%)	3.09	11 (91%)	0	0	30, 46, 62, 68	12 (100%)
2	2-D	12/12 (100%)	3.09	11 (91%)	0	0	30, 46, 62, 68	12 (100%)
3	1-A	133/139 (95%)	-0.26	3 (2%)	60	58	19, 38, 55, 65	0
3	1-B	133/139 (95%)	-0.28	0	100	100	20, 38, 56, 61	2 (1%)
3	2-A	133/139 (95%)	-0.26	3 (2%)	60	58	19, 38, 55, 65	0
3	2-B	133/139 (95%)	-0.28	0	100	100	20, 38, 56, 61	2 (1%)
All	All	580/604 (96%)	0.02	52 (8%)	9	8	19, 39, 57, 68	52 (8%)

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	1-C	9	A	4.1
1	2-C	9	A	4.1
2	1-D	13	DG	3.9
2	2-D	13	DG	3.9
2	1-D	19	DG	3.9

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	MG	1-B	203	1/1	0.93	0.21	31,31,31,31	0
4	MG	2-B	203	1/1	0.93	0.21	31,31,31,31	0
4	MG	1-B	204	1/1	0.96	0.05	30,30,30,30	0
4	MG	2-B	204	1/1	0.96	0.05	30,30,30,30	0
4	MG	1-A	201	1/1	0.97	0.16	33,33,33,33	0
4	MG	2-A	201	1/1	0.97	0.16	33,33,33,33	0
4	MG	1-A	202	1/1	0.98	0.05	33,33,33,33	0
4	MG	2-A	202	1/1	0.98	0.05	33,33,33,33	0

6.5 Other polymers (i)

There are no such residues in this entry.