



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2026 – 02:23 PM EDT

PDB ID : 12QA / pdb_000012qa
Title : Crystal structure of CM10 Fab in complex with an orthommarburgvirus GP2 peptide
Authors : Niyongabo, A.; Janus, B.M.; Metcalf, M.; Ofek, G.
Deposited on : 2026-04-14
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	219	<div> <div>2%</div> <div> <div></div> <div>91%</div> <div>9%</div> </div> </div>
2	H	226	<div> <div>5%</div> <div> <div></div> <div>88%</div> <div>12%</div> </div> </div>
3	P	15	<div> <div>13%</div> <div> <div></div> <div>47%</div> <div>13%</div> <div>40%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7203 atoms, of which 3366 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CM10 Fab light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	L	219	Total	C	H	N	O	S	0	5	0
			3378	1072	1670	286	343	7			

- Molecule 2 is a protein called CM10 Fab heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	H	226	Total	C	H	N	O	S	0	3	0
			3293	1043	1631	285	327	7			

- Molecule 3 is a protein called GP2 peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	9	Total	C	H	N	O	0	0	0
			137	47	65	10	15			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	1	Total	Ca	0	0
			1	1		
4	H	1	Total	Ca	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	219	Total	O	0	0
			219	219		
5	H	160	Total	O	0	0
			160	160		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	P	14	Total	O	0	0
			14	14		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CM10 Fab light chain



- Molecule 2: CM10 Fab heavy chain



- Molecule 3: GP2 peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.22Å 41.69Å 80.64Å 90.00° 102.17° 90.00°	Depositor
Resolution (Å)	36.85 – 1.60 36.85 – 1.60	Depositor EDS
% Data completeness (in resolution range)	87.5 (36.85-1.60) 87.5 (36.85-1.60)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.20	Depositor
R, R_{free}	0.175 , 0.207 0.177 , 0.211	Depositor DCC
R_{free} test set	1879 reflections (3.24%)	wwPDB-VP
Wilson B-factor (Å ²)	17.7	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7203	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	1.04	4/1766 (0.2%)	0.85	0/2401
2	H	1.11	10/1716 (0.6%)	0.84	0/2335
3	P	0.92	0/74	0.88	0/102
All	All	1.07	14/3556 (0.4%)	0.85	0/4838

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	200	LYS	C-O	-7.88	1.17	1.24
1	L	88	CYS	C-O	-6.89	1.15	1.23
2	H	145	PHE	C-O	-6.82	1.16	1.24
2	H	93	ARG	C-O	-6.78	1.15	1.23
1	L	53	ASN	C-O	-6.17	1.16	1.24
2	H	92	SER	C-O	-5.82	1.16	1.23
2	H	89	TYR	C-O	-5.82	1.17	1.24
2	H	198	ASN	C-O	-5.78	1.17	1.24
2	H	72	ASN	C-O	-5.44	1.17	1.24
1	L	209	PHE	C-O	-5.39	1.17	1.23
1	L	207	LYS	C-O	-5.33	1.17	1.24
2	H	94	GLY	C-O	-5.22	1.16	1.23
2	H	102	TRP	C-O	-5.19	1.17	1.23
2	H	74	LYS	C-O	-5.03	1.17	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1708	1670	1650	11	0
2	H	1662	1631	1613	13	0
3	P	72	65	65	4	0
4	H	1	0	0	0	0
4	L	1	0	0	0	0
5	H	160	0	0	0	2
5	L	219	0	0	1	3
5	P	14	0	0	0	0
All	All	3837	3366	3328	23	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:126:SER:O	2:H:130:THR:HG23	1.92	0.68
1:L:80:ALA:HA	1:L:106:ILE:CD1	2.26	0.65
2:H:57:TYR:CE2	3:P:461:ILE:HD13	2.33	0.63
1:L:96:TRP:CZ3	2:H:100(e):LEU:HD11	2.36	0.61
2:H:6:GLU:HG2	2:H:22:CYS:SG	2.44	0.57
2:H:97:PRO:HG3	3:P:456:ILE:HG21	1.87	0.56
1:L:2:ILE:HG12	1:L:27:GLN:HG3	1.92	0.51
1:L:96:TRP:HZ3	2:H:100(e):LEU:HD11	1.79	0.47
1:L:175:LEU:C	1:L:175:LEU:HD23	2.42	0.45
1:L:107:LYS:HA	1:L:140:TYR:OH	2.17	0.44
2:H:99:GLY:HA2	2:H:100(b):ARG:HG3	2.00	0.43
2:H:57:TYR:CE2	3:P:461:ILE:CD1	3.01	0.42
1:L:145:LYS:HB3	1:L:197:THR:HB	2.01	0.42
2:H:128:LYS:HG2	2:H:215:CYS:SG	2.60	0.42
3:P:461:ILE:N	3:P:461:ILE:HD12	2.35	0.42
2:H:21:SER:HA	2:H:77:LEU:O	2.19	0.41
1:L:210:ASN:O	1:L:213[A]:GLU:HG2	2.20	0.41
2:H:17:SER:HA	2:H:81:MET:O	2.20	0.41
2:H:127:SER:OG	2:H:215:CYS:O	2.38	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:37:LEU:HD13	1:L:86:TYR:CZ	2.56	0.41
1:L:204:PRO:HD2	5:L:414:HOH:O	2.20	0.41
2:H:18:LEU:HD12	2:H:18:LEU:C	2.45	0.41
1:L:80:ALA:HA	1:L:106:ILE:HD12	2.02	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:L:423:HOH:O	5:L:496:HOH:O[2_546]	1.73	0.47
5:L:544:HOH:O	5:H:542:HOH:O[2_556]	1.90	0.30
5:L:569:HOH:O	5:H:472:HOH:O[2_556]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	222/219 (101%)	217 (98%)	5 (2%)	0	100	100
2	H	227/226 (100%)	225 (99%)	2 (1%)	0	100	100
3	P	7/15 (47%)	6 (86%)	1 (14%)	0	100	100
All	All	456/460 (99%)	448 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	198/193 (103%)	198 (100%)	0	100	100
2	H	188/186 (101%)	188 (100%)	0	100	100
3	P	9/14 (64%)	9 (100%)	0	100	100
All	All	395/393 (100%)	395 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	93	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	219/219 (100%)	0.05	5 (2%) 61 64	12, 22, 44, 76	3 (1%)
2	H	226/226 (100%)	0.18	12 (5%) 32 33	11, 23, 55, 84	1 (0%)
3	P	9/15 (60%)	0.64	2 (22%) 2 2	19, 20, 41, 51	0
All	All	454/460 (98%)	0.12	19 (4%) 40 42	11, 22, 51, 84	4 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	215	CYS	6.0
1	L	96	TRP	4.4
1	L	194	CYS	4.0
2	H	130	THR	3.7
2	H	131	SER	3.5
3	P	456	ILE	3.2
3	P	464	THR	2.9
2	H	132	GLY	2.7
1	L	214	CYS	2.7
1	L	69	THR	2.5
2	H	190	THR	2.4
2	H	160	SER	2.3
1	L	106	ILE	2.3
2	H	189	GLY	2.3
2	H	192	THR	2.3
2	H	22	CYS	2.3
2	H	214	SER	2.2
2	H	158	LEU	2.2
2	H	157	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CA	H	301	1/1	0.93	0.09	58,58,58,58	0
4	CA	L	301	1/1	0.97	0.07	27,27,27,27	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CA H 301:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.