



wwPDB EM Validation Summary Report ⓘ

Jul 10, 2026 – 03:34 PM JST

PDB ID : 24QP / pdb_000024qp
EMDB ID : EMD-69770
Title : Structure of yeast Pol II in complex with an oxaliplatin-ICL lesion at +7 position.
Authors : Xu, J.; Zhu, L.
Deposited on : 2026-03-17
Resolution : 2.73 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at
<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev133
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	10432 (2.23 - 3.23)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1733	
2	B	1224	
3	C	318	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	221	
5	E	215	
6	F	155	
7	G	171	
8	H	146	
9	I	122	
10	J	70	
11	K	120	
12	L	70	
13	N	18	
14	P	6	
15	Q	64	
16	T	25	

2 Entry composition

There are 20 unique types of molecules in this entry. The entry contains 33308 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1424	Total	C	N	O	S	0	0
			11205	7060	1958	2125	62		

- Molecule 2 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	1187	Total	C	N	O	S	0	0
			9453	5973	1656	1768	56		

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	265	Total	C	N	O	S	0	0
			2086	1312	347	414	13		

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB4.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	165	Total	C	N	O	S	0	0
			1326	820	237	267	2		

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	214	Total	C	N	O	S	0	0
			1752	1111	309	321	11		

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	87	Total	C	N	O	S	0	0
			705	451	119	132	3		

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	171	Total	C	N	O	S	0	0
			1339	861	222	248	8		

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	139	Total	C	N	O	S	0	0
			1113	699	187	223	4		

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	I	116	Total	C	N	O	S	0	0
			944	581	172	181	10		

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	J	69	Total	C	N	O	S	0	0
			569	362	101	100	6		

- Molecule 11 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	K	115	Total	C	N	O	S	0	0
			924	593	157	172	2		

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	L	45	Total	C	N	O	S	0	0
			359	221	71	63	4		

- Molecule 13 is a DNA chain called DNA (NTS, non-template strand).

Mol	Chain	Residues	Atoms					AltConf	Trace
13	N	18	Total	C	N	O	P	0	0
			385	179	88	100	18		

- Molecule 14 is a RNA chain called RNA (5'-R(P*GP*GP*GP*AP*AP*A)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
14	P	6	Total	C	N	O	P	0	0
			135	60	30	39	6		

- Molecule 15 is a protein called Transcription elongation factor 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	Q	64	Total	C	N	O	S	0	0
			505	316	84	100	5		

- Molecule 16 is a DNA chain called DNA (TS, template strand).

Mol	Chain	Residues	Atoms					AltConf	Trace
16	T	25	Total	C	N	O	P	0	0
			489	237	66	161	25		


- Molecule 17 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
17	A	2	Total	Zn	0
			2	2	
17	B	1	Total	Zn	0
			1	1	
17	C	1	Total	Zn	0
			1	1	
17	I	2	Total	Zn	0
			2	2	
17	J	1	Total	Zn	0
			1	1	
17	L	1	Total	Zn	0
			1	1	
17	Q	1	Total	Zn	0
			1	1	

- Molecule 18 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
18	A	1	Total	Mg	0
			1	1	

- Molecule 19 is 1R,2R-DIAMINOCYCLOHEXANE (CCD ID: DNH) (formula: C₆H₁₄N₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
19	N	1	Total	C	N	0
			8	6	2	


- Molecule 20 is PLATINUM (II) ION (CCD ID: PT) (formula: Pt) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
20	N	1	Total	Pt	0
			1	1	

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA-directed RNA polymerase II subunit RPB1






- Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2

Chain F: 

MET SER ASP GLN TYR GLU ALA PHE ASN ASP GLY ASN GLU ASN PHE GLU ASP PHE ASP VAL GLU HIS PHE SER ASP GLU THR TYR GLU LYS PRO GLN PHE LYS ASP GLY THR ASP ALA ASN GLY LYS THR ILE VAL THR GLY GLY ASN GLY PRO ASP PHE GLN

HIS GLU GLN ILE ARG LYS THR L69 P75 K76 D77 Q78 R79 M85 E89 V133 I134 R135 R136 F139 W146 L156

- Molecule 7: DNA-directed RNA polymerase II subunit RPB7

Chain G: 

M1 F2 I11 H14 P15 K23 L26 E32 E35 L46 L49 D52 R53 R59 I61 K73 V77 K80 P81 F82 K83 G84 E85 V86 V87 D88 G89 T90 V91 V92 S93 C94 S95 Q96 H97 G98 F99 E100 V101 Q102 V103 G104 K107 V108 F109

V110 T111 K112 H113 I114 M115 P116 Q117 D118 L119 T120 F121 M122 A123 G124 S125 M126 P127 S128 S129 Y130 Q131 S132 S133 E134 D135 V136 I137 T138 I139 K140 S141 R142 T143 R144 V145 K146 T147 E148 G149 C150 I151 S152 Q153 V154 H158 A159 I160 G161 S162 I163 K164 E165 D166 Y167 L168 G169 A170 I171

- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3

Chain H: 

MET S2 R3 T4 L5 F6 D7 F10 V15 Y20 I26 E27 A28 L38 T39 L40 L46 L49 L59 L63 D67 T68 PRO ALA ASN ASP SER A75 Q83 D91 N97 Y102 K103 Y115 Y116 L122 M123 K124 L125 R130 K136 Q137 E138

L142 L143 R146

- Molecule 9: DNA-directed RNA polymerase II subunit RPB9

Chain I: 

MET T2 R5 D9 C10 N11 Y15 E21 E27 R30 P41 R45 H46 T49 Q60 E74 F85 T95 F101 V102 I109 S112 K117 ARG THR GLN PHE SER

- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5

Chain J: 


H1 I2 V3 P4 V5 R6 C7 V13 K17 W18 L36 L39 G40 L41 Y44 C45 C46 R47 R48 M49 L56 F60 R69 ASP

- Molecule 11: DNA-directed RNA polymerase II subunit RPB11


Chain K: 



- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4





- Molecule 13: DNA (NTS, non-template strand)



- Molecule 14: RNA (5'-R(P*GP*GP*GP*AP*AP*A)-3')



- Molecule 15: Transcription elongation factor 1



- Molecule 16: DNA (TS, template strand)



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	156371	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.802	Depositor
Minimum map value	-0.415	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.027	Depositor
Recommended contour level	0.1	Depositor
Map size (Å)	297.6, 297.6, 297.6	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.93, 0.93, 0.93	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PT, MG, DNH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.16	0/11406	0.38	3/15423 (0.0%)
2	B	0.22	0/9640	0.47	8/13002 (0.1%)
3	C	0.11	0/2124	0.28	0/2879
4	D	0.15	0/1334	0.42	0/1789
5	E	0.12	0/1788	0.32	0/2406
6	F	0.11	0/717	0.32	0/967
7	G	0.10	0/1367	0.30	0/1844
8	H	0.08	0/1131	0.23	0/1532
9	I	0.10	0/962	0.31	0/1295
10	J	0.11	0/578	0.29	0/775
11	K	0.12	0/942	0.27	0/1272
12	L	0.11	0/361	0.32	0/478
13	N	0.46	0/437	0.91	0/675
14	P	0.10	0/152	0.21	0/236
15	Q	0.08	0/513	0.22	0/690
16	T	0.53	0/539	1.10	1/825 (0.1%)
All	All	0.18	0/33991	0.42	12/46088 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	2
All	All	0	3

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1132	GLU	CB-CG-CD	-8.93	97.42	112.60
2	B	502	ILE	CA-C-N	-7.62	115.75	121.61
2	B	502	ILE	C-N-CA	-7.62	115.75	121.61
2	B	1130	PHE	N-CA-CB	7.00	120.39	110.24
2	B	1130	PHE	CA-CB-CG	6.34	120.14	113.80

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	532	ARG	Sidechain
2	B	504	ARG	Sidechain
2	B	512	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11205	0	11277	208	0
2	B	9453	0	9455	179	0
3	C	2086	0	2045	41	0
4	D	1326	0	1348	37	0
5	E	1752	0	1776	40	0
6	F	705	0	731	8	0
7	G	1339	0	1357	21	0
8	H	1113	0	1079	19	0
9	I	944	0	899	11	0
10	J	569	0	585	16	0
11	K	924	0	934	21	0
12	L	359	0	381	5	0
13	N	385	0	199	12	0
14	P	135	0	66	2	0
15	Q	505	0	491	5	0
16	T	489	0	287	16	0
17	A	2	0	0	0	0
17	B	1	0	0	0	0
17	C	1	0	0	0	0
17	I	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
17	J	1	0	0	0	0
17	L	1	0	0	0	0
17	Q	1	0	0	0	0
18	A	1	0	0	0	0
19	N	8	0	13	0	0
20	N	1	0	0	0	0
All	All	33308	0	32923	560	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 560 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:LYS:HB3	2:B:1220:ARG:HE	1.41	0.84
16:T:8:DT:H2"	16:T:9:DC:H5"	1.59	0.84
2:B:1152:MET:HE1	2:B:1195:HIS:HB3	1.61	0.81
5:E:100:ILE:HG23	5:E:105:PHE:HB2	1.62	0.81
2:B:593:PRO:HG2	2:B:617:ARG:HH21	1.46	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1416/1733 (82%)	1385 (98%)	31 (2%)	0	100	100
2	B	1179/1224 (96%)	1136 (96%)	40 (3%)	3 (0%)	36	54
3	C	263/318 (83%)	252 (96%)	11 (4%)	0	100	100
4	D	161/221 (73%)	156 (97%)	5 (3%)	0	100	100
5	E	212/215 (99%)	206 (97%)	6 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
6	F	85/155 (55%)	82 (96%)	3 (4%)	0	100	100
7	G	169/171 (99%)	161 (95%)	8 (5%)	0	100	100
8	H	135/146 (92%)	134 (99%)	1 (1%)	0	100	100
9	I	114/122 (93%)	108 (95%)	6 (5%)	0	100	100
10	J	67/70 (96%)	65 (97%)	2 (3%)	0	100	100
11	K	113/120 (94%)	111 (98%)	2 (2%)	0	100	100
12	L	43/70 (61%)	43 (100%)	0	0	100	100
15	Q	62/64 (97%)	62 (100%)	0	0	100	100
All	All	4019/4629 (87%)	3901 (97%)	115 (3%)	3 (0%)	49	69

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	505	ASP
2	B	506	GLY
2	B	1118	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1246/1520 (82%)	1245 (100%)	1 (0%)	88	94
2	B	1029/1061 (97%)	1024 (100%)	5 (0%)	81	88
3	C	233/274 (85%)	233 (100%)	0	100	100
4	D	147/200 (74%)	147 (100%)	0	100	100
5	E	196/197 (100%)	196 (100%)	0	100	100
6	F	77/137 (56%)	77 (100%)	0	100	100
7	G	152/152 (100%)	152 (100%)	0	100	100
8	H	122/128 (95%)	122 (100%)	0	100	100
9	I	110/116 (95%)	110 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
10	J	64/65 (98%)	64 (100%)	0	100	100
11	K	99/102 (97%)	99 (100%)	0	100	100
12	L	40/57 (70%)	40 (100%)	0	100	100
15	Q	61/61 (100%)	61 (100%)	0	100	100
All	All	3576/4070 (88%)	3570 (100%)	6 (0%)	85	94

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	1123	SER
2	B	1124	ARG
2	B	1128	LEU
2	B	505	ASP
1	A	336	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	325	GLN
7	G	122	ASN
2	B	975	GLN
9	I	22	ASN
4	D	146	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
14	P	5/6 (83%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 12 ligands modelled in this entry, 11 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
19	DNH	N	101	20	8,8,8	0.48	0	10,10,10	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
19	DNH	N	101	20	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-69770. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 148




Y Index: 141




Z Index: 163


6.3.2 Raw map



X Index: 148



Y Index: 141




Z Index: 163


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.1. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_69770_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 243 nm³; this corresponds to an approximate mass of 220 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.366 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.366 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.73	-	-
Author-provided FSC curve	2.73	3.13	2.77
Unmasked-calculated*	3.33	4.00	3.40

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.33 differs from the reported value 2.73 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-69770 and PDB model 24QP. Per-residue inclusion information can be found in section [3](#) on page [8](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.1 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.1).

9.4 Atom inclusion [i](#)



At the recommended contour level, 90% of all backbone atoms, 84% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.1) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8390	 0.4990
A	 0.8920	 0.5330
B	 0.8730	 0.5230
C	 0.8990	 0.5370
D	 0.3440	 0.2200
E	 0.8680	 0.4750
F	 0.9160	 0.5590
G	 0.4620	 0.3110
H	 0.8840	 0.5380
I	 0.8020	 0.4480
J	 0.9040	 0.5340
K	 0.9220	 0.5670
L	 0.8420	 0.4800
N	 0.8680	 0.4190
P	 0.9560	 0.5360
Q	 0.5260	 0.3140
T	 0.8510	 0.4380

