



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 30, 2024 – 11:29 pm BST

PDB ID : 5A1S
Title : Crystal structure of the sodium-dependent citrate symporter SeCitS form *Salmonella enterica*.
Authors : Woehlert, D.; Groetzinger, M.J.; Kuhlbrandt, W.; Yildiz, O.
Deposited on : 2015-05-05
Resolution : 2.50 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	 10% 85% 12% •
1	B	448	 18% 80% 15% 5%
1	C	448	 13% 79% 16% ••
1	D	448	 15% 82% 14% ••

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FLC	B	1449	-	-	-	X
4	BOG	A	1456	-	-	-	X
4	BOG	C	1448	-	-	-	X
4	BOG	D	1454	-	-	-	X
7	PTY	A	1455	-	-	-	X

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 13270 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CITRATE-SODIUM SYMPORTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	433	3256	2151	516	567	22	0	0	0
1	B	426	3201	2117	508	554	22	0	0	0
1	C	428	3206	2118	511	555	22	0	0	0
1	D	434	3253	2147	517	567	22	0	0	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	447	LEU	-	expression tag	UNP G4BX92
A	448	GLU	-	expression tag	UNP G4BX92
B	447	LEU	-	expression tag	UNP G4BX92
B	448	GLU	-	expression tag	UNP G4BX92
C	447	LEU	-	expression tag	UNP G4BX92
C	448	GLU	-	expression tag	UNP G4BX92
D	447	LEU	-	expression tag	UNP G4BX92
D	448	GLU	-	expression tag	UNP G4BX92

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Na 2	0	0
2	B	1	Total 1	Na 1	0	0
2	C	2	Total 2	Na 2	0	0
2	D	2	Total 2	Na 2	0	0

- Molecule 3 is CITRATE ANION (three-letter code: FLC) (formula: $C_6H_5O_7$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 13 6 7	0	0
3	A	1	Total C O 13 6 7	0	0
3	A	1	Total C O 13 6 7	0	0
3	B	1	Total C O 13 6 7	0	0
3	B	1	Total C O 13 6 7	0	0
3	C	1	Total C O 13 6 7	0	0
3	D	1	Total C O 13 6 7	0	0

- Molecule 4 is octyl beta-D-glucopyranoside (three-letter code: BOG) (formula: $C_{14}H_{28}O_6$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C O	0	0
			20	14 6		
4	A	1	Total	C O	0	0
			20	14 6		
4	C	1	Total	C O	0	0
			20	14 6		
4	C	1	Total	C O	0	0
			20	14 6		
4	D	1	Total	C O	0	0
			20	14 6		

- Molecule 5 is UNDECANE (three-letter code: UND) (formula: $C_{11}H_{24}$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C 11 11	0	0
5	D	1	Total C 11 11	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0
6	D	2	Total Cl 2 2	0	0

- Molecule 7 is PHOSPHATIDYLETHANOLAMINE (three-letter code: PTY) (formula: C₄₀H₈₀NO₈P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C N O P 50 40 1 8 1	0	0

- Molecule 8 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
8	D	1	12	6	1	4	1	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	24	Total	O	0	0
			24	24		
9	B	14	Total	O	0	0
			14	14		
9	C	13	Total	O	0	0
			13	13		
9	D	18	Total	O	0	0
			18	18		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CITRATE-SODIUM SYMPORTER




• Molecule 1: CITRATE-SODIUM SYMPORTER





• Molecule 1: CITRATE-SODIUM SYMPORTER





● Molecule 1: CITRATE-SODIUM SYMPORTER



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	86.38Å 89.94Å 91.84Å 90.44° 113.79° 99.55°	Depositor
Resolution (Å)	47.98 – 2.50 47.98 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.6 (47.98-2.50) 98.7 (47.98-2.50)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 2.51Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.208 , 0.246 0.209 , 0.248	Depositor DCC
R_{free} test set	4200 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	62.4	Xtrriage
Anisotropy	0.118	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 64.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	13270	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PTY, BOG, NA, UND, MES, CL, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/3320	0.40	0/4503
1	B	0.23	0/3263	0.41	0/4425
1	C	0.23	0/3268	0.41	0/4430
1	D	0.23	0/3316	0.40	0/4498
All	All	0.23	0/13167	0.40	0/17856

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3256	0	3414	34	0
1	B	3201	0	3365	36	0
1	C	3206	0	3369	39	0
1	D	3253	0	3407	34	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	39	0	15	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	26	0	10	2	0
3	C	13	0	5	0	0
3	D	13	0	5	1	0
4	A	40	0	56	2	0
4	C	40	0	56	0	0
4	D	20	0	28	1	0
5	A	11	0	24	0	0
5	D	11	0	24	0	0
6	A	1	0	0	0	0
6	D	2	0	0	0	0
7	A	50	0	79	4	0
8	D	12	0	13	2	0
9	A	24	0	0	3	0
9	B	14	0	0	0	0
9	C	13	0	0	0	0
9	D	18	0	0	0	0
All	All	13270	0	13870	145	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 145 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:136:LEU:HD21	1:C:250:ALA:HB1	1.65	0.78
1:A:340:LEU:HD21	1:A:408:LEU:HD13	1.66	0.77
1:C:340:LEU:HD21	1:C:408:LEU:HD13	1.71	0.73
1:A:67:GLU:HB3	1:A:71:ARG:HH21	1.55	0.72
1:C:288:LYS:NZ	1:D:44:TYR:O	2.26	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	431/448 (96%)	412 (96%)	19 (4%)	0	100	100
1	B	422/448 (94%)	395 (94%)	24 (6%)	3 (1%)	22	39
1	C	424/448 (95%)	402 (95%)	20 (5%)	2 (0%)	29	48
1	D	432/448 (96%)	409 (95%)	22 (5%)	1 (0%)	47	68
All	All	1709/1792 (95%)	1618 (95%)	85 (5%)	6 (0%)	34	54

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	259	LYS
1	B	73	PRO
1	D	143	GLY
1	B	186	ASN
1	C	260	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/354 (97%)	339 (99%)	4 (1%)	71	88
1	B	337/354 (95%)	330 (98%)	7 (2%)	53	78
1	C	336/354 (95%)	329 (98%)	7 (2%)	53	78
1	D	341/354 (96%)	330 (97%)	11 (3%)	39	65
All	All	1357/1416 (96%)	1328 (98%)	29 (2%)	53	78

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	132	VAL
1	D	327	ARG
1	C	402	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	263	ILE
1	C	353	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	262	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 10 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FLC	C	1446	-	12,12,12	1.25	0	17,17,17	1.42	1 (5%)
4	BOG	C	1447	-	20,20,20	0.53	0	25,25,25	0.87	1 (4%)
4	BOG	A	1450	-	20,20,20	0.57	0	25,25,25	0.99	1 (4%)
3	FLC	B	1450	-	12,12,12	1.21	0	17,17,17	1.45	1 (5%)
3	FLC	B	1449	-	12,12,12	1.20	0	17,17,17	1.45	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FLC	A	1449	-	12,12,12	1.26	0	17,17,17	1.38	1 (5%)
3	FLC	A	1454	-	12,12,12	1.26	0	17,17,17	1.49	1 (5%)
3	FLC	A	1453	-	12,12,12	1.18	0	17,17,17	1.44	1 (5%)
3	FLC	D	1449	-	12,12,12	1.24	0	17,17,17	1.50	1 (5%)
4	BOG	A	1456	-	20,20,20	0.47	0	25,25,25	1.02	1 (4%)
5	UND	A	1451	-	10,10,10	0.10	0	9,9,9	0.80	0
4	BOG	C	1448	-	20,20,20	0.50	0	25,25,25	0.87	1 (4%)
4	BOG	D	1454	-	20,20,20	0.42	0	25,25,25	1.37	2 (8%)
7	PTY	A	1455	-	49,49,49	1.02	3 (6%)	52,54,54	1.09	3 (5%)
5	UND	D	1450	-	10,10,10	0.10	0	9,9,9	0.81	0
8	MES	D	1453	-	12,12,12	1.28	2 (16%)	14,16,16	2.04	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FLC	C	1446	-	-	0/16/16/16	-
4	BOG	C	1447	-	-	4/11/31/31	0/1/1/1
4	BOG	A	1450	-	-	5/11/31/31	0/1/1/1
3	FLC	B	1450	-	-	6/16/16/16	-
3	FLC	B	1449	-	-	8/16/16/16	-
3	FLC	A	1449	-	-	0/16/16/16	-
3	FLC	A	1454	-	-	5/16/16/16	-
3	FLC	A	1453	-	-	8/16/16/16	-
3	FLC	D	1449	-	-	7/16/16/16	-
4	BOG	A	1456	-	-	7/11/31/31	0/1/1/1
5	UND	A	1451	-	-	0/8/8/8	-
4	BOG	C	1448	-	-	3/11/31/31	0/1/1/1
4	BOG	D	1454	-	-	2/11/31/31	0/1/1/1
7	PTY	A	1455	-	-	23/53/53/53	-
5	UND	D	1450	-	-	2/8/8/8	-
8	MES	D	1453	-	-	3/6/14/14	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1455	PTY	O4-C30	3.60	1.43	1.33
7	A	1455	PTY	O7-C8	3.21	1.43	1.34
8	D	1453	MES	O2S-S	2.40	1.52	1.45
7	A	1455	PTY	O7-C6	-2.22	1.41	1.46
8	D	1453	MES	O1S-S	2.03	1.51	1.45

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	D	1453	MES	O3S-S-O2S	-4.32	100.72	111.27
4	D	1454	BOG	O5-C5-C4	4.29	117.49	109.69
3	D	1449	FLC	OB2-CBC-CB	4.21	120.35	113.05
7	A	1455	PTY	O7-C8-C11	4.16	120.46	111.50
3	C	1446	FLC	OB2-CBC-CB	3.96	119.93	113.05

There are no chirality outliers.

5 of 83 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1453	FLC	CAC-CA-CB-CBC
3	A	1453	FLC	CAC-CA-CB-CG
3	A	1454	FLC	CAC-CA-CB-CBC
3	A	1454	FLC	CAC-CA-CB-CG
3	A	1454	FLC	CAC-CA-CB-OHB


There are no ring outliers.



7 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1450	BOG	2	0
3	B	1450	FLC	1	0
3	B	1449	FLC	1	0
3	D	1449	FLC	1	0
4	D	1454	BOG	1	0
7	A	1455	PTY	4	0
8	D	1453	MES	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	433/448 (96%)	0.50	44 (10%) 6 6	34, 56, 99, 136	0
1	B	426/448 (95%)	0.87	79 (18%) 1 1	46, 80, 120, 132	0
1	C	428/448 (95%)	0.62	59 (13%) 2 2	38, 62, 109, 143	0
1	D	434/448 (96%)	0.74	67 (15%) 2 1	38, 59, 101, 142	0
All	All	1721/1792 (96%)	0.68	249 (14%) 2 2	34, 63, 110, 143	0

The worst 5 of 249 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	16	SER	9.6
1	A	360	PHE	9.1
1	B	17	ASP	8.5
1	B	166	PHE	7.8
1	B	244	GLY	7.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	FLC	B	1449	13/13	0.52	0.48	76,87,98,105	0
3	FLC	A	1453	13/13	0.56	0.25	79,108,121,121	0
7	PTY	A	1455	50/50	0.60	0.65	66,90,101,110	50
4	BOG	A	1456	20/20	0.64	0.56	71,113,125,131	0
3	FLC	A	1454	13/13	0.64	0.40	98,120,125,128	0
8	MES	D	1453	12/12	0.68	0.23	75,103,106,113	0
4	BOG	C	1448	20/20	0.69	0.42	80,99,115,118	0
4	BOG	D	1454	20/20	0.71	0.41	84,105,112,113	0
3	FLC	B	1450	13/13	0.79	0.24	98,108,114,116	0
5	UND	D	1450	11/11	0.80	0.35	55,61,81,83	0
3	FLC	D	1449	13/13	0.87	0.29	68,82,91,93	0
6	CL	A	1452	1/1	0.89	0.07	108,108,108,108	0
5	UND	A	1451	11/11	0.90	0.26	55,62,68,75	0
2	NA	D	502	1/1	0.91	0.21	50,50,50,50	0
2	NA	C	501	1/1	0.92	0.23	54,54,54,54	0
4	BOG	A	1450	20/20	0.92	0.22	44,57,70,77	0
2	NA	B	502	1/1	0.93	0.14	72,72,72,72	0
2	NA	A	501	1/1	0.94	0.11	39,39,39,39	0
2	NA	C	502	1/1	0.95	0.12	56,56,56,56	0
6	CL	D	1451	1/1	0.95	0.14	62,62,62,62	0
4	BOG	C	1447	20/20	0.95	0.17	57,63,76,76	0
3	FLC	C	1446	13/13	0.95	0.20	44,57,65,66	0
2	NA	D	501	1/1	0.96	0.19	54,54,54,54	0
2	NA	A	502	1/1	0.97	0.06	42,42,42,42	0
3	FLC	A	1449	13/13	0.98	0.10	38,43,52,54	0
6	CL	D	1452	1/1	0.98	0.07	56,56,56,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around BOG C 1448:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BOG D 1454:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BOG A 1450:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around BOG C 1447:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.