Full wwPDB X-ray Structure Validation Report (i)

May 16, 2020 – 01:46 am BST

PDB ID	:	1AGF
Title	:	ANTAGONIST HIV-1 GAG PEPTIDES INDUCE STRUCTURAL CHANGES IN HLA B8-HIV-1 GAG PEPTIDE (GGKKRYKL-5R MUTATION)
Authors	:	Reid, S.W.; Mcadam, S.; Smith, K.J.; Klenerman, P.; O'Callaghan, C.A.; Harlos, K.; Jakobsen, B.K.; Mcmichael, A.J.; Bell, J.; Stuart, D.I.; Jones, E.Y.
Deposited on	:	1997-03-24
Resolution	:	2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	276	86% 13% •
2	B	99	89% 10% •
3	C	8	100%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4805 atoms, of which 1355 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called B*0801.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	276	2787	1395	535	411	439	7	5	0	0

- Molecule 2 is a protein called BETA-2 MICROGLOBULIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	99	1020	528	191	140	158	3	5	0	0

- Molecule 3 is a protein called HIV-1 GAG PEPTIDE (GGKKRYKL - 5R MUTATION).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	C	8	92	43	25	14	10	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
4	A	226	678	452	226	0	0
4	B	67	201	134	67	0	0
4	C	9	27	18	9	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: B*0801

Chain A:



- Molecule 2: BETA-2 MICROGLOBULIN

Chain B:



- Molecule 3: HIV-1 GAG PEPTIDE (GGKKRYKL - 5R MUTATION)

Chain C:



There are no outlier residues recorded for this chain.

4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	50.60Å 81.30Å 110.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.00 – 2.20	Depositor
% Data completeness (in resolution range)	96.3 (14.00-2.20)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.181 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4805	wwPDB-VP
Average B, all atoms (Å^2)	12.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.57	0/2314	0.80	2/3146 (0.1%)
2	B	0.62	1/852 (0.1%)	0.79	1/1152 (0.1%)
3	C	0.60	0/67	0.79	0/83
All	All	0.58	1/3233 (0.0%)	0.80	3/4381 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	80	CYS	CB-SG	-5.21	1.73	1.81

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	28	VAL	N-CA-C	-5.76	95.44	111.00
1	A	272	LEU	CA-CB-CG	5.51	127.96	115.30
2	B	25	CYS	CA-CB-SG	-5.27	104.52	114.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	116	TYR	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2252	535	2095	19	0
2	B	829	191	794	3	1
3	C	67	25	80	0	0
4	A	226	452	0	2	2
4	B	67	134	0	1	0
4	C	9	18	0	0	0
All	All	3450	1355	2969	21	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:GLY:HA3	1:A:110:LEU:HD11	1.23	1.17
1:A:1:GLY:CA	1:A:110:LEU:HD11	2.14	0.68
1:A:1:GLY:HA2	1:A:104:GLY:HA2	1.77	0.65
1:A:49:ALA:O	1:A:52:ILE:HG22	1.99	0.62
1:A:263:HIS:CD2	1:A:265:GLY:H	2.21	0.58
1:A:12:MET:HB3	4:A:344:HOH:O	2.06	0.55
1:A:210:PRO:O	1:A:263:HIS:HE1	1.91	0.54
1:A:131:ARG:HH11	1:A:157:ARG:HH12	1.57	0.52
1:A:263:HIS:HD2	1:A:265:GLY:H	1.58	0.52
1:A:206:LEU:HD23	1:A:242:GLN:HB3	1.93	0.49
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.48	0.49
1:A:219:ARG:O	1:A:219:ARG:HG3	2.12	0.49
1:A:82:ARG:HD2	4:A:482:HOH:O	2.13	0.48
1:A:84:TYR:HB3	1:A:139:ALA:HB1	1.97	0.47
2:B:25:CYS:HB2	2:B:39:LEU:HD21	1.98	0.45
1:A:123:TYR:CZ	1:A:140:ALA:HA	2.51	0.45
1:A:23:ILE:HD12	1:A:36:PHE:O	2.19	0.42
1:A:238:ASP:O	1:A:239:ARG:HB3	2.20	0.41
1:A:110:LEU:O	1:A:111:ARG:HB3	2.21	0.41
2:B:45:ARG:HD2	4:B:144:HOH:O	2.21	0.41
1:A:9:ASP:O	1:A:96:GLN:HA	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:12:ARG:HH11	4:A:485:HOH:H2[1_655]	1.14	0.46
4:A:303:HOH:H2	4:A:491:HOH:O[2_344]	1.53	0.07

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/276 (99%)	267 (97%)	7 (3%)	0	100	100
2	B	97/99 (98%)	96 (99%)	1 (1%)	0	100	100
3	C	6/8 (75%)	6 (100%)	0	0	100	100
All	All	377/383 (98%)	369 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/235 (100%)	222 (94%)	13 (6%)	21	26
2	B	94/94 (100%)	89 (95%)	5 (5%)	22	27
3	C	6/6 (100%)	6 (100%)	0	100	100
All	All	335/335 (100%)	317 (95%)	18 (5%)	22	26

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ARG
1	A	12	MET
1	A	23	ILE
1	A	42	SER
1	A	44	ARG
1	A	58	GLU
1	A	81	LEU
1	A	186	LYS
1	A	197	HIS
1	A	219	ARG
1	A	222	GLU
1	A	239	ARG
1	A	273	ARG
2	B	70	PHE
2	B	83	ASN
2	B	88	SER
2	B	89	GLN
2	B	97	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	115	GLN
1	A	174	ASN
1	A	192	HIS
1	A	263	HIS
2	B	83	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.