Full wwPDB X-ray Structure Validation Report (i)

May 29, 2020 – 04:16 am BST

PDB ID	:	4AO7
Title	:	Zinc bound structure of a novel cold-adapted esterase from an Arctic intertidal metagenomic library
Authors	:	Fu, J.; Leiros, H.-K.S.; Pascale, D.d.; Johnson, K.A.; Blencke, H.M.; Landfald, B.
Deposited on	:	2012-03-23
Resolution	:	1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1890 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ESTERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	223	1764	1116	306	332	10	38	8	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	expression tag	PDB 4AO7
A	-10	ARG	-	expression tag	PDB 4AO7
A	-9	GLY	-	expression tag	PDB 4AO7
A	-8	SER	-	expression tag	PDB 4AO7
A	-7	HIS	-	expression tag	PDB 4AO7
A	-6	HIS	-	expression tag	PDB 4AO7
A	-5	HIS	-	expression tag	PDB 4AO7
A	-4	HIS	-	expression tag	PDB 4AO7
A	-3	HIS	-	expression tag	PDB 4AO7
A	-2	HIS	-	expression tag	PDB 4AO7
A	-1	GLY	-	expression tag	PDB 4AO7
A	0	SER	-	expression tag	PDB 4AO7

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	A	3	3	3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	123	123	123	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ESTERASE

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α, β, γ	55.66Å 70.03Å 53.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.85 24.73 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.5 (25.00-1.85) 98.5 (24.73-1.85)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.11 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.176 , 0.228 0.180 , 0.223	Depositor DCC
R_{free} test set	939 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å^2)	23.5	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 50.4	EDS
L-test for twinning[2]	$< \|L\| > = 0.47$, $< L^2 > = 0.30$	Xtriage
Estimated twinning fraction	0.025 for l,-k,h	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	1890	wwPDB-VP
Average B, all atoms (Å^2)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.93% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.78	13/1804 (0.7%)	1.35	22/2441 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	5

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	12	GLU	CA-CB	38.05	2.37	1.53
1	A	176	GLU	CD-OE2	33.48	1.62	1.25
1	A	14	LYS	CA-C	20.87	2.07	1.52
1	A	61	TYR	CD2-CE2	-19.05	1.10	1.39
1	A	171	GLU	CD-OE2	-18.63	1.05	1.25
1	A	27	ASP	CG-OD1	17.15	1.64	1.25
1	A	111	GLU	CG-CD	-16.56	1.27	1.51
1	A	11	ASP	CB-CG	-15.97	1.18	1.51
1	A	61	TYR	CE2-CZ	14.48	1.57	1.38
1	A	13	ARG	CA-C	-13.77	1.17	1.52
1	A	82	PRO	C-O	-9.42	1.04	1.23
1	A	13	ARG	N-CA	6.31	1.58	1.46
1	A	107	ARG	CB-CG	-6.13	1.36	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	171	GLU	OE1-CD-OE2	-28.50	89.10	123.30
1	A	176	GLU	OE1-CD-OE2	-18.59	100.99	123.30
1	A	40	GLU	CB-CA-C	17.80	145.99	110.40
1	A	12	GLU	CB-CA-C	-17.44	75.52	110.40
1	A	13	ARG	CA-C-O	-16.25	85.97	120.10
1	A	13	ARG	CA-C-N	15.16	150.56	117.20
1	A	13	ARG	CB-CA-C	-14.21	81.98	110.40
1	A	12	GLU	N-CA-CB	-13.05	87.10	110.60
1	A	13	ARG	N-CA-C	10.88	140.37	111.00
1	A	61	TYR	CZ-CE2-CD2	10.28	129.06	119.80
1	A	13	ARG	N-CA-CB	7.62	124.31	110.60
1	A	171	GLU	CG-CD-OE2	7.53	133.36	118.30
1	A	61	TYR	CE1-CZ-CE2	-7.30	108.11	119.80
1	A	27	ASP	CB-CG-OD1	-6.93	112.07	118.30
1	A	11	ASP	CB-CG-OD2	6.54	124.19	118.30
1	A	11	ASP	CA-CB-CG	5.89	126.36	113.40
1	A	176	GLU	CG-CD-OE2	-5.88	106.54	118.30
1	A	14	LYS	CA-C-O	-5.87	107.77	120.10
1	A	107	ARG	CA-CB-CG	5.87	126.31	113.40
1	A	11	ASP	CB-CG-OD1	-5.84	113.05	118.30
1	A	14	LYS	CA-C-N	5.48	129.26	117.20
1	A	61	TYR	OH-CZ-CE2	5.19	134.10	120.10

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	40	GLU	CA

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	12	GLU	Peptide
1	A	13	ARG	Peptide
1	A	171	GLU	Sidechain
1	A	176	GLU	Sidechain
1	A	27	ASP	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1764	0	1740	14	0
2	A	3	0	0	0	0
3	A	123	0	0	3	0
All	All	1890	0	1740	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:MET:N	3:A:2001:HOH:O	2.09	0.84
1:A:247:LYS:NZ	3:A:2122:HOH:O	2.28	0.67
1:A:49:LEU:HB3	1:A:62:ILE:HD12	1.82	0.59
1:A:51:HIS:HB3	1:A:62:ILE:HG13	1.85	0.57
1:A:29:ARG:NH1	1:A:121:ASP:OD1	2.37	0.56
1:A:5:MET:HG2	1:A:24:LEU:HD22	1.89	0.55
1:A:7:TRP:O	1:A:8:ASN:HB2	2.06	0.54
1:A:216[B]:GLN:OE1	3:A:2078:HOH:O	2.18	0.50
1:A:15:LEU:HD12	1:A:16:SER:H	1.79	0.47
1:A:11:ASP:HB2	1:A:19:GLU:HB3	1.98	0.45
1:A:46:LEU:O	1:A:138:THR:HA	2.16	0.44
1:A:158:LYS:NZ	1:A:184:GLN:O	2.50	0.44
1:A:241:TYR:O	1:A:245:ARG:HB2	2.19	0.42
1:A:62:ILE:HD13	1:A:62:ILE:HA	1.91	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	225/259 (87%)	216 (96%)	7 (3%)	2 (1%)	17 6

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	8	ASN
1	A	13	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	185/206 (90%)	184 (100%)	1 (0%)	88 86

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	245	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	223/259 (86%)	0.44	24 (10%) 5 5	12, 24, 54, 102	26 (11%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	14	LYS	6.0
1	A	12	GLU	5.4
1	A	15	LEU	5.4
1	A	30	THR	5.3
1	A	97	THR	5.1
1	A	39	ALA	5.1
1	A	11	ASP	4.8
1	A	25	GLU	4.5
1	A	9	GLY	4.4
1	A	26	VAL	4.3
1	A	16	SER	4.1
1	A	41	GLY	4.1
1	A	27	ASP	3.9
1	A	8	ASN	3.8
1	A	13	ARG	3.6
1	A	10	LYS	3.4
1	A	5	MET	3.4
1	A	42	SER	3.3
1	A	28	GLY	3.1
1	A	104	ALA	3.1
1	A	40	GLU	3.0
1	A	17	VAL	2.7
1	A	103	ASP	2.1
1	A	108	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	ZN	A	1249	1/1	0.96	0.09	34,34,34,34	1
2	ZN	A	1248	1/1	0.98	0.12	26,26,26,26	1
2	ZN	A	1250	1/1	0.99	0.04	33,33,33,33	1

6.5 Other polymers (i)

There are no such residues in this entry.