



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 27, 2024 – 10:47 pm BST

PDB ID : 4AOY
Title : Open CtIDH. The complex structures of Isocitrate dehydrogenase from *Clostridium thermocellum* and *Desulfotalea psychrophila*, support a new active site locking mechanism
Authors : Leiros, H.-K.S.; Fedoy, A.-E.; Leiros, I.; Steen, I.H.
Deposited on : 2012-03-30
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	 3% 61% 26% 6% 6%
1	B	402	 3% 60% 25% 7% 7%
1	C	402	 3% 69% 19% 8% 8%
1	D	402	 3% 60% 27% 6% 7%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 11678 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ISOCITRATE DEHYDROGENASE [NADP].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	376	2917	1854	488	558	17	0	0	0
1	B	372	2859	1817	474	552	16	0	0	0
1	C	371	2884	1837	477	554	16	0	0	0
1	D	375	2930	1869	485	560	16	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	21	Total	O	0	0
			21	21		
2	B	15	Total	O	0	0
			15	15		
2	C	22	Total	O	0	0
			22	22		
2	D	30	Total	O	0	0
			30	30		

- Molecule 1: ISOCITRATE DEHYDROGENASE [NADP]



- Molecule 1: ISOCITRATE DEHYDROGENASE [NADP]



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.13Å 106.84Å 154.50Å 90.00° 93.63° 90.00°	Depositor
Resolution (Å)	20.00 – 2.35 48.02 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-2.35) 99.7 (48.02-2.35)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.210 , 0.258 0.211 , 0.225	Depositor DCC
R_{free} test set	2280 reflections (3.01%)	wwPDB-VP
Wilson B-factor (Å ²)	40.4	Xtrriage
Anisotropy	0.050	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11678	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/2970	0.80	0/4011
1	B	0.64	0/2913	0.81	5/3936 (0.1%)
1	C	0.76	1/2938 (0.0%)	0.81	3/3966 (0.1%)
1	D	0.75	1/2985 (0.0%)	0.80	4/4029 (0.1%)
All	All	0.71	2/11806 (0.0%)	0.80	12/15942 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	267	CYS	CB-SG	-7.07	1.70	1.82
1	D	297	ASP	CB-CG	-5.26	1.40	1.51

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	150	GLY	N-CA-C	7.74	132.45	113.10
1	B	379	VAL	C-N-CA	-7.08	92.28	122.00
1	C	297	ASP	CB-CG-OD1	-6.96	112.03	118.30
1	B	13	MET	CG-SD-CE	-6.59	89.66	100.20
1	B	149	ALA	N-CA-C	6.43	128.37	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	379	VAL	Peptide
1	B	380	PRO	Peptide
1	C	273	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2917	0	2820	144	0
1	B	2859	0	2723	149	0
1	C	2884	0	2788	81	0
1	D	2930	0	2855	132	0
2	A	21	0	0	2	0
2	B	15	0	0	0	0
2	C	22	0	0	3	0
2	D	30	0	0	4	0
All	All	11678	0	11186	492	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 492 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:379:VAL:HG12	1:A:380:PRO:CB	1.38	1.52
1:A:379:VAL:HG12	1:A:380:PRO:CG	1.10	1.52
1:A:379:VAL:CG1	1:A:380:PRO:CG	1.97	1.43
1:A:379:VAL:CG1	1:A:380:PRO:CB	2.00	1.38
1:A:379:VAL:CG1	1:A:380:PRO:HB3	1.54	1.36

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	368/402 (92%)	338 (92%)	24 (6%)	6 (2%)	9 8
1	B	364/402 (90%)	329 (90%)	28 (8%)	7 (2%)	8 6
1	C	363/402 (90%)	342 (94%)	16 (4%)	5 (1%)	11 9
1	D	367/402 (91%)	347 (95%)	17 (5%)	3 (1%)	19 20
All	All	1462/1608 (91%)	1356 (93%)	85 (6%)	21 (1%)	11 9

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	148	SER
1	A	380	PRO
1	B	122	LYS
1	B	150	GLY
1	B	285	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	302/347 (87%)	270 (89%)	32 (11%)	6 6
1	B	291/347 (84%)	254 (87%)	37 (13%)	4 4
1	C	299/347 (86%)	276 (92%)	23 (8%)	13 12
1	D	306/347 (88%)	269 (88%)	37 (12%)	5 4
All	All	1198/1388 (86%)	1069 (89%)	129 (11%)	6 6

5 of 129 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	258	ARG
1	D	285	SER
1	B	219	HIS
1	B	217	TYR
1	D	289	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	358	GLN
1	D	160	ASN
1	C	386	ASN
1	D	182	ASN
1	B	87	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	376/402 (93%)	0.27	12 (3%) 47 59	31, 50, 80, 121	0
1	B	372/402 (92%)	0.32	14 (3%) 40 53	33, 53, 86, 104	0
1	C	371/402 (92%)	0.18	11 (2%) 50 61	26, 42, 75, 97	0
1	D	375/402 (93%)	0.19	11 (2%) 51 62	27, 44, 75, 100	0
All	All	1494/1608 (92%)	0.24	48 (3%) 47 59	26, 47, 81, 121	0

The worst 5 of 48 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	211	ASP	5.3
1	A	212	THR	4.9
1	B	217	TYR	4.5
1	D	84	GLU	3.8
1	D	91	MET	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.