wwPDB X-ray Structure Validation Summary Report (i)

Sep 26, 2023 – 01:44 AM EDT

PDB ID	:	6ATB
Title	:	Crystal Structure of human NAMPT in complex with NVP-LOD812
Authors	:	Weihofen, W.A.; Thigale, S.
Deposited on	:	2017-08-28
Resolution	:	2.53 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5743 (2.54-2.50)
Clashscore	141614	6463 (2.54-2.50)
Ramachandran outliers	138981	6335 (2.54-2.50)
Sidechain outliers	138945	6337 (2.54-2.50)
RSRZ outliers	127900	5630 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	501	83% 9% 7%
1	B	501	87% 7% 6%
1	C	501	2% 80% 12% • 7%
1	D	501	84% 9% • 7%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 16259 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nicotinamide phosphoribosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	466	3729	2400	617	705	7	0	0	0
1	B	470	3762	2421	621	713	7	0	0	0
1	C	466	3729	2400	617	705	7	0	0	0
1	D	468	3746	2411	619	709	7	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	492	LEU	-	expression tag	UNP P43490
A	493	GLU	-	expression tag	UNP P43490
A	494	HIS	-	expression tag	UNP P43490
A	495	HIS	-	expression tag	UNP P43490
A	496	HIS	-	expression tag	UNP P43490
A	497	HIS	-	expression tag	UNP P43490
A	498	HIS	-	expression tag	UNP P43490
A	499	HIS	-	expression tag	UNP P43490
A	500	HIS	-	expression tag	UNP P43490
A	501	HIS	-	expression tag	UNP P43490
B	492	LEU	-	expression tag	UNP P43490
B	493	GLU	-	expression tag	UNP P43490
B	494	HIS	-	expression tag	UNP P43490
B	495	HIS	-	expression tag	UNP P43490
B	496	HIS	-	expression tag	UNP P43490
B	497	HIS	-	expression tag	UNP P43490
B	498	HIS	-	expression tag	UNP P43490
B	499	HIS	-	expression tag	UNP P43490
B	500	HIS	-	expression tag	UNP P43490
B	501	HIS	-	expression tag	UNP P43490
C	492	LEU	-	expression tag	UNP P43490

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	493	GLU	-	expression tag	UNP P43490
C	494	HIS	-	expression tag	UNP P43490
C	495	HIS	-	expression tag	UNP P43490
C	496	HIS	-	expression tag	UNP P43490
C	497	HIS	-	expression tag	UNP P43490
C	498	HIS	-	expression tag	UNP P43490
C	499	HIS	-	expression tag	UNP P43490
C	500	HIS	-	expression tag	UNP P43490
C	501	HIS	-	expression tag	UNP P43490
D	492	LEU	-	expression tag	UNP P43490
D	493	GLU	-	expression tag	UNP P43490
D	494	HIS	-	expression tag	UNP P43490
D	495	HIS	-	expression tag	UNP P43490
D	496	HIS	-	expression tag	UNP P43490
D	497	HIS	-	expression tag	UNP P43490
D	498	HIS	-	expression tag	UNP P43490
D	499	HIS	-	expression tag	UNP P43490
D	500	HIS	-	expression tag	UNP P43490
D	501	HIS	-	expression tag	UNP P43490

- Molecule 2 is N-{4-[(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)methyl]phenyl}-N'-[(pyridin-3-yl)methyl]urea (three-letter code: BWA) (formula: $C_{22}H_{18}N_4O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	29	22	4	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	B	1	29	22	4	3	0	0
			Total	C	N	O		
2	C	1	29	22	4	3	0	0
			Total	C	N	O		
2	D	1	29	22	4	3	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O			
3	A	1	6	3	3		0	0
			Total	C	O			
3	B	1	6	3	3		0	0
			Total	C	O			
3	C	1	6	3	3		0	0
			Total	C	O			
3	C	1	6	3	3		0	0
			Total	C	O			
3	D	1	6	3	3		0	0

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0
4	D	1	Total O P 5 4 1	0	0

- Molecule 5 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	S		
5	C	1	4	2	1	1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	301	301	301	0	0
			Total	O		
6	B	299	299	299	0	0
			Total	O		
6	C	221	221	221	0	0
			Total	O		
6	D	302	302	302	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nicotinamide phosphoribosyltransferase

Chain A:



• Molecule 1: Nicotinamide phosphoribosyltransferase

Chain B:



• Molecule 1: Nicotinamide phosphoribosyltransferase

Chain C:



• Molecule 1: Nicotinamide phosphoribosyltransferase

Chain D:

84%
9%
7%


MET ASN PRO ALA ALA GLU ALA GLU F9 L12 Y18 Q25 K32 K42 LYS THR GLU ASN SER LYS LEU ARG LYS VAL K53 Y54 E55 E82 K88 F96 N102 K107 P113 P120 F123 T137 S165 Q168 L172 Y175 Y195
R196 S209 D219 I227 K255 K259 Y281 D294 P312 L318 V321 E326 I351 Q352 I358 L361 I364 I375 I378 K400 V404 V405 L409 H433 R434 L443 E452 N464 L486 GLU ALA ALA HIS HIS LEU
GLU HIS HIS HIS HIS HIS HIS HIS HIS

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	98.46Å 208.40Å 98.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	104.20 – 2.53 104.20 – 2.43	Depositor EDS
% Data completeness (in resolution range)	100.0 (104.20-2.53) 100.0 (104.20-2.43)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.91 (at 2.42Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.152 , 0.215 0.151 , 0.211	Depositor DCC
R_{free} test set	3897 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å^2)	32.9	Xtriage
Anisotropy	0.510	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 65.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.023 for l,-k,h	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	16259	wwPDB-VP
Average B, all atoms (Å^2)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.69% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, DMS, BWA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.50	0/3817	0.70	0/5171
1	B	0.50	0/3850	0.69	0/5216
1	C	0.51	0/3817	0.72	0/5171
1	D	0.52	0/3834	0.70	0/5194
All	All	0.51	0/15318	0.70	0/20752

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3729	0	3712	21	0
1	B	3762	0	3744	13	0
1	C	3729	0	3712	34	0
1	D	3746	0	3729	23	0
2	A	29	0	0	0	0
2	B	29	0	0	0	0
2	C	29	0	0	0	0
2	D	29	0	0	0	0
3	A	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	6	0	8	0	0
3	C	12	0	16	1	0
3	D	6	0	8	1	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
4	C	5	0	0	0	0
4	D	5	0	0	0	0
5	C	4	0	6	0	0
6	A	301	0	0	1	0
6	B	299	0	0	2	0
6	C	221	0	0	0	0
6	D	302	0	0	1	0
All	All	16259	0	14943	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 84 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:153:VAL:CG1	1:D:196:ARG:HB2	2.19	0.72
1:A:299:ILE:HD12	1:A:308:LEU:HD22	1.70	0.71
1:A:245:ALA:HA	1:B:25:GLN:HE22	1.56	0.71
1:C:316:ASN:HD22	1:C:319:ASP:H	1.39	0.69
1:C:153:VAL:HG11	1:D:196:ARG:HB2	1.75	0.68

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	462/501 (92%)	446 (96%)	16 (4%)	0	100	100
1	B	466/501 (93%)	451 (97%)	15 (3%)	0	100	100
1	C	462/501 (92%)	442 (96%)	19 (4%)	1 (0%)	47	67
1	D	464/501 (93%)	451 (97%)	13 (3%)	0	100	100
All	All	1854/2004 (92%)	1790 (96%)	63 (3%)	1 (0%)	51	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	389	LYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/440 (93%)	390 (95%)	20 (5%)	25	45
1	B	414/440 (94%)	398 (96%)	16 (4%)	32	55
1	C	410/440 (93%)	387 (94%)	23 (6%)	21	38
1	D	412/440 (94%)	397 (96%)	15 (4%)	35	59
All	All	1646/1760 (94%)	1572 (96%)	74 (4%)	27	48

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	469	LYS
1	D	434	ARG
1	D	18	TYR
1	D	195	TYR
1	B	119	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	25	GLN
1	D	164	ASN
1	D	396	ASN
1	B	362	GLN
1	B	164	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	GOL	A	602	-	5,5,5	0.06	0	5,5,5	0.35	0
2	BWA	D	602	-	32,32,32	1.30	2 (6%)	44,44,44	1.90	11 (25%)
3	GOL	C	604	-	5,5,5	0.09	0	5,5,5	0.58	0
4	PO4	D	601	-	4,4,4	1.25	1 (25%)	6,6,6	0.36	0
3	GOL	D	603	-	5,5,5	0.11	0	5,5,5	0.07	0
2	BWA	C	602	-	32,32,32	1.27	2 (6%)	44,44,44	1.79	11 (25%)
4	PO4	C	605	-	4,4,4	1.38	1 (25%)	6,6,6	0.53	0
3	GOL	C	603	-	5,5,5	0.04	0	5,5,5	0.17	0
2	BWA	B	601	-	32,32,32	1.30	3 (9%)	44,44,44	1.96	12 (27%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	PO4	A	603	-	4,4,4	1.32	1 (25%)	6,6,6	0.43	0
3	GOL	B	602	-	5,5,5	0.02	0	5,5,5	0.24	0
5	DMS	C	601	-	3,3,3	0.29	0	3,3,3	0.14	0
2	BWA	A	601	-	32,32,32	1.12	2 (6%)	44,44,44	1.89	11 (25%)
4	PO4	B	603	-	4,4,4	1.14	0	6,6,6	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	602	-	-	0/4/4/4	-
2	BWA	D	602	-	-	0/13/29/29	0/4/4/4
3	GOL	C	604	-	-	0/4/4/4	-
3	GOL	D	603	-	-	0/4/4/4	-
2	BWA	C	602	-	-	0/13/29/29	0/4/4/4
3	GOL	C	603	-	-	0/4/4/4	-
2	BWA	B	601	-	-	0/13/29/29	0/4/4/4
3	GOL	B	602	-	-	0/4/4/4	-
2	BWA	A	601	-	-	0/13/29/29	0/4/4/4

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	602	BWA	C20-N19	-3.50	1.35	1.39
2	C	602	BWA	C20-N19	-3.20	1.35	1.39
2	B	601	BWA	C23-N19	-3.02	1.36	1.39
2	B	601	BWA	C20-N19	-2.93	1.36	1.39
2	C	602	BWA	C23-N19	-2.92	1.36	1.39

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	601	BWA	C21-C20-N19	5.07	109.48	105.88
2	D	602	BWA	C22-C23-N19	4.92	109.38	105.88
2	B	601	BWA	C22-C23-N19	4.85	109.32	105.88
2	A	601	BWA	C22-C23-N19	4.60	109.15	105.88
2	C	602	BWA	C22-C23-N19	4.57	109.13	105.88

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	603	GOL	1	0
3	C	603	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand BWA D 602



Bond lengths



Bond angles



Torsions



Rings

Ligand BWA C 602



Bond lengths



Bond angles



Torsions



Rings

Ligand BWA B 601



Bond lengths



Bond angles



Torsions



Rings

Ligand BWA A 601



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	466/501 (93%)	-0.57	0 100 100	21, 35, 55, 84	0
1	B	470/501 (93%)	-0.65	0 100 100	20, 34, 55, 83	0
1	C	466/501 (93%)	-0.24	10 (2%) 63 67	20, 40, 78, 102	0
1	D	468/501 (93%)	-0.63	2 (0%) 92 93	18, 32, 64, 91	0
All	All	1870/2004 (93%)	-0.52	12 (0%) 89 90	18, 35, 69, 102	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	179	THR	4.7
1	C	367	GLY	3.5
1	C	319	ASP	3.2
1	C	324	VAL	3.2
1	C	368	MET	3.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	PO4	D	601	5/5	0.84	0.18	126,126,127,127	0
5	DMS	C	601	4/4	0.84	0.18	103,103,104,104	0
3	GOL	C	603	6/6	0.86	0.29	62,67,68,69	0
3	GOL	C	604	6/6	0.90	0.23	63,66,66,67	0
3	GOL	B	602	6/6	0.91	0.21	53,54,57,59	0
4	PO4	B	603	5/5	0.91	0.22	73,73,75,75	0
3	GOL	A	602	6/6	0.92	0.21	52,54,60,63	0
4	PO4	A	603	5/5	0.94	0.26	71,73,75,76	0
2	BWA	C	602	29/29	0.94	0.23	40,51,53,56	0
3	GOL	D	603	6/6	0.95	0.12	49,55,58,62	0
2	BWA	A	601	29/29	0.96	0.18	24,35,45,47	0
4	PO4	C	605	5/5	0.97	0.21	66,67,68,68	0
2	BWA	D	602	29/29	0.97	0.13	18,30,39,41	0
2	BWA	B	601	29/29	0.97	0.15	24,28,51,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BWA C 602:

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around BWA A 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BWA D 602:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around BWA B 601:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.