wwPDB X-ray Structure Validation Summary Report (i)

Jan 27, 2024 – 01:22 PM EST

PDB ID	:	1AVO
Title	:	PROTEASOME ACTIVATOR REG(ALPHA)
Authors	:	Hill, C.P.; Knowlton, J.R.
Deposited on	:	1997-09-18
Resolution	:	2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	60	62% 37% •
1	C	60	67% 32% •
1	E	60	58% 40% •
1	G	60	62% 37% •
1	I	60	67% 32% •
1	K	60	62% 37% •
1	M	60	65% 33% •
2	B	140	71% 28% •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	140	71% 26% •
2	F	140	69% 29% •
2	H	140	74% 25% •
2	J	140	71% 27% •
2	L	140	78% 21% •
2	N	140	71% 26% •

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 11361 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 11S REGULATOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	60	475	303	78	93	1	0	0	0
1	C	60	475	303	78	93	1	0	0	0
1	E	60	475	303	78	93	1	0	0	0
1	G	60	475	303	78	93	1	0	0	0
1	I	60	475	303	78	93	1	0	0	0
1	K	60	475	303	78	93	1	0	0	0
1	M	60	475	303	78	93	1	0	0	0

- Molecule 2 is a protein called 11S REGULATOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	140	1148	734	200	210	4	0	0	0
2	D	140	1148	734	200	210	4	0	0	0
2	F	140	1148	734	200	210	4	0	0	0
2	H	140	1148	734	200	210	4	0	0	0
2	J	140	1148	734	200	210	4	0	0	0
2	L	140	1148	734	200	210	4	0	0	0
2	N	140	1148	734	200	210	4	0	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: 11S REGULATOR

Chain A:



• Molecule 1: 11S REGULATOR

Chain C:



• Molecule 1: 11S REGULATOR

Chain E:



• Molecule 1: 11S REGULATOR

Chain G:



• Molecule 1: 11S REGULATOR

Chain I:



• Molecule 1: 11S REGULATOR

Chain K:

• Molecule 1: 11S REGULATOR

Chain M:



• Molecule 2: 11S REGULATOR

Chain B:



• Molecule 2: 11S REGULATOR

Chain D:



• Molecule 2: 11S REGULATOR

Chain F:



• Molecule 2: 11S REGULATOR

Chain H:



- Molecule 2: 11S REGULATOR

Chain J:



- Molecule 2: 11S REGULATOR

Chain L:



- Molecule 2: 11S REGULATOR

Chain N:



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	162.60Å 134.30Å 116.20Å 90.00° 90.60° 90.00°	Depositor
Resolution (Å)	10.00 – 2.80	Depositor
% Data completeness (in resolution range)	98.4 (10.00-2.80)	Depositor
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.249 , 0.289	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	11361	wwPDB-VP
Average B, all atoms (Å^2)	28.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.89	0/482	0.82	0/650
1	C	0.90	0/482	0.87	0/650
1	E	0.87	0/482	0.84	0/650
1	G	0.86	0/482	0.83	0/650
1	I	0.81	0/482	0.82	0/650
1	K	0.83	0/482	0.79	0/650
1	M	0.87	0/482	0.84	0/650
2	B	0.99	0/1167	0.82	0/1572
2	D	1.02	1/1167 (0.1%)	0.84	0/1572
2	F	0.96	0/1167	0.82	0/1572
2	H	0.94	0/1167	0.82	0/1572
2	J	0.99	1/1167 (0.1%)	0.83	0/1572
2	L	0.94	0/1167	0.85	1/1572 (0.1%)
2	N	1.03	1/1167 (0.1%)	0.86	1/1572 (0.1%)
All	All	0.95	3/11543 (0.0%)	0.83	2/15554 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	N	157	PHE	CB-CG	-5.40	1.42	1.51
2	J	154	GLU	CG-CD	5.20	1.59	1.51
2	D	106	CYS	CB-SG	-5.11	1.73	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	N	210	ARG	NE-CZ-NH1	7.11	123.86	120.30
2	L	210	ARG	NE-CZ-NH1	6.25	123.42	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	475	0	485	15	0
1	C	475	0	485	15	0
1	E	475	0	485	20	0
1	G	475	0	485	15	0
1	I	475	0	485	14	0
1	K	475	0	485	16	0
1	M	475	0	485	15	0
2	B	1148	0	1177	27	0
2	D	1148	0	1177	29	0
2	F	1148	0	1177	36	0
2	H	1148	0	1177	30	0
2	J	1148	0	1177	32	0
2	L	1148	0	1177	21	0
2	N	1148	0	1177	26	0
All	All	11361	0	11634	266	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 266 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:63:ILE:HB	2:J:198:ARG:NH2	1.73	1.03
1:E:63:ILE:HB	2:F:198:ARG:NH1	1.86	0.90
1:K:63:ILE:HB	2:L:198:ARG:NH2	1.91	0.86
1:M:63:ILE:HB	2:N:198:ARG:NH2	1.91	0.85
2:B:122:LYS:HD2	2:N:210:ARG:HH11	1.46	0.80

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	58/60 (97%)	56 (97%)	2 (3%)	0	100	100
1	C	58/60 (97%)	53 (91%)	5 (9%)	0	100	100
1	E	58/60 (97%)	55 (95%)	3 (5%)	0	100	100
1	G	58/60 (97%)	56 (97%)	2 (3%)	0	100	100
1	I	58/60 (97%)	54 (93%)	4 (7%)	0	100	100
1	K	58/60 (97%)	55 (95%)	3 (5%)	0	100	100
1	M	58/60 (97%)	55 (95%)	3 (5%)	0	100	100
2	B	138/140 (99%)	136 (99%)	2 (1%)	0	100	100
2	D	138/140 (99%)	137 (99%)	1 (1%)	0	100	100
2	F	138/140 (99%)	136 (99%)	2 (1%)	0	100	100
2	H	138/140 (99%)	136 (99%)	2 (1%)	0	100	100
2	J	138/140 (99%)	136 (99%)	2 (1%)	0	100	100
2	L	138/140 (99%)	137 (99%)	1 (1%)	0	100	100
2	N	138/140 (99%)	137 (99%)	1 (1%)	0	100	100
All	All	1372/1400 (98%)	1339 (98%)	33 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	53/53 (100%)	50 (94%)	3 (6%)	20	50
1	C	53/53 (100%)	51 (96%)	2 (4%)	33	67
1	E	53/53 (100%)	50 (94%)	3 (6%)	20	50
1	G	53/53 (100%)	51 (96%)	2 (4%)	33	67
1	I	53/53 (100%)	50 (94%)	3 (6%)	20	50
1	K	53/53 (100%)	51 (96%)	2 (4%)	33	67
1	M	53/53 (100%)	51 (96%)	2 (4%)	33	67
2	B	126/126 (100%)	122 (97%)	4 (3%)	39	73
2	D	126/126 (100%)	121 (96%)	5 (4%)	31	65
2	F	126/126 (100%)	122 (97%)	4 (3%)	39	73
2	H	126/126 (100%)	122 (97%)	4 (3%)	39	73
2	J	126/126 (100%)	122 (97%)	4 (3%)	39	73
2	L	126/126 (100%)	121 (96%)	5 (4%)	31	65
2	N	126/126 (100%)	121 (96%)	5 (4%)	31	65
All	All	1253/1253 (100%)	1205 (96%)	48 (4%)	33	67

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	28	LEU
1	K	31	SER
1	I	31	SER
2	J	159	LEU
2	L	138	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
2	J	193	HIS
2	L	193	HIS
2	N	202	HIS
1	M	56	ASN
2	L	115	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.