



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 25, 2023 – 06:26 PM EDT

PDB ID : 6AV7
Title : Structure of human endothelial nitric oxide synthase heme domain in complex with HW69
Authors : Li, H.; Poulos, T.L.
Deposited on : 2017-09-01
Resolution : 1.92 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	 29% 74% 17% 8%
1	B	440	 10% 81% 10% 9%
1	C	440	 20% 80% 10% 9%
1	D	440	 10% 80% 11% 9%

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		
3	C	1	Total	C	N	O	0	0
			17	9	5	3		
3	D	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 6-(2-{3-[3-(dimethylamino)propyl]-5-fluorophenyl}ethyl)-4-methylpyridin-2-amine (three-letter code: W69) (formula: $C_{19}H_{26}FN_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	F	N		
4	A	1	23	19	1	3	0	0
4	B	1	23	19	1	3	0	0
4	C	1	23	19	1	3	0	0
4	D	1	23	19	1	3	0	0

- Molecule 5 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: C₈H₁₉NO₅).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		
6	C	1	Total	Zn	0	0
			1	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	C O	0	0
			6	3 3		
7	C	1	Total	C O	0	0
			6	3 3		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Cl	0	0
			1	1		
8	B	1	Total	Cl	0	0
			1	1		
8	C	1	Total	Cl	0	0
			1	1		
8	D	1	Total	Cl	0	0
			1	1		

- Molecule 9 is GADOLINIUM ATOM (three-letter code: GD) (formula: Gd).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	Gd	0	0
			1	1		
9	B	1	Total	Gd	0	0
			1	1		
9	C	1	Total	Gd	0	0
			1	1		
9	D	1	Total	Gd	0	0
			1	1		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	34	Total	O	0	0
			34	34		
10	B	117	Total	O	0	0
			117	117		
10	C	73	Total	O	0	0
			73	73		
10	D	130	Total	O	0	0
			130	130		



• Molecule 1: Nitric oxide synthase, endothelial



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.45Å 152.35Å 108.66Å 90.00° 90.79° 90.00°	Depositor
Resolution (Å)	39.03 – 1.92 39.03 – 1.92	Depositor EDS
% Data completeness (in resolution range)	94.5 (39.03-1.92) 95.0 (39.03-1.92)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 1.92Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.204 , 0.254 0.206 , 0.251	Depositor DCC
R_{free} test set	7085 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	39.5	Xtrriage
Anisotropy	0.482	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 55.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.105 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13722	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BTB, HEM, GD, GOL, W69, CL, H4B

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/3335	0.49	0/4543
1	B	0.39	0/3319	0.52	0/4523
1	C	0.35	0/3307	0.49	0/4507
1	D	0.43	0/3319	0.54	0/4523
All	All	0.38	0/13280	0.51	0/18096

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	89	GLN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3237	0	3146	56	0
1	B	3221	0	3126	27	0
1	C	3209	0	3109	27	0
1	D	3221	0	3126	28	0
2	A	43	0	30	5	0
2	B	43	0	30	4	0
2	C	43	0	30	5	0
2	D	43	0	30	3	0
3	A	17	0	15	2	0
3	B	17	0	15	1	0
3	C	17	0	15	1	0
3	D	17	0	15	0	0
4	A	23	0	0	2	0
4	B	23	0	0	3	0
4	C	23	0	0	3	0
4	D	23	0	0	2	0
5	A	28	0	38	6	0
5	B	28	0	36	5	0
5	C	42	0	56	6	0
5	D	28	0	36	5	0
6	A	1	0	0	0	0
6	C	1	0	0	0	0
7	A	6	0	8	1	0
7	C	6	0	8	1	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
9	A	1	0	0	0	0
9	B	1	0	0	0	0
9	C	1	0	0	0	0
9	D	1	0	0	0	0
10	A	34	0	0	0	0
10	B	117	0	0	1	0
10	C	73	0	0	3	0
10	D	130	0	0	1	0
All	All	13722	0	12869	161	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 161 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:235:CYS:H	1:C:238:ARG:HD3	1.36	0.88
1:D:275:ILE:HD12	1:D:281:PRO:HG3	1.63	0.80
1:A:242:ARG:NH2	1:A:477:PRO:O	2.15	0.79
1:A:125:SER:HA	1:A:128:ARG:HE	1.50	0.76
1:A:382:CYS:HA	5:A:504:BTB:H11	1.70	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/440 (91%)	372 (92%)	26 (6%)	4 (1%)	15	6
1	B	401/440 (91%)	394 (98%)	7 (2%)	0	100	100
1	C	399/440 (91%)	378 (95%)	21 (5%)	0	100	100
1	D	401/440 (91%)	395 (98%)	6 (2%)	0	100	100
All	All	1603/1760 (91%)	1539 (96%)	60 (4%)	4 (0%)	41	38

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	90	GLN
1	A	283	ASN
1	A	238	ARG
1	A	281	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/373 (92%)	334 (97%)	11 (3%)	39	29
1	B	344/373 (92%)	334 (97%)	10 (3%)	42	33
1	C	342/373 (92%)	334 (98%)	8 (2%)	50	43
1	D	344/373 (92%)	340 (99%)	4 (1%)	71	69
All	All	1375/1492 (92%)	1342 (98%)	33 (2%)	50	41

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	302	LEU
1	D	71	VAL
1	D	258	ASP
1	B	90	GLN
1	B	67	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 33 ligands modelled in this entry, 10 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BTB	C	505	-	13,13,13	0.65	0	7,16,16	1.02	1 (14%)
2	HEM	C	501	1	41,50,50	2.02	8 (19%)	45,82,82	1.82	9 (20%)
2	HEM	D	501	1	41,50,50	1.98	7 (17%)	45,82,82	1.86	12 (26%)
5	BTB	D	505	-	13,13,13	0.48	0	7,16,16	0.74	0
4	W69	B	503	-	24,24,24	0.60	0	32,32,32	1.66	6 (18%)
3	H4B	D	502	-	16,18,18	0.82	0	11,26,26	2.65	6 (54%)
3	H4B	B	502	-	16,18,18	0.94	0	11,26,26	2.77	5 (45%)
3	H4B	A	502	-	16,18,18	0.98	0	11,26,26	2.69	6 (54%)
5	BTB	A	505	-	13,13,13	0.44	0	7,16,16	0.61	0
7	GOL	C	508	-	5,5,5	0.40	0	5,5,5	0.43	0
7	GOL	A	507	-	5,5,5	0.36	0	5,5,5	0.39	0
3	H4B	C	502	-	16,18,18	0.89	0	11,26,26	2.67	6 (54%)
5	BTB	A	504	9	13,13,13	0.38	0	7,16,16	0.46	0
4	W69	A	503	-	24,24,24	0.46	0	32,32,32	1.42	4 (12%)
5	BTB	B	505	-	13,13,13	0.40	0	7,16,16	0.78	0
4	W69	C	503	-	24,24,24	0.45	0	32,32,32	1.61	4 (12%)
5	BTB	D	504	9	13,13,13	0.39	0	7,16,16	0.60	0
2	HEM	A	501	1	41,50,50	1.97	6 (14%)	45,82,82	1.84	10 (22%)
2	HEM	B	501	1	41,50,50	1.95	8 (19%)	45,82,82	1.85	10 (22%)
5	BTB	C	504	9	13,13,13	0.36	0	7,16,16	0.61	0
5	BTB	B	504	9	13,13,13	0.50	0	7,16,16	0.41	0
5	BTB	C	506	-	13,13,13	0.37	0	7,16,16	0.36	0
4	W69	D	503	-	24,24,24	0.60	0	32,32,32	1.52	6 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BTB	C	505	-	-	5/21/21/21	-
2	HEM	C	501	1	-	9/12/54/54	-
2	HEM	D	501	1	-	4/12/54/54	-
5	BTB	D	505	-	-	8/21/21/21	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	W69	B	503	-	-	3/11/11/11	0/2/2/2
3	H4B	D	502	-	-	3/8/17/17	0/2/2/2
3	H4B	B	502	-	-	0/8/17/17	0/2/2/2
3	H4B	A	502	-	-	3/8/17/17	0/2/2/2
5	BTB	A	505	-	-	7/21/21/21	-
7	GOL	C	508	-	-	4/4/4/4	-
7	GOL	A	507	-	-	4/4/4/4	-
3	H4B	C	502	-	-	0/8/17/17	0/2/2/2
5	BTB	A	504	9	-	9/21/21/21	-
4	W69	A	503	-	-	3/11/11/11	0/2/2/2
5	BTB	B	505	-	-	2/21/21/21	-
4	W69	C	503	-	-	5/11/11/11	0/2/2/2
5	BTB	D	504	9	-	5/21/21/21	-
2	HEM	A	501	1	-	7/12/54/54	-
2	HEM	B	501	1	-	7/12/54/54	-
5	BTB	C	504	9	-	1/21/21/21	-
5	BTB	B	504	9	-	1/21/21/21	-
5	BTB	C	506	-	-	8/21/21/21	-
4	W69	D	503	-	-	6/11/11/11	0/2/2/2

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	HEM	C3D-C2D	8.02	1.53	1.36
2	C	501	HEM	C3D-C2D	8.00	1.53	1.36
2	D	501	HEM	C3D-C2D	7.70	1.53	1.36
2	B	501	HEM	C3D-C2D	7.60	1.52	1.36
2	B	501	HEM	C3C-CAC	4.00	1.56	1.47

The worst 5 of 85 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	HEM	C4D-ND-C1D	6.15	111.42	105.07
2	B	501	HEM	C4D-ND-C1D	6.04	111.31	105.07
2	C	501	HEM	C4D-ND-C1D	5.87	111.14	105.07
2	D	501	HEM	C4D-ND-C1D	5.86	111.13	105.07
4	C	503	W69	C02-N01-C06	5.85	122.54	118.10

There are no chirality outliers.

5 of 104 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	HEM	C1A-C2A-CAA-CBA
2	A	501	HEM	C3A-C2A-CAA-CBA
2	B	501	HEM	C1A-C2A-CAA-CBA
2	B	501	HEM	C3A-C2A-CAA-CBA
2	C	501	HEM	C1A-C2A-CAA-CBA


There are no ring outliers.


22 monomers are involved in 48 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	505	BTB	3	0
2	C	501	HEM	5	0
2	D	501	HEM	3	0
5	D	505	BTB	2	0
4	B	503	W69	3	0
3	B	502	H4B	1	0
3	A	502	H4B	2	0
5	A	505	BTB	2	0
7	C	508	GOL	1	0
7	A	507	GOL	1	0
3	C	502	H4B	1	0
5	A	504	BTB	4	0
4	A	503	W69	2	0
5	B	505	BTB	4	0
4	C	503	W69	3	0
5	D	504	BTB	3	0
2	A	501	HEM	5	0
2	B	501	HEM	4	0
5	C	504	BTB	2	0
5	B	504	BTB	1	0
5	C	506	BTB	1	0
4	D	503	W69	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and


any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/440 (91%)	1.66	129 (31%) 0 0	37, 75, 139, 169	0
1	B	402/440 (91%)	0.66	43 (10%) 6 7	32, 51, 93, 141	0
1	C	401/440 (91%)	1.17	90 (22%) 0 0	38, 66, 112, 156	0
1	D	402/440 (91%)	0.53	42 (10%) 6 7	31, 51, 85, 138	0
All	All	1609/1760 (91%)	1.00	304 (18%) 1 1	31, 59, 117, 169	0

The worst 5 of 304 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	109	LEU	9.6
1	A	153	VAL	8.3
1	A	107	ARG	7.9
1	A	244	TRP	7.8
1	A	480	TRP	7.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	H4B	C	502	17/17	0.74	0.32	73,86,103,104	0
3	H4B	A	502	17/17	0.75	0.35	74,86,103,103	0
5	BTB	C	506	14/14	0.78	0.13	88,97,103,109	0
3	H4B	D	502	17/17	0.82	0.27	59,76,91,91	0
5	BTB	B	504	14/14	0.82	0.17	44,68,90,98	0
3	H4B	B	502	17/17	0.82	0.25	51,78,100,102	0
5	BTB	D	504	14/14	0.82	0.17	54,74,87,88	0
4	W69	C	503	23/23	0.84	0.41	47,91,126,128	0
5	BTB	C	505	14/14	0.85	0.18	59,76,90,99	0
4	W69	A	503	23/23	0.88	0.45	56,104,133,134	0
5	BTB	D	505	14/14	0.89	0.20	70,85,93,100	0
5	BTB	B	505	14/14	0.91	0.23	42,74,89,95	0
9	GD	A	509	1/1	0.91	0.05	131,131,131,131	0
4	W69	D	503	23/23	0.92	0.23	30,78,111,112	0
5	BTB	A	505	14/14	0.92	0.12	64,80,89,91	0
7	GOL	A	507	6/6	0.92	0.23	59,83,102,111	0
4	W69	B	503	23/23	0.92	0.22	33,71,101,101	0
7	GOL	C	508	6/6	0.93	0.21	50,77,81,86	0
5	BTB	C	504	14/14	0.93	0.25	35,85,109,114	0
9	GD	C	510	1/1	0.93	0.05	106,106,106,106	0
2	HEM	A	501	43/43	0.94	0.25	53,70,98,110	0
2	HEM	C	501	43/43	0.95	0.21	37,55,86,110	0
8	CL	A	508	1/1	0.95	0.22	63,63,63,63	0
8	CL	C	509	1/1	0.95	0.20	56,56,56,56	0
2	HEM	D	501	43/43	0.95	0.14	30,41,85,102	0
5	BTB	A	504	14/14	0.95	0.28	52,91,102,115	0
2	HEM	B	501	43/43	0.96	0.15	33,41,94,97	0
8	CL	D	507	1/1	0.97	0.10	45,45,45,45	0
9	GD	B	507	1/1	0.98	0.14	54,54,54,54	0
6	ZN	A	506	1/1	0.98	0.08	53,53,53,53	0
9	GD	D	506	1/1	0.98	0.12	54,54,54,54	0
8	CL	B	506	1/1	0.99	0.14	44,44,44,44	0
6	ZN	C	507	1/1	1.00	0.10	46,46,46,46	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around W69 C 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around W69 A 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around W69 D 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around W69 B 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM C 501:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.