Full wwPDB X-ray Structure Validation Report (i)

Oct 23, 2023 – 03:35 PM EDT

PDB ID	:	3B8S
Title	:	Crystal structure of wild-type chitinase A from Vibrio harveyi
Authors	:	Songsiriritthigul, C.; Pantoom, S.; Aguda, A.H.; Robinson, R.C.; Suginta, W.
Deposited on	:	2007-11-01
Resolution	:	2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	584	
1	B	584	

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 9799 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chitinase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	567	4354	2760	706	863	25	0	0	0
1	B	567	4354	2760	706	863	25	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	598	ARG	-	expression tag	UNP Q9AMP1
A	599	SER	-	expression tag	UNP Q9AMP1
A	600	HIS	-	expression tag	UNP Q9AMP1
A	601	HIS	-	expression tag	UNP Q9AMP1
A	602	HIS	-	expression tag	UNP Q9AMP1
A	603	HIS	-	expression tag	UNP Q9AMP1
A	604	HIS	-	expression tag	UNP Q9AMP1
A	605	HIS	-	expression tag	UNP Q9AMP1
B	598	ARG	-	expression tag	UNP Q9AMP1
B	599	SER	-	expression tag	UNP Q9AMP1
B	600	HIS	-	expression tag	UNP Q9AMP1
B	601	HIS	-	expression tag	UNP Q9AMP1
B	602	HIS	-	expression tag	UNP Q9AMP1
B	603	HIS	-	expression tag	UNP Q9AMP1
B	604	HIS	-	expression tag	UNP Q9AMP1
B	605	HIS	-	expression tag	UNP Q9AMP1

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	562	562	562	0	0
2	B	529	529	529	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chitinase A

Chain A:





- Molecule 1: Chitinase A

Chain B:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	60.27Å 64.28Å 83.52Å 91.74° 91.18° 112.91°	Depositor
Resolution (Å)	24.62 – 2.00 24.62 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.6 (24.62-2.00) 94.6 (24.62-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.08	Depositor
$< I/\sigma(I) >$ [1]	4.32 (at 1.99Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.168 , 0.205 0.170 , 0.207	Depositor DCC
R_{free} test set	3724 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å^2)	15.8	Xtriage
Anisotropy	0.511	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.37 , 48.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.033 for -h,-k,l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	9799	wwPDB-VP
Average B, all atoms (Å^2)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.36	0/4467	0.50	0/6075
1	B	0.37	0/4467	0.50	0/6075
All	All	0.36	0/8934	0.50	0/12150

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4354	0	4103	19	0
1	B	4354	0	4103	29	0
2	A	562	0	0	1	1
2	B	529	0	0	4	0
All	All	9799	0	8206	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:53:ASP:HB2	2:B:806:HOH:O	1.32	1.24

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22:ALA:N	1:B:119:THR:HG1	1.76	0.83
1:B:313:ASP:HB3	2:B:702:HOH:O	1.86	0.74
1:A:22:ALA:N	1:A:119:THR:HG1	1.90	0.69
1:B:344:ARG:O	1:B:344:ARG:HD3	1.96	0.65
1:B:238:PHE:H	1:B:243:HIS:CD2	2.16	0.62
1:B:238:PHE:H	1:B:243:HIS:HD2	1.46	0.61
1:A:231:TRP:HA	1:A:235:GLN:HB2	1.83	0.60
1:A:344:ARG:O	1:A:344:ARG:HD3	2.04	0.58
1:A:231:TRP:HA	1:A:235:GLN:HE21	1.68	0.58
1:B:271:SER:OG	1:B:313:ASP:OD2	2.23	0.57
1:A:238:PHE:H	1:A:243:HIS:HD2	1.53	0.56
1:B:231:TRP:HA	1:B:235:GLN:HB2	1.86	0.55
1:B:270:PRO:HD3	1:B:307:TYR:CD2	2.42	0.54
1:A:238:PHE:H	1:A:243:HIS:CD2	2.25	0.54
1:A:380:VAL:HG22	1:A:453:LYS:HD2	1.93	0.51
1:A:410:GLY:HA2	1:A:530:ALA:HB2	1.93	0.51
1:A:176:THR:H	1:A:179:ASN:ND2	2.10	0.50
1:B:194:PRO:HB3	1:B:277:LEU:HD22	1.95	0.49
1:B:209:SER:O	1:B:228:HIS:HE1	1.95	0.49
1:B:188:ILE:HD12	1:B:266:LEU:HD21	1.93	0.49
1:A:270:PRO:HD3	1:A:307:TYR:CD2	2.48	0.49
1:A:313:ASP:HB3	2:A:739:HOH:O	2.12	0.49
1:B:165:PHE:HA	1:B:572:ILE:HD11	1.95	0.48
1:B:461:TYR:CE1	1:B:570:TRP:HD1	2.33	0.47
1:B:491:SER:H	1:B:494:GLN:HE21	1.62	0.47
1:A:407:LEU:HA	1:A:437:ALA:HB3	1.96	0.46
1:B:407:LEU:HA	1:B:437:ALA:HB3	1.98	0.46
1:B:419:ASP:HB2	2:B:755:HOH:O	2.14	0.46
1:B:572:ILE:H	1:B:572:ILE:HG13	1.67	0.45
1:B:117:ASP:OD1	1:B:119:THR:HG23	2.17	0.44
1:A:461:TYR:CE1	1:A:570:TRP:HD1	2.36	0.44
1:B:231:TRP:HA	1:B:235:GLN:HE21	1.83	0.44
1:A:188:ILE:HD12	1:A:266:LEU:HD21	1.99	0.44
1:B:78:TYR:HA	1:B:114:GLU:O	2.18	0.43
1:B:176:THR:H	1:B:179:ASN:ND2	2.16	0.43
1:A:191:GLY:HA2	1:A:192:PHE:HA	1.72	0.43
1:B:247:THR:HA	1:B:248:PRO:HD3	1.88	0.42
1:B:81:TYR:O	1:B:111:MET:HA	2.20	0.41
1:B:286:ASP:HB2	2:B:772:HOH:O	2.20	0.41
1:B:351:GLU:HG2	1:B:356:ARG:O	2.19	0.41
1:B:57:TYR:HB2	1:B:303:THR:HG23	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:209:SER:O	1:A:228:HIS:HE1	2.03	0.41
1:A:491:SER:H	1:A:494:GLN:HE21	1.67	0.41
1:A:234:TYR:HA	1:A:250:LYS:O	2.21	0.41
1:A:570:TRP:HA	1:A:571:GLU:HA	1.90	0.40
1:B:344:ARG:HD3	1:B:344:ARG:C	2.39	0.40
1:B:508:LYS:HA	1:B:512:LEU:HB2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1108:HOH:O	2:A:1112:HOH:O[1_445]	2.00	0.20

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	565/584 (97%)	552 (98%)	12 (2%)	1 (0%)	47 44
1	B	565/584 (97%)	547 (97%)	18 (3%)	0	100 100
All	All	1130/1168 (97%)	1099 (97%)	30 (3%)	1 (0%)	51 49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	171	TYR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	454/469 (97%)	450 (99%)	4 (1%)	78	83
1	B	454/469 (97%)	451 (99%)	3 (1%)	84	88
All	All	908/938 (97%)	901 (99%)	7 (1%)	81	86

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	ASN
1	A	404	GLN
1	A	423	VAL
1	A	572	ILE
1	B	119	THR
1	B	423	VAL
1	B	572	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	ASN
1	A	179	ASN
1	A	228	HIS
1	A	235	GLN
1	A	243	HIS
1	A	403	HIS
1	A	494	GLN
1	A	516	ASN
1	A	529	GLN
1	B	179	ASN
1	B	228	HIS
1	B	235	GLN
1	B	243	HIS
1	B	403	HIS
1	B	494	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	567/584 (97%)	-0.45	6 (1%)	80	79	9, 14, 21, 29	0
1	B	567/584 (97%)	-0.23	22 (3%)	39	38	8, 14, 32, 38	0
All	All	1134/1168 (97%)	-0.34	28 (2%)	57	56	8, 14, 27, 38	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	22	ALA	8.1
1	B	118	ALA	6.7
1	B	119	THR	5.6
1	B	31	TYR	4.7
1	B	205	VAL	4.6
1	B	514	ALA	4.6
1	B	121	CYS	4.6
1	A	119	THR	4.0
1	B	120	GLY	3.8
1	A	245	TYR	3.7
1	B	428	GLU	3.2
1	B	116	CYS	3.1
1	B	92	ILE	2.8
1	B	117	ASP	2.7
1	A	425	GLU	2.7
1	B	204	SER	2.7
1	B	427	GLY	2.7
1	A	325	ASP	2.7
1	B	245	TYR	2.6
1	B	425	GLU	2.6
1	B	73	THR	2.5
1	B	426	ASN	2.5
1	A	154	ASN	2.5
1	B	93	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	325	ASP	2.3
1	A	428	GLU	2.2
1	B	206	GLY	2.1
1	B	76	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.