



wwPDB X-ray Structure Validation Summary Report

Dec 18, 2023 – 12:55 am GMT

PDB ID : 4BMM
Title : Crystal structure of Trypanosoma cruzi CYP51 bound to the inhibitor (R)-N-(3-(1H-indol-3-yl)-1-oxo-1-(pyridin-4-ylamino)propan-2-yl)-2',3, 5'-trifluoro-(1,1'-biphenyl)-4-carboxamide
Authors : Choi, J.Y.; Calvet, C.M.; Gunatilleke, S.S.; Roush, W.R.; McKerrow, J.H.; Podust, L.M.
Deposited on : 2013-05-09
Resolution : 2.84 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1031 (2.86-2.82)
Clashscore	141614	1078 (2.86-2.82)
Ramachandran outliers	138981	1050 (2.86-2.82)
Sidechain outliers	138945	1051 (2.86-2.82)
RSRZ outliers	127900	1019 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	467	 53% 35% 6% 6%
1	B	467	 52% 37% 6% 5%
1	C	467	 57% 32% 6% 6%
1	D	467	 53% 36% 5% 5%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 14088 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called STEROL 14-ALPHA DEMETHYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	441	3431	2203	590	610	28	0	0	0
1	B	442	3432	2205	596	603	28	0	0	0
1	C	440	3415	2193	595	599	28	0	0	0
1	D	442	3448	2219	591	610	28	0	0	0

There are 68 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	21	MET	-	expression tag	UNP Q7Z1V1
A	22	ALA	-	expression tag	UNP Q7Z1V1
A	23	LYS	-	expression tag	UNP Q7Z1V1
A	24	LYS	-	expression tag	UNP Q7Z1V1
A	25	THR	-	expression tag	UNP Q7Z1V1
A	26	SER	-	expression tag	UNP Q7Z1V1
A	27	SER	-	expression tag	UNP Q7Z1V1
A	28	LYS	-	expression tag	UNP Q7Z1V1
A	29	GLY	-	expression tag	UNP Q7Z1V1
A	30	LYS	-	expression tag	UNP Q7Z1V1
A	31	LEU	-	expression tag	UNP Q7Z1V1
A	482	HIS	-	expression tag	UNP Q7Z1V1
A	483	HIS	-	expression tag	UNP Q7Z1V1
A	484	HIS	-	expression tag	UNP Q7Z1V1
A	485	HIS	-	expression tag	UNP Q7Z1V1
A	486	HIS	-	expression tag	UNP Q7Z1V1
A	487	HIS	-	expression tag	UNP Q7Z1V1
B	21	MET	-	expression tag	UNP Q7Z1V1
B	22	ALA	-	expression tag	UNP Q7Z1V1
B	23	LYS	-	expression tag	UNP Q7Z1V1
B	24	LYS	-	expression tag	UNP Q7Z1V1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	25	THR	-	expression tag	UNP Q7Z1V1
B	26	SER	-	expression tag	UNP Q7Z1V1
B	27	SER	-	expression tag	UNP Q7Z1V1
B	28	LYS	-	expression tag	UNP Q7Z1V1
B	29	GLY	-	expression tag	UNP Q7Z1V1
B	30	LYS	-	expression tag	UNP Q7Z1V1
B	31	LEU	-	expression tag	UNP Q7Z1V1
B	482	HIS	-	expression tag	UNP Q7Z1V1
B	483	HIS	-	expression tag	UNP Q7Z1V1
B	484	HIS	-	expression tag	UNP Q7Z1V1
B	485	HIS	-	expression tag	UNP Q7Z1V1
B	486	HIS	-	expression tag	UNP Q7Z1V1
B	487	HIS	-	expression tag	UNP Q7Z1V1
C	21	MET	-	expression tag	UNP Q7Z1V1
C	22	ALA	-	expression tag	UNP Q7Z1V1
C	23	LYS	-	expression tag	UNP Q7Z1V1
C	24	LYS	-	expression tag	UNP Q7Z1V1
C	25	THR	-	expression tag	UNP Q7Z1V1
C	26	SER	-	expression tag	UNP Q7Z1V1
C	27	SER	-	expression tag	UNP Q7Z1V1
C	28	LYS	-	expression tag	UNP Q7Z1V1
C	29	GLY	-	expression tag	UNP Q7Z1V1
C	30	LYS	-	expression tag	UNP Q7Z1V1
C	31	LEU	-	expression tag	UNP Q7Z1V1
C	482	HIS	-	expression tag	UNP Q7Z1V1
C	483	HIS	-	expression tag	UNP Q7Z1V1
C	484	HIS	-	expression tag	UNP Q7Z1V1
C	485	HIS	-	expression tag	UNP Q7Z1V1
C	486	HIS	-	expression tag	UNP Q7Z1V1
C	487	HIS	-	expression tag	UNP Q7Z1V1
D	21	MET	-	expression tag	UNP Q7Z1V1
D	22	ALA	-	expression tag	UNP Q7Z1V1
D	23	LYS	-	expression tag	UNP Q7Z1V1
D	24	LYS	-	expression tag	UNP Q7Z1V1
D	25	THR	-	expression tag	UNP Q7Z1V1
D	26	SER	-	expression tag	UNP Q7Z1V1
D	27	SER	-	expression tag	UNP Q7Z1V1
D	28	LYS	-	expression tag	UNP Q7Z1V1
D	29	GLY	-	expression tag	UNP Q7Z1V1
D	30	LYS	-	expression tag	UNP Q7Z1V1
D	31	LEU	-	expression tag	UNP Q7Z1V1
D	482	HIS	-	expression tag	UNP Q7Z1V1


Continued on next page...




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total	C	F	N	O	0	0
			38	29	3	4	2		
3	B	1	Total	C	F	N	O	0	0
			38	29	3	4	2		
3	C	1	Total	C	F	N	O	0	0
			38	29	3	4	2		
3	D	1	Total	C	F	N	O	0	0
			38	29	3	4	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	13	Total	O	0	0
			13	13		
4	B	6	Total	O	0	0
			6	6		
4	B	1	Total	O	0	0
			1	1		
4	C	9	Total	O	0	0
			9	9		
4	C	2	Total	O	0	0
			2	2		
4	D	7	Total	O	0	0
			7	7		



● Molecule 1: STEROL 14-ALPHA DEMETHYLASE



● Molecule 1: STEROL 14-ALPHA DEMETHYLASE



G463	N467	L470	K477	LYS
P464	Q468	V471	K477	LEU
	C469	K472	K477	PRO
	L470	Y473	K477	SER
	V471	T474	K477	HIS
	K472		K477	HIS
	T474		K477	HIS
	K477		K477	HIS
	LYS		K477	HIS
	LEU		K477	HIS
	PRO		K477	HIS
	SER		K477	HIS
	HIS		K477	HIS
	HIS		K477	HIS
	HIS		K477	HIS
	HIS		K477	HIS
	HIS		K477	HIS

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	272.47Å 66.45Å 122.22Å 90.00° 110.65° 90.00°	Depositor
Resolution (Å)	114.63 – 2.84 114.37 – 2.84	Depositor EDS
% Data completeness (in resolution range)	99.7 (114.63-2.84) 99.7 (114.37-2.84)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 2.86Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.189 , 0.290 0.189 , 0.289	Depositor DCC
R_{free} test set	2469 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	68.6	Xtrriage
Anisotropy	0.638	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 56.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14088	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: TU1, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/3515	0.73	0/4770
1	B	0.53	0/3516	0.71	0/4770
1	C	0.49	0/3497	0.70	1/4739 (0.0%)
1	D	0.59	1/3531 (0.0%)	0.77	2/4785 (0.0%)
All	All	0.54	1/14059 (0.0%)	0.73	3/19064 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	33	PRO	N-CD	5.11	1.55	1.47

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	107	THR	C-N-CD	6.41	141.86	128.40
1	D	32	PRO	C-N-CD	5.87	140.72	128.40
1	D	31	LEU	C-N-CD	5.40	139.75	128.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	39	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3431	0	3347	167	0
1	B	3432	0	3356	166	0
1	C	3415	0	3348	194	0
1	D	3448	0	3402	177	0
2	A	43	0	30	11	0
2	B	43	0	30	6	0
2	C	43	0	30	8	0
2	D	43	0	30	13	0
3	A	38	0	21	7	0
3	B	38	0	21	2	0
3	C	38	0	21	2	0
3	D	38	0	21	5	0
4	A	13	0	0	1	0
4	B	7	0	0	0	0
4	C	11	0	0	0	0
4	D	7	0	0	0	0
All	All	14088	0	13657	726	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 726 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:235:GLU:O	1:D:239:ILE:HG13	1.40	1.17
1:A:109:VAL:HG13	1:A:204:MET:CE	1.79	1.11
1:B:215:MET:HB2	1:B:218:LEU:HD12	1.24	1.10
1:A:109:VAL:HG13	1:A:204:MET:HE2	1.08	1.08
1:D:162:GLU:HB3	1:D:474:THR:HG23	1.38	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	437/467 (94%)	403 (92%)	32 (7%)	2 (0%)	29	51
1	B	438/467 (94%)	424 (97%)	13 (3%)	1 (0%)	47	69
1	C	436/467 (93%)	403 (92%)	33 (8%)	0	100	100
1	D	438/467 (94%)	414 (94%)	23 (5%)	1 (0%)	47	69
All	All	1749/1868 (94%)	1644 (94%)	101 (6%)	4 (0%)	47	69

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	91	SER
1	B	39	VAL
1	A	92	PRO
1	D	136	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/408 (87%)	312 (88%)	44 (12%)	4	9
1	B	354/408 (87%)	307 (87%)	47 (13%)	4	7
1	C	351/408 (86%)	311 (89%)	40 (11%)	5	11
1	D	359/408 (88%)	319 (89%)	40 (11%)	6	12
All	All	1420/1632 (87%)	1249 (88%)	171 (12%)	5	10

5 of 171 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	266	LEU
1	D	161	ASP
1	C	348	GLU
1	C	474	THR
1	D	233	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	334	ASN
1	D	334	ASN
1	C	44	HIS
1	D	148	HIS
1	B	458	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TU1	C	1460	2	41,42,42	1.99	3 (7%)	55,59,59	2.00	16 (29%)
2	HEM	B	1450	3,1	41,50,50	1.38	6 (14%)	45,82,82	1.94	11 (24%)
3	TU1	B	1460	2	41,42,42	1.97	6 (14%)	55,59,59	2.14	23 (41%)
3	TU1	A	1460	2	41,42,42	1.86	5 (12%)	55,59,59	1.65	11 (20%)
2	HEM	C	1450	3,1	41,50,50	1.34	6 (14%)	45,82,82	1.92	8 (17%)
3	TU1	D	1460	2	41,42,42	1.86	4 (9%)	55,59,59	1.61	9 (16%)
2	HEM	D	1450	3,1	41,50,50	1.43	6 (14%)	45,82,82	2.10	13 (28%)
2	HEM	A	1450	3,1	41,50,50	1.46	5 (12%)	45,82,82	2.29	14 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TU1	C	1460	2	-	0/23/24/24	0/5/5/5
2	HEM	B	1450	3,1	-	0/12/54/54	-
3	TU1	B	1460	2	-	0/23/24/24	0/5/5/5
3	TU1	A	1460	2	-	0/23/24/24	0/5/5/5
2	HEM	C	1450	3,1	-	4/12/54/54	-
3	TU1	D	1460	2	-	1/23/24/24	0/5/5/5
2	HEM	D	1450	3,1	-	4/12/54/54	-
2	HEM	A	1450	3,1	-	6/12/54/54	-

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	1460	TU1	C13-C14	8.52	1.50	1.39
3	C	1460	TU1	C13-C14	8.35	1.50	1.39
3	B	1460	TU1	C13-C14	7.52	1.49	1.39
3	B	1460	TU1	C9-C20	7.38	1.49	1.38
3	A	1460	TU1	C13-C14	7.24	1.49	1.39

The worst 5 of 105 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1450	HEM	CAD-C3D-C4D	6.42	135.87	124.66
2	A	1450	HEM	C1B-NB-C4B	5.79	111.05	105.07
2	D	1450	HEM	C1B-NB-C4B	5.63	110.89	105.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1450	HEM	C1B-NB-C4B	5.54	110.80	105.07
3	B	1460	TU1	C26-C25-C24	5.05	124.87	118.17

There are no chirality outliers.

5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1450	HEM	C3A-C2A-CAA-CBA
2	A	1450	HEM	C2D-C3D-CAD-CBD
2	A	1450	HEM	C4D-C3D-CAD-CBD
2	C	1450	HEM	C2D-C3D-CAD-CBD
2	C	1450	HEM	C4D-C3D-CAD-CBD


There are no ring outliers.



8 monomers are involved in 46 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1460	TU1	2	0
2	B	1450	HEM	6	0
3	B	1460	TU1	2	0
3	A	1460	TU1	7	0
2	C	1450	HEM	8	0
3	D	1460	TU1	5	0
2	D	1450	HEM	13	0
2	A	1450	HEM	11	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	441/467 (94%)	-0.51	0 100 100	47, 69, 109, 142	0
1	B	442/467 (94%)	-0.51	2 (0%) 91 89	57, 80, 114, 146	0
1	C	440/467 (94%)	-0.38	2 (0%) 91 89	60, 88, 128, 180	0
1	D	442/467 (94%)	-0.52	0 100 100	46, 68, 107, 140	0
All	All	1765/1868 (94%)	-0.48	4 (0%) 95 94	46, 77, 118, 180	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	214	PHE	4.1
1	C	158	TRP	2.4
1	B	375	PRO	2.2
1	C	164	VAL	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	TU1	B	1460	38/38	0.94	0.19	60,67,85,107	0
3	TU1	A	1460	38/38	0.95	0.24	51,55,77,99	0
3	TU1	C	1460	38/38	0.96	0.22	59,64,91,120	0
3	TU1	D	1460	38/38	0.97	0.17	51,55,67,82	0
2	HEM	A	1450	43/43	0.98	0.20	47,51,56,64	0
2	HEM	B	1450	43/43	0.98	0.18	56,59,62,68	0
2	HEM	C	1450	43/43	0.98	0.20	60,62,68,71	0
2	HEM	D	1450	43/43	0.98	0.19	50,52,55,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around TU1 A 1460:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around TU1 C 1460:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around TU1 D 1460:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around HEM A 1450:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM B 1450:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM C 1450:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.