Full wwPDB X-ray Structure Validation Report (i)

Jun 12, 2024 – 08:31 AM EDT

PDB ID	:	2BNA
Title	:	STRUCTURE OF A B-DNA DODECAMER AT 16 KELVIN
Authors	:	Drew, H.R.; Samson, S.; Dickerson, R.E.
Deposited on	:	1981-11-12
Resolution	:	2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3122 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	12	58% 42%
1	B	12	67% 33%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 569 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	12	243	116	46	70	11	0	0	0
1	B	12	243	116	46	70	11	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	37	37	37	0	0
2	B	46	46	46	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3')

Chain A:





- Molecule 1: DNA (5'-D(*CP*GP*CP*GP*AP*AP*TP*TP*CP*GP*CP*G)-3')

Chain B:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	23.44Å 39.31Å 65.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.70 20.13 – 2.71	Depositor EDS
% Data completeness (in resolution range)	(Not available) (15.00-2.70) 99.7 (20.13-2.71)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$	-	Xtriage
Refinement program	JACK-LEVITT	Depositor
R, R_{free}	0.151 , (Not available) 0.201 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å^2)	22.3	Xtriage
Anisotropy	1.916	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.13 , 53.7	EDS
L-test for twinning[1]	$< \|L\| > = 0.48$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	569	wwPDB-VP
Average B, all atoms (Å^2)	7.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.70% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	2.29	18/272 (6.6%)	2.20	18/418 (4.3%)
1	B	2.21	19/272 (7.0%)	2.16	16/418 (3.8%)
All	All	2.25	37/544 (6.8%)	2.18	34/836 (4.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	20	DT	C5-C6	8.47	1.40	1.34
1	A	7	DT	C5-C6	8.46	1.40	1.34
1	A	8	DT	C5-C6	8.31	1.40	1.34
1	A	4	DG	C8-N7	7.82	1.35	1.30
1	A	11	DC	C5-C6	7.73	1.40	1.34
1	A	5	DA	N7-C5	-7.56	1.34	1.39
1	A	12	DG	C8-N7	7.53	1.35	1.30
1	B	18	DA	N7-C5	-7.44	1.34	1.39
1	B	19	DT	C5-C6	7.40	1.39	1.34
1	A	10	DG	C8-N7	7.30	1.35	1.30
1	B	22	DG	C8-N7	6.84	1.35	1.30
1	A	7	DT	C4-C5	-6.84	1.38	1.45
1	B	24	DG	N7-C5	-6.72	1.35	1.39
1	B	21	DC	C5-C6	6.47	1.39	1.34
1	A	6	DA	N7-C5	-6.46	1.35	1.39
1	B	16	DG	C8-N7	6.45	1.34	1.30
1	B	14	DG	N7-C5	-6.29	1.35	1.39
1	B	13	DC	C5-C6	6.28	1.39	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	23	DC	C5-C6	6.26	1.39	1.34
1	A	2	DG	N7-C5	-6.26	1.35	1.39
1	A	3	DC	C5-C6	6.21	1.39	1.34
1	B	24	DG	C8-N7	6.13	1.34	1.30
1	A	10	DG	N7-C5	-6.03	1.35	1.39
1	B	15	DC	C5-C6	5.96	1.39	1.34
1	B	17	DA	N7-C5	-5.94	1.35	1.39
1	B	14	DG	C8-N7	5.90	1.34	1.30
1	A	12	DG	N7-C5	-5.89	1.35	1.39
1	A	2	DG	C8-N7	5.83	1.34	1.30
1	A	9	DC	C5-C6	5.81	1.39	1.34
1	B	17	DA	C8-N7	5.74	1.35	1.31
1	A	1	DC	C5-C6	5.74	1.39	1.34
1	A	8	DT	C4-C5	-5.71	1.39	1.45
1	B	22	DG	N7-C5	-5.70	1.35	1.39
1	B	20	DT	C4-C5	-5.46	1.40	1.45
1	B	19	DT	C4-C5	-5.45	1.40	1.45
1	B	16	DG	N7-C5	-5.35	1.36	1.39
1	A	4	DG	N7-C5	-5.13	1.36	1.39

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	19	DT	C2-N3-C4	-7.90	122.46	127.20
1	B	20	DT	C5-C6-N1	-7.84	118.99	123.70
1	A	8	DT	C2-N3-C4	-7.74	122.55	127.20
1	B	20	DT	C2-N3-C4	-7.62	122.62	127.20
1	B	19	DT	C5-C6-N1	-7.54	119.18	123.70
1	A	7	DT	C2-N3-C4	-7.39	122.77	127.20
1	B	15	DC	O4'-C4'-C3'	-7.36	101.56	104.50
1	A	8	DT	C5-C6-N1	-7.35	119.29	123.70
1	A	7	DT	C5-C6-N1	-6.91	119.56	123.70
1	A	8	DT	N1-C2-N3	6.77	118.66	114.60
1	B	19	DT	N3-C4-C5	6.42	119.05	115.20
1	B	20	DT	N1-C2-N3	6.06	118.24	114.60
1	A	1	DC	C3'-C2'-C1'	-6.01	95.29	102.50
1	B	19	DT	N1-C2-N3	5.91	118.14	114.60
1	A	1	DC	C6-N1-C2	5.86	122.64	120.30
1	B	24	DG	C4'-C3'-C2'	-5.79	97.89	103.10
1	A	7	DT	N3-C4-C5	5.76	118.66	115.20
1	A	11	DC	C5-C6-N1	-5.64	118.18	121.00
1	A	10	DG	C8-N9-C1'	5.58	134.26	127.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	5	DA	O4'-C1'-N9	-5.52	104.14	108.00
1	A	7	DT	N1-C2-N3	5.49	117.89	114.60
1	A	10	DG	C3'-C2'-C1'	-5.45	95.96	102.50
1	B	20	DT	N3-C4-C5	5.43	118.46	115.20
1	B	21	DC	C5-C6-N1	-5.41	118.30	121.00
1	A	8	DT	N3-C4-C5	5.32	118.39	115.20
1	A	1	DC	C5-C6-N1	-5.26	118.37	121.00
1	B	22	DG	N3-C4-N9	-5.26	122.85	126.00
1	A	2	DG	O4'-C1'-N9	-5.20	104.36	108.00
1	B	19	DT	C6-C5-C7	-5.20	119.78	122.90
1	B	19	DT	N3-C4-O4	-5.19	116.79	119.90
1	A	10	DG	N3-C4-N9	-5.18	122.89	126.00
1	B	13	DC	C5-C6-N1	-5.17	118.41	121.00
1	B	22	DG	C3'-C2'-C1'	-5.12	96.35	102.50
1	A	3	DC	O4'-C1'-N1	-5.05	104.47	108.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	10	DG	Sidechain
1	A	5	DA	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	243	0	136	2	0
1	B	243	0	136	6	0
2	A	37	0	0	1	0
2	B	46	0	0	6	0
All	All	569	0	272	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:DC:OP1	2:B:104:HOH:O	2.01	0.78
1:B:20:DT:C7	2:B:57:HOH:O	2.34	0.76
1:B:22:DG:H5"	2:B:94:HOH:O	1.84	0.75
1:A:11:DC:OP2	2:A:68:HOH:O	2.04	0.74
1:B:18:DA:OP2	2:B:85:HOH:O	2.09	0.70
1:A:3:DC:H2'	1:A:4:DG:C8	2.35	0.61
1:B:22:DG:C5'	2:B:94:HOH:O	2.52	0.47
1:B:20:DT:H73	2:B:57:HOH:O	2.10	0.44

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

There are no protein molecules in this entry.

5.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.