



wwPDB EM Validation Summary Report ⓘ

Mar 20, 2024 – 11:40 AM JST

PDB ID : 7C7A
EMDB ID : EMD-30297
Title : Cryo-EM structure of yeast Ribonuclease MRP with substrate ITS1
Authors : Lan, P.; Wu, J.; Lei, M.
Deposited on : 2020-05-24
Resolution : 2.80 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev70
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	
2	B	875	
3	C	195	
4	D	279	
5	E	173	
6	F	158	
7	G	140	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
8	H	133	
9	I	293	
9	J	293	
10	K	198	
11	L	201	
12	R	21	

2 Entry composition [i](#)

There are 14 unique types of molecules in this entry. The entry contains 27500 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called Ribonuclease MRP RNA subunit NME1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
1	A	330	6988	3130	1202	2326	330	0	0

- Molecule 2 is a protein called Ribonucleases P/MRP protein subunit POP1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	793	6453	4085	1170	1162	36	0	0

- Molecule 3 is a protein called Ribonucleases P/MRP protein subunit POP3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	152	1246	812	206	219	9	0	0

- Molecule 4 is a protein called RNases MRP/P 32.9 kDa subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	D	251	2086	1338	352	386	10	0	0

- Molecule 5 is a protein called Ribonuclease P/MRP protein subunit POP5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	E	145	1135	716	201	210	8	0	0

- Molecule 6 is a protein called Ribonucleases P/MRP protein subunit POP6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	F	157	1271	804	222	241	4	0	0

- Molecule 7 is a protein called Ribonucleases P/MRP protein subunit POP7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	G	121	960	609	167	182	2	0	0

- Molecule 8 is a protein called Ribonucleases P/MRP protein subunit POP8.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	H	131	1079	683	177	215	4	0	0

- Molecule 9 is a protein called Ribonuclease P/MRP protein subunit RPP1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	I	242	1881	1188	339	345	9	0	0
9	J	292	2253	1410	412	421	10	0	0

- Molecule 10 is a protein called Ribonuclease MRP protein subunit SNM1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	K	104	840	527	148	156	9	0	0

- Molecule 11 is a protein called Ribonuclease MRP protein subunit RMP1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	L	131	1110	711	210	182	7	0	0

- Molecule 12 is a RNA chain called Internal Transcribed Spacer 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
12	R	9	194	88	41	56	9	0	0

- Molecule 13 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
13	A	2	Total 2	Mg 2	0
13	R	1	Total 1	Mg 1	0


- Molecule 14 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
14	K	1	Total 1	Zn 1	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribonuclease MRP RNA subunit NME1




- Molecule 2: Ribonucleases P/MRP protein subunit POP1







• Molecule 3: Ribonucleases P/MRP protein subunit POP3



• Molecule 4: RNases MRP/P 32.9 kDa subunit



• Molecule 5: Ribonuclease P/MRP protein subunit POP5



• Molecule 6: Ribonucleases P/MRP protein subunit POP6

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	1037868	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.085	Depositor
Minimum map value	-0.036	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.006	Depositor
Map size (Å)	475.2, 475.2, 475.2	wwPDB
Map dimensions	432, 432, 432	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.1, 1.1, 1.1	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/7806	0.72	1/12148 (0.0%)
2	B	0.23	0/6593	0.39	0/8887
3	C	0.27	0/1281	0.52	0/1744
4	D	0.24	0/2123	0.41	0/2851
5	E	0.24	0/1148	0.39	0/1542
6	F	0.23	0/1289	0.41	0/1733
7	G	0.23	0/972	0.41	0/1305
8	H	0.23	0/1101	0.42	0/1488
9	I	0.23	0/1908	0.42	0/2577
9	J	0.23	0/2283	0.42	0/3080
10	K	0.24	0/856	0.46	0/1145
11	L	0.23	0/1132	0.40	0/1523
12	R	0.13	0/218	0.66	0/337
All	All	0.21	0/28710	0.53	1/40360 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
10	K	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	302	C	N3-C2-O2	-5.31	118.18	121.90

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
10	K	22	HIS	Peptide
10	K	99	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6988	0	3524	99	0
2	B	6453	0	6530	124	0
3	C	1246	0	1261	55	0
4	D	2086	0	2132	47	0
5	E	1135	0	1199	7	0
6	F	1271	0	1305	5	0
7	G	960	0	1005	14	0
8	H	1079	0	1037	25	0
9	I	1881	0	1975	36	0
9	J	2253	0	2344	37	0
10	K	840	0	830	24	0
11	L	1110	0	1138	26	0
12	R	194	0	100	0	0
13	A	2	0	0	0	0
13	R	1	0	0	0	0
14	K	1	0	0	0	0
All	All	27500	0	24380	445	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 445 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:671:ILE:HG22	2:B:675:GLN:HE22	1.37	0.88
11:L:63:LYS:HG2	11:L:66:GLU:HB2	1.61	0.81
2:B:311:LYS:HG2	2:B:323:ASN:HD21	1.46	0.80
9:J:120:SER:HA	9:J:124:LEU:HB2	1.62	0.79
2:B:353:LEU:HG	2:B:374:ILE:HD11	1.63	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	785/875 (90%)	731 (93%)	53 (7%)	1 (0%)	51	81
3	C	150/195 (77%)	128 (85%)	22 (15%)	0	100	100
4	D	247/279 (88%)	239 (97%)	8 (3%)	0	100	100
5	E	143/173 (83%)	139 (97%)	4 (3%)	0	100	100
6	F	155/158 (98%)	154 (99%)	1 (1%)	0	100	100
7	G	117/140 (84%)	112 (96%)	5 (4%)	0	100	100
8	H	129/133 (97%)	122 (95%)	7 (5%)	0	100	100
9	I	240/293 (82%)	224 (93%)	15 (6%)	1 (0%)	34	66
9	J	290/293 (99%)	274 (94%)	16 (6%)	0	100	100
10	K	100/198 (50%)	91 (91%)	8 (8%)	1 (1%)	15	44
11	L	129/201 (64%)	122 (95%)	7 (5%)	0	100	100
All	All	2485/2938 (85%)	2336 (94%)	146 (6%)	3 (0%)	54	81

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
10	K	23	VAL
2	B	593	ASN
9	I	138	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	718/785 (92%)	716 (100%)	2 (0%)	92	98
3	C	144/185 (78%)	143 (99%)	1 (1%)	84	95
4	D	235/261 (90%)	235 (100%)	0	100	100
5	E	133/160 (83%)	133 (100%)	0	100	100
6	F	148/149 (99%)	147 (99%)	1 (1%)	84	95
7	G	110/127 (87%)	110 (100%)	0	100	100
8	H	122/123 (99%)	122 (100%)	0	100	100
9	I	216/258 (84%)	216 (100%)	0	100	100
9	J	257/258 (100%)	256 (100%)	1 (0%)	91	97
10	K	95/184 (52%)	94 (99%)	1 (1%)	73	92
11	L	118/182 (65%)	117 (99%)	1 (1%)	81	94
All	All	2296/2672 (86%)	2289 (100%)	7 (0%)	92	98

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	F	132	ARG
9	J	144	LYS
11	L	16	ARG
10	K	3	LYS
3	C	175	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
9	J	40	HIS
9	J	95	GLN
10	K	41	HIS
3	C	59	GLN
3	C	32	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	328/340 (96%)	79 (24%)	16 (4%)
12	R	8/21 (38%)	2 (25%)	0
All	All	336/361 (93%)	81 (24%)	16 (4%)

5 of 81 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	16	A
1	A	25	A
1	A	26	A
1	A	36	U
1	A	40	A

5 of 16 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	299	U
1	A	290	U
1	A	188	U
1	A	289	C
1	A	143	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-30297. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 210



Y Index: 219




Z Index: 226


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.006. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 358 nm^3 ; this corresponds to an approximate mass of 323 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.357\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-30297 and PDB model 7C7A. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.006 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.006).

9.4 Atom inclusion [i](#)



At the recommended contour level, 94% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.006) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9020	 0.4040
A	 0.9160	 0.3850
B	 0.9240	 0.4370
C	 0.5970	 0.2150
D	 0.8930	 0.3940
E	 0.9340	 0.4500
F	 0.9450	 0.4450
G	 0.9350	 0.4670
H	 0.8720	 0.3460
I	 0.9330	 0.4460
J	 0.9130	 0.3980
K	 0.8570	 0.3730
L	 0.9120	 0.4130
R	 0.9800	 0.4660

