Full wwPDB X-ray Structure Validation Report (i)

Jun 13, 2024 – 02:53 AM EDT

PDB ID	:	4CDN
Title	:	Crystal structure of M. mazei photolyase with its in vivo reconstituted 8-HDF antenna chromophore
Authors	:	Kiontke, S.; Batschauer, A.; Essen, L.-O.
Deposited on	:	2013-11-01
Resolution	:	1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	4% 93% . .
1	B	482	4% 85% 6% 9%

2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 8267 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DEOXYRIBODIPYRIMIDINE PHOTOLYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	463	3762	2420	630	697	15	0	9	0
1	B	438	3564	2291	596	662	15	0	7	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	expression tag	UNP Q8PYK9
A	-16	GLY	-	expression tag	UNP Q8PYK9
A	-15	SER	-	expression tag	UNP Q8PYK9
A	-14	SER	-	expression tag	UNP Q8PYK9
A	-13	HIS	-	expression tag	UNP Q8PYK9
A	-12	HIS	-	expression tag	UNP Q8PYK9
A	-11	HIS	-	expression tag	UNP Q8PYK9
A	-10	HIS	-	expression tag	UNP Q8PYK9
A	-9	HIS	-	expression tag	UNP Q8PYK9
A	-8	HIS	-	expression tag	UNP Q8PYK9
A	-7	SER	-	expression tag	UNP Q8PYK9
A	-6	SER	-	expression tag	UNP Q8PYK9
A	-5	GLY	-	expression tag	UNP Q8PYK9
A	-4	LEU	-	expression tag	UNP Q8PYK9
A	-3	VAL	-	expression tag	UNP Q8PYK9
A	-2	PRO	-	expression tag	UNP Q8PYK9
A	-1	ARG	-	expression tag	UNP Q8PYK9
A	0	GLY	-	expression tag	UNP Q8PYK9
A	1	SER	-	expression tag	UNP Q8PYK9
A	2	HIS	-	expression tag	UNP Q8PYK9
A	377	THR	MET	cloning artifact	UNP Q8PYK9
B	-17	MET	-	expression tag	UNP Q8PYK9
B	-16	GLY	-	expression tag	UNP Q8PYK9
B	-15	SER	-	expression tag	UNP Q8PYK9
B	-14	SER	-	expression tag	UNP Q8PYK9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-13	HIS	-	expression tag	UNP Q8PYK9
B	-12	HIS	-	expression tag	UNP Q8PYK9
B	-11	HIS	-	expression tag	UNP Q8PYK9
B	-10	HIS	-	expression tag	UNP Q8PYK9
B	-9	HIS	-	expression tag	UNP Q8PYK9
B	-8	HIS	-	expression tag	UNP Q8PYK9
B	-7	SER	-	expression tag	UNP Q8PYK9
B	-6	SER	-	expression tag	UNP Q8PYK9
B	-5	GLY	-	expression tag	UNP Q8PYK9
B	-4	LEU	-	expression tag	UNP Q8PYK9
B	-3	VAL	-	expression tag	UNP Q8PYK9
B	-2	PRO	-	expression tag	UNP Q8PYK9
B	-1	ARG	-	expression tag	UNP Q8PYK9
B	0	GLY	-	expression tag	UNP Q8PYK9
B	1	SER	-	expression tag	UNP Q8PYK9
B	2	HIS	-	expression tag	UNP Q8PYK9
B	377	THR	MET	cloning artifact	UNP Q8PYK9

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0
			Total	C	N	O	P		
2	B	1	53	27	9	15	2	0	0

- Molecule 3 is 1-deoxy-1-(8-hydroxy-2,4-dioxo-3,4-dihydropyrimido[4,5-b]quinolin-10(2H)-yl) -D-ribitol (three-letter code: FO1) (formula: $C_{16}H_{17}N_3O_7$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	26	16	3	7	0	0
			Total	C	N	O		
3	B	1	26	16	3	7	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	6	3	3	0	0
			Total	C	O		
4	A	1	6	3	3	0	0
			Total	C	O		
4	A	1	6	3	3	0	0
			Total	C	O		
4	B	1	6	3	3	0	0
			Total	C	O		
4	B	1	6	3	3	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
5	A	1	5	4	1	0	0
			Total	O	S		
5	B	1	5	4	1	0	0
			Total	O	S		
5	B	1	5	4	1	0	0

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	A	1	10	6	4	0	0
6	A	1	10	6	4	0	0

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	A	1	7	4	3	0	0
7	A	1	7	4	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	A	1	7	4	3	0	0
			Total	C	O		
7	A	1	7	4	3	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
8	A	377	377	377	0	0
			Total	O		
8	B	313	313	313	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DEOXYRIBODIPYRIMIDINE PHOTOLYASE

Chain A:





• Molecule 1: DEOXYRIBODIPYRIMIDINE PHOTOLYASE

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	78.97Å 114.74Å 141.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.47 – 1.90 34.47 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.1 (34.47-1.90) 98.1 (34.47-1.90)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.18 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.149 , 0.178 0.158 , 0.188	Depositor DCC
R_{free} test set	2025 reflections (2.03%)	wwPDB-VP
Wilson B-factor (Å^2)	25.8	Xtriage
Anisotropy	0.492	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.40 , 51.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	8267	wwPDB-VP
Average B, all atoms (Å^2)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PGE, FAD, GOL, FO1, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.51	0/3890	0.65	0/5266
1	B	0.50	0/3685	0.63	0/4990
All	All	0.51	0/7575	0.64	0/10256

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3762	0	3681	14	0
1	B	3564	0	3444	25	0
2	A	53	0	31	2	0
2	B	53	0	31	3	0
3	A	26	0	16	0	0
3	B	26	0	16	0	0
4	A	18	0	24	0	0
4	B	12	0	16	2	0
5	A	5	0	0	0	0
5	B	10	0	0	0	0
6	A	20	0	28	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	28	0	40	3	0
8	A	377	0	0	3	0
8	B	313	0	0	7	0
All	All	8267	0	7327	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:PRO:HB3	6:A:1469:PGE:H4	1.63	0.79
1:A:387:GLU:HG2	7:A:1470:PEG:H11	1.73	0.70
1:A:387:GLU:HG2	7:A:1470:PEG:C1	2.26	0.65
1:B:336:HIS:HD2	1:B:339:ASP:OD2	1.84	0.61
1:B:365[A]:MET:HA	1:B:365[A]:MET:HE3	1.82	0.61
1:A:51:VAL:H	1:A:184:ASN:HD21	1.48	0.60
1:B:374:HIS:CD2	1:B:376:TYR:H	2.21	0.58
1:B:336:HIS:HE1	1:B:433:GLU:OE2	1.87	0.58
1:B:103:ARG:CB	8:B:2092:HOH:O	2.53	0.57
1:B:374:HIS:HD2	1:B:376:TYR:H	1.53	0.55
1:B:3:MET:HB3	1:B:39:PHE:CZ	2.44	0.53
1:A:462:SER:C	8:A:2371:HOH:O	2.48	0.51
1:B:336:HIS:CD2	1:B:339:ASP:OD2	2.62	0.51
1:A:459:GLU:HG3	8:A:2368:HOH:O	2.14	0.47
1:B:244:PHE:CD1	1:B:265:SER:HA	2.49	0.47
1:B:366:GLU:OE2	4:B:1463:GOL:H32	2.15	0.47
1:B:58:THR:HG22	1:B:97:PRO:HD3	1.97	0.46
1:B:365[A]:MET:HA	1:B:365[A]:MET:CE	2.43	0.46
1:B:156:HIS:CD2	1:B:316:ASN:HD21	2.34	0.46
1:B:263:MET:HB3	8:B:2179:HOH:O	2.16	0.45
1:A:68:GLN:HG2	1:A:411:ARG:NH2	2.31	0.45
1:A:244:PHE:CD1	1:A:265:SER:HA	2.51	0.45
1:A:268:SER:HB3	2:A:998:FAD:H5'2	1.98	0.44
1:B:268:SER:HB3	2:B:998:FAD:H5'2	2.00	0.44
1:B:188:PRO:CD	8:B:2083:HOH:O	2.64	0.43
1:B:336:HIS:CE1	1:B:433:GLU:OE2	2.69	0.43
1:A:323:GLU:OE2	8:A:2279:HOH:O	2.21	0.43
1:A:410:GLY:HA2	2:A:998:FAD:O4	2.19	0.42
1:B:263:MET:CG	8:B:2179:HOH:O	2.66	0.42
1:B:145:ASN:HD22	1:B:307:GLU:HG2	1.84	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:374:HIS:HE1	8:B:2185:HOH:O	2.01	0.42
1:A:58[A]:THR:HG22	1:A:97:PRO:HD3	2.01	0.42
1:A:387:GLU:HG2	7:A:1470:PEG:H12	2.00	0.42
1:A:-4:LEU:HG	6:A:1469:PGE:H32	2.00	0.42
1:B:188:PRO:HD3	8:B:2083:HOH:O	2.19	0.42
1:B:301:GLU:HB3	2:B:998:FAD:C8A	2.51	0.41
1:B:68:GLN:HG2	1:B:411:ARG:NH2	2.36	0.41
2:B:998:FAD:H62A	4:B:1464:GOL:H2	1.85	0.41
1:B:263:MET:CB	8:B:2179:HOH:O	2.69	0.41
1:B:171:LEU:N	1:B:172:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	468/482 (97%)	462 (99%)	6 (1%)	0	100	100
1	B	441/482 (92%)	433 (98%)	8 (2%)	0	100	100
All	All	909/964 (94%)	895 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	404/416 (97%)	402 (100%)	2 (0%)	88 89
1	B	381/416 (92%)	379 (100%)	2 (0%)	88 89
All	All	785/832 (94%)	781 (100%)	4 (0%)	88 89

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-4	LEU
1	A	196	MET
1	B	1	SER
1	B	220	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	145	ASN
1	A	184	ASN
1	A	316	ASN
1	B	145	ASN
1	B	156	HIS
1	B	257	ASN
1	B	336	HIS
1	B	343	HIS
1	B	374	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
7	PEG	A	1470	-	6,6,6	0.52	0	5,5,5	0.60	0
7	PEG	A	1468	-	6,6,6	0.50	0	5,5,5	0.34	0
4	GOL	B	1464	-	5,5,5	0.19	0	5,5,5	1.24	1 (20%)
4	GOL	B	1463	-	5,5,5	0.25	0	5,5,5	0.45	0
5	SO4	B	1466	-	4,4,4	0.43	0	6,6,6	0.19	0
5	SO4	B	1465	-	4,4,4	0.51	0	6,6,6	0.25	0
6	PGE	A	1469	-	9,9,9	0.55	0	8,8,8	0.41	0
5	SO4	A	1466	-	4,4,4	0.46	0	6,6,6	0.24	0
6	PGE	A	1467	-	9,9,9	0.65	0	8,8,8	0.47	0
7	PEG	A	1471	-	6,6,6	0.53	0	5,5,5	0.50	0
7	PEG	A	1472	-	6,6,6	0.57	0	5,5,5	0.40	0
4	GOL	A	1465	-	5,5,5	0.30	0	5,5,5	0.24	0
3	FO1	B	999	-	28,28,28	1.50	5 (17%)	38,41,41	1.81	10 (26%)
2	FAD	A	998	-	54,58,58	1.23	5 (9%)	71,89,89	1.34	11 (15%)
4	GOL	A	1464	-	5,5,5	0.32	0	5,5,5	0.55	0
2	FAD	B	998	-	54,58,58	1.24	6 (11%)	71,89,89	1.28	8 (11%)
3	FO1	A	999	-	28,28,28	1.43	3 (10%)	38,41,41	1.51	6 (15%)
4	GOL	A	1463	-	5,5,5	0.40	0	5,5,5	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	PEG	A	1470	-	-	3/4/4/4	-
7	PEG	A	1468	-	-	2/4/4/4	-
4	GOL	B	1464	-	-	2/4/4/4	-
4	GOL	B	1463	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PGE	A	1469	-	-	4/7/7/7	-
6	PGE	A	1467	-	-	5/7/7/7	-
7	PEG	A	1471	-	-	1/4/4/4	-
7	PEG	A	1472	-	-	3/4/4/4	-
4	GOL	A	1465	-	-	0/4/4/4	-
3	FO1	B	999	-	-	2/14/14/14	0/3/3/3
2	FAD	A	998	-	-	1/30/50/50	0/6/6/6
4	GOL	A	1464	-	-	0/4/4/4	-
2	FAD	B	998	-	-	1/30/50/50	0/6/6/6
3	FO1	A	999	-	-	3/14/14/14	0/3/3/3
4	GOL	A	1463	-	-	4/4/4/4	-

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	998	FAD	C9A-C5X	4.74	1.48	1.41
2	A	998	FAD	C9A-C5X	4.71	1.48	1.41
3	B	999	FO1	C5-C12	4.52	1.48	1.41
3	A	999	FO1	C5-C12	4.41	1.48	1.41
3	A	999	FO1	C14-C15	-3.27	1.48	1.52
2	A	998	FAD	C8-C7	3.09	1.48	1.40
2	B	998	FAD	C8-C7	3.01	1.48	1.40
2	A	998	FAD	C4X-N5	2.88	1.37	1.30
3	B	999	FO1	C4-C3	2.72	1.41	1.35
2	B	998	FAD	C4X-N5	2.53	1.36	1.30
2	A	998	FAD	C2-N3	-2.47	1.33	1.39
2	A	998	FAD	C2A-N3A	2.41	1.35	1.32
3	B	999	FO1	C3-C13	2.36	1.49	1.44
3	B	999	FO1	C3-C2	2.36	1.49	1.45
2	B	998	FAD	C4-N3	-2.06	1.35	1.38
3	A	999	FO1	C4-C3	2.06	1.40	1.35
2	B	998	FAD	O4B-C1B	2.04	1.43	1.40
2	B	998	FAD	P-O3P	2.02	1.61	1.59
3	B	999	FO1	C1-N2	-2.02	1.34	1.39

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	999	FO1	C2-C3-C13	-5.79	114.45	117.48
2	A	998	FAD	N3A-C2A-N1A	-4.77	122.20	128.67
3	B	999	FO1	C5-C4-C3	-3.72	118.43	122.06

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	B	998	FAD	N3A-C2A-N1A	-3.67	123.69	128.67
3	A	999	FO1	O2-C2-C3	-3.59	118.08	125.98
3	B	999	FO1	C3-C2-N2	3.34	120.29	114.45
3	A	999	FO1	C5-C4-C3	-3.33	118.81	122.06
3	A	999	FO1	C2-C3-C13	-3.19	115.81	117.48
2	A	998	FAD	O4-C4-C4X	-2.95	118.73	126.53
2	B	998	FAD	C9A-C5X-N5	-2.82	119.47	122.45
2	A	998	FAD	C4-C4X-N5	2.80	122.08	118.21
2	B	998	FAD	O4B-C1B-N9A	2.59	112.18	108.75
2	B	998	FAD	O4-C4-C4X	-2.53	119.85	126.53
2	A	998	FAD	C10-N1-C2	2.51	122.29	116.85
3	B	999	FO1	O2-C2-C3	-2.50	120.47	125.98
3	A	999	FO1	C17-C16-C15	-2.49	109.43	113.57
3	B	999	FO1	C13-N1-C1	2.43	122.11	116.85
2	A	998	FAD	C9A-C5X-N5	-2.43	119.88	122.45
2	A	998	FAD	O2P-P-O1P	2.33	123.31	112.44
2	A	998	FAD	N6A-C6A-N1A	2.33	123.31	118.33
2	A	998	FAD	C4X-C4-N3	2.31	119.13	113.25
3	B	999	FO1	C4-C3-C2	2.27	124.21	119.30
2	A	998	FAD	C5X-C9A-N10	2.26	120.01	117.97
3	B	999	FO1	C17-C16-C15	-2.25	109.83	113.57
3	A	999	FO1	C3-C2-N2	2.22	118.34	114.45
4	B	1464	GOL	O1-C1-C2	-2.18	100.57	110.38
3	B	999	FO1	C15-C14-N3	2.16	120.43	110.20
2	B	998	FAD	O2-C2-N1	-2.11	118.30	121.80
2	A	998	FAD	C4X-C10-N10	2.11	119.50	116.48
2	B	998	FAD	N3-C2-N1	2.11	123.98	119.50
3	B	999	FO1	C5-C12-N3	2.10	121.19	118.72
2	B	998	FAD	C4X-C10-N1	-2.09	119.46	124.59
3	B	999	FO1	C2-N2-C1	-2.07	121.97	125.64
2	B	998	FAD	C4-N3-C2	-2.05	122.01	125.64
2	A	998	FAD	C5'-C4'-C3'	-2.03	108.39	112.22
3	A	999	FO1	C15-C14-N3	2.00	119.66	110.20

There are no chirality outliers.

All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	999	FO1	N3-C14-C15-O3
3	A	999	FO1	N3-C14-C15-C16
3	B	999	FO1	N3-C14-C15-O3
3	B	999	FO1	N3-C14-C15-C16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	1463	GOL	O1-C1-C2-C3
4	A	1463	GOL	C1-C2-C3-O3
4	A	1463	GOL	O2-C2-C3-O3
4	B	1464	GOL	O1-C1-C2-C3
7	A	1470	PEG	O1-C1-C2-O2
6	A	1469	PGE	C6-C5-O3-C4
6	A	1467	PGE	C3-C4-O3-C5
6	A	1467	PGE	O3-C5-C6-O4
7	A	1472	PEG	O2-C3-C4-O4
4	A	1463	GOL	O1-C1-C2-O2
4	B	1464	GOL	O1-C1-C2-O2
7	A	1470	PEG	O2-C3-C4-O4
6	A	1467	PGE	C6-C5-O3-C4
6	A	1469	PGE	O2-C3-C4-O3
4	B	1463	GOL	O1-C1-C2-O2
6	A	1467	PGE	O1-C1-C2-O2
7	A	1472	PEG	O1-C1-C2-O2
6	A	1467	PGE	C1-C2-O2-C3
2	A	998	FAD	C4'-C5'-O5'-P
6	A	1469	PGE	C3-C4-O3-C5
6	A	1469	PGE	C4-C3-O2-C2
7	A	1470	PEG	C4-C3-O2-C2
4	B	1463	GOL	O1-C1-C2-C3
7	A	1468	PEG	C4-C3-O2-C2
7	A	1468	PEG	O1-C1-C2-O2
2	B	998	FAD	C4'-C5'-O5'-P
7	A	1472	PEG	C4-C3-O2-C2
3	A	999	FO1	O4-C16-C17-C18
7	A	1471	PEG	O1-C1-C2-O2

There are no ring outliers.

6 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1470	PEG	3	0
4	B	1464	GOL	1	0
4	B	1463	GOL	1	0
6	A	1469	PGE	2	0
2	A	998	FAD	2	0
2	B	998	FAD	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FO1 B 999



Bond lengths



Bond angles



Torsions



Rings

Ligand FAD A 998



Bond lengths



Bond angles



Torsions



Rings

Ligand FAD B 998



Bond lengths



Bond angles



Torsions



Rings

Ligand FO1 A 999



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	463/482 (96%)	-0.09	18 (3%)	39	42	17, 26, 45, 90	6 (1%)
1	B	438/482 (90%)	0.00	20 (4%)	32	35	18, 29, 53, 83	3 (0%)
All	All	901/964 (93%)	-0.04	38 (4%)	36	39	17, 28, 50, 90	9 (0%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	187	THR	6.8
1	A	189	GLU	6.8
1	B	220	ASN	6.7
1	A	195	GLY	6.5
1	A	190	LEU	6.4
1	B	186	VAL	5.7
1	A	187	THR	5.2
1	A	196	MET	4.9
1	B	188	PRO	4.8
1	B	219	LYS	4.7
1	B	218	LEU	4.6
1	B	-1	ARG	3.9
1	B	14	GLY	3.9
1	B	221	LYS	3.6
1	A	-3	VAL	3.5
1	A	218	LEU	3.3
1	B	0	GLY	3.1
1	A	-4	LEU	3.1
1	A	228	TRP	3.0
1	B	18	ASP	3.0
1	A	462	SER	2.9
1	B	108	TYR	2.9
1	A	215	ARG	2.9
1	A	450	ARG	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	185	SER	2.7
1	B	-3	VAL	2.5
1	B	-2	PRO	2.4
1	A	188	PRO	2.4
1	B	158	TYR	2.4
1	A	-1	ARG	2.2
1	A	273	PHE	2.2
1	A	274	GLY	2.2
1	B	66	ILE	2.2
1	A	18[A]	ASP	2.1
1	A	186	VAL	2.1
1	B	47	ALA	2.1
1	B	109	ASN	2.1
1	B	87	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
7	PEG	A	1471	7/7	0.63	0.29	56,67,75,75	0
7	PEG	A	1472	7/7	0.72	0.25	60,63,68,68	0
6	PGE	A	1469	10/10	0.82	0.22	47,52,55,57	0
6	PGE	A	1467	10/10	0.87	0.14	43,51,54,55	0
5	SO4	B	1466	5/5	0.88	0.16	82,85,91,94	0
7	PEG	A	1470	7/7	0.88	0.17	32,44,56,57	0
4	GOL	A	1465	6/6	0.90	0.12	53,54,55,58	0
4	GOL	B	1464	6/6	0.91	0.17	28,33,38,38	0
7	PEG	A	1468	7/7	0.92	0.14	36,47,57,59	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	GOL	A	1463	6/6	0.92	0.16	39,43,43,48	0
4	GOL	B	1463	6/6	0.94	0.12	32,40,42,43	0
3	FO1	B	999	26/26	0.95	0.12	19,22,25,26	0
5	SO4	B	1465	5/5	0.95	0.12	50,60,62,62	0
4	GOL	A	1464	6/6	0.95	0.15	30,35,36,38	0
5	SO4	A	1466	5/5	0.97	0.15	44,56,59,61	0
2	FAD	A	998	53/53	0.98	0.14	17,19,22,23	0
2	FAD	B	998	53/53	0.98	0.13	18,20,22,23	0
3	FO1	A	999	26/26	0.98	0.16	18,19,21,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FO1 B 999:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FAD A 998:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around FAD B 998:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FO1 A 999:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.