wwPDB NMR Structure Validation Summary Report (i)

Feb 8, 2022 – 02:55 PM EST

PDB ID	:	1COP
Title	:	THREE-DIMENSIONAL DIMER STRUCTURE OF THE LAMBDA-CRO REPRESSOR IN SOLUTION AS DETERMINED BY HETERONUCLEAR MULTIDIMENSIONAL NMR
Authors	:	Matsuo, H.; Shirakawa, M.; Kyogoku, Y.
Deposited on	:	1995-06-23

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.26
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	D	66	18% 61% 9% 12%
1	E	66	30% 47% 9% 14%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 7 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	D:2-D:59, E:3-E:59 (115)	1.00	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 4, 7, 8, 12, 14, 15, 17, 19
2	5, 10, 11, 13, 16, 20
3	1, 6
4	9, 18
Single-model clusters	3

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2104 atoms, of which 1068 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called CRO REPRESSOR.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	D	66	1052	328	534	91	97	2	0
1	E	66	1052	328	534	91	97	2	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: CRO REPRESSOR

Chain D: 18% 61% 9% 12%



• Molecule 1: CRO REPRESSOR

Chain E: 30% 47% 9% 14%



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 7. Colouring as in section 4.1 above.

• Molecule 1: CRO REPRESSOR

Chain D: 32% 41% 14% • 12%



• Molecule 1: CRO REPRESSOR

Chain E: 32% 45% 9% 14%


M1
E2
Q3
R4
I5
T6
L7
K8
D9
Y10
A11
M12
R13
F14
G15
Q16
T17
K18
T19
A20
K21
D22
L23
G24
V25
Y26
Q27
S28
A29
I30
R38
K39
I40
F41
L42
T43
I44
N45
A46
D47
G48
S49
E53
E54
V55
K56
P57
F58
P59
S60
N61
K62
K63
T64
T65
A66

5 Refinement protocol and experimental data overview (i)

Of the ? calculated structures, 20 were deposited, based on the following criterion: *?*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.1

No chemical shift data was provided.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	D	0.0±0.0	2.9±0.3
1	E	0.0±0.0	2.8±0.4
All	All	0	114

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

5 of 6 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	D	4	ARG	Sidechain	20
1	E	4	ARG	Sidechain	20
1	D	13	ARG	Sidechain	19
1	D	38	ARG	Sidechain	19
1	E	13	ARG	Sidechain	18

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	D	458	467	467	41±6
1	E	449	461	461	36±5
All	All	18140	18560	18560	1429

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

5 of 522 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:D:33:ALA:HB1	1:D:40:ILE:CD1	1.01	1.85	19	10
1:E:42:LEU:HD12	1:E:52:ALA:HB2	0.97	1.32	8	7
1:E:4:ARG:O	1:E:5:ILE:HD13	0.95	1.61	9	4
1:D:58:PHE:CZ	1:E:25:VAL:HG11	0.94	1.97	17	3
1:D:55:VAL:CG1	1:E:55:VAL:HG13	0.93	1.93	9	3

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	58/66 (88%)	52±1 (89±2%)	5±1 (9±2%)	1±1 (2±2%)	11	52
1	E	57/66 (86%)	51±2 (90±3%)	4±2 (8±3%)	1±1 (2±2%)	9	46
All	All	2300/2640 (87%)	2057 (89%)	193 (8%)	50 (2%)	10	49

5 of 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	E	46	ALA	15
1	D	46	ALA	10
1	D	25	VAL	6
1	D	2	GLU	3
1	E	3	GLN	3

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	D	47/54 (87%)	34±2 (71±4%)	13±2 (29±4%)	2 18
1	E	46/54 (85%)	33±2 (72±5%)	13±2 (28±5%)	2 19
All	All	1860/2160 (86%)	1334 (72%)	526 (28%)	2 19

5 of 70 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	D	55	VAL	20
1	E	55	VAL	20
1	D	26	TYR	18
1	E	26	TYR	18
1	E	49	SER	17

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided