wwPDB X-ray Structure Validation Summary Report (i)

Jun 19, 2024 – 05:04 AM EDT

PDB ID	:	4CO6
Title	:	Crystal structure of the Nipah virus RNA free nucleoprotein- phosphoprotein complex
Authors	:	Yabukarksi, F.; Lawrence, P.; Tarbouriech, N.; Bourhis, J.M.; Jensen, M.R.; Ruigrok, R.W.H.; Blackledge, M.; Volchkov, V.; Jamin, M.
Deposited on	:	2014-01-27
Resolution	:	2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	3%, 71%, 18%, •, 7%
1	B	356	%, 70%, 19%, •, 9%
1	C	356	14%, 46%, 9%, •, 44%
2	D	52	2%, 63%, 13%, 23%
2	E	52	6%, 62%, 13%, 25%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	52	2% 48% 21% • 29%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7650 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NUCLEOPROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	330	2599	1666	443	474	16	0	1	0
1	B	324	2574	1649	441	469	15	0	0	0
1	C	200	1444	920	248	267	9	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	GLY	-	expression tag	UNP Q9IK92
A	29	ALA	-	expression tag	UNP Q9IK92
A	30	MSE	-	expression tag	UNP Q9IK92
A	31	ALA	-	expression tag	UNP Q9IK92
B	28	GLY	-	expression tag	UNP Q9IK92
B	29	ALA	-	expression tag	UNP Q9IK92
B	30	MSE	-	expression tag	UNP Q9IK92
B	31	ALA	-	expression tag	UNP Q9IK92
C	28	GLY	-	expression tag	UNP Q9IK92
C	29	ALA	-	expression tag	UNP Q9IK92
C	30	MSE	-	expression tag	UNP Q9IK92
C	31	ALA	-	expression tag	UNP Q9IK92

- Molecule 2 is a protein called PHOSPHOPROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	40	317	198	55	63	1	0	0	0
2	E	39	309	192	54	62	1	0	0	0
2	F	37	299	187	52	59	1	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-1	GLY	-	expression tag	UNP Q9IK91
D	0	ALA	-	expression tag	UNP Q9IK91
E	-1	GLY	-	expression tag	UNP Q9IK91
E	0	ALA	-	expression tag	UNP Q9IK91
F	-1	GLY	-	expression tag	UNP Q9IK91
F	0	ALA	-	expression tag	UNP Q9IK91

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
3	A	5	5	5	0	0
			Total	Cl		
3	B	2	2	2	0	0

- Molecule 4 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Br		
4	A	1	1	1	0	0
			Total	Br		
4	C	1	1	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	42	42	42	0	0
			Total	O		
5	B	33	33	33	0	0
			Total	O		
5	C	17	17	17	0	0
			Total	O		
5	D	2	2	2	0	0
			Total	O		
5	E	3	3	3	0	0
			Total	O		
5	F	2	2	2	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NUCLEOPROTEIN

Chain A:



- Molecule 1: NUCLEOPROTEIN

Chain B:



- Molecule 1: NUCLEOPROTEIN

Chain C:







- Molecule 2: PHOSPHOPROTEIN

Chain D:



- Molecule 2: PHOSPHOPROTEIN

Chain E:



- Molecule 2: PHOSPHOPROTEIN

Chain F:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	82.89Å 98.96Å 156.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.19 – 2.50 47.19 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.5 (47.19-2.50) 99.6 (47.19-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.00 (at 2.51Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.192 , 0.259 0.203 , 0.264	Depositor DCC
R_{free} test set	2288 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å^2)	46.9	Xtriage
Anisotropy	0.618	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.31 , 49.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	7650	wwPDB-VP
Average B, all atoms (Å^2)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.52% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BR, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.48	0/2635	0.60	0/3536
1	B	0.47	0/2603	0.58	0/3487
1	C	0.45	0/1458	0.57	0/1958
2	D	0.46	0/319	0.51	0/426
2	E	0.41	0/311	0.53	0/415
2	F	0.45	0/301	0.54	0/402
All	All	0.47	0/7627	0.58	0/10224

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2599	0	2669	42	0
1	B	2574	0	2645	42	0
1	C	1444	0	1322	23	0
2	D	317	0	327	3	0
2	E	309	0	316	4	0
2	F	299	0	308	8	0
3	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	2	0	0	1	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	42	0	0	0	0
5	B	33	0	0	0	0
5	C	17	0	0	1	0
5	D	2	0	0	0	0
5	E	3	0	0	0	0
5	F	2	0	0	1	0
All	All	7650	0	7587	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 115 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:193:ARG:HD2	1:A:194:TRP:H	1.34	0.93
1:A:335[B]:MSE:SE	1:A:348:LEU:HG	2.26	0.85
1:A:32:THR:OG1	1:A:33:THR:N	2.10	0.83
1:C:302:ARG:NH1	2:E:8:ASN:OD1	2.13	0.82
1:B:42:THR:HG23	1:B:44:SER:H	1.48	0.78

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	325/356 (91%)	314 (97%)	10 (3%)	1 (0%)	41 61
1	B	314/356 (88%)	304 (97%)	9 (3%)	1 (0%)	41 61

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	191/356 (54%)	184 (96%)	7 (4%)	0	100	100
2	D	38/52 (73%)	38 (100%)	0	0	100	100
2	E	37/52 (71%)	37 (100%)	0	0	100	100
2	F	35/52 (67%)	35 (100%)	0	0	100	100
All	All	940/1224 (77%)	912 (97%)	26 (3%)	2 (0%)	47	68

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	140	LYS
1	B	343	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/278 (101%)	259 (92%)	21 (8%)	13	26
1	B	279/278 (100%)	257 (92%)	22 (8%)	12	24
1	C	128/278 (46%)	119 (93%)	9 (7%)	15	29
2	D	36/47 (77%)	33 (92%)	3 (8%)	11	22
2	E	35/47 (74%)	33 (94%)	2 (6%)	20	39
2	F	34/47 (72%)	31 (91%)	3 (9%)	10	19
All	All	792/975 (81%)	732 (92%)	60 (8%)	13	25

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	139	SER
2	E	2	ASP
1	B	259	VAL
2	D	38	ILE
2	F	35	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	122	GLN
1	C	211	GLN
1	C	257	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	315/356 (88%)	0.11	11 (3%)	44	47	27, 53, 98, 155	0
1	B	309/356 (86%)	0.07	4 (1%)	77	79	28, 56, 93, 122	0
1	C	190/356 (53%)	1.13	50 (26%)	0	0	31, 81, 131, 165	0
2	D	40/52 (76%)	0.02	1 (2%)	57	61	35, 55, 104, 121	0
2	E	39/52 (75%)	0.14	3 (7%)	13	13	43, 58, 96, 114	0
2	F	37/52 (71%)	0.20	1 (2%)	54	58	34, 55, 97, 106	0
All	All	930/1224 (75%)	0.31	70 (7%)	14	14	27, 56, 117, 165	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	251	ILE	7.8
1	C	157	ALA	7.7
1	C	75	THR	7.6
1	C	220	LEU	7.5
1	C	207	PHE	7.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	CL	A	1375	1/1	0.94	0.28	48,48,48,48	0
3	CL	B	1371	1/1	0.94	0.17	56,56,56,56	0
3	CL	A	1377	1/1	0.97	0.17	40,40,40,40	0
3	CL	B	1370	1/1	0.98	0.19	40,40,40,40	0
4	BR	C	1369	1/1	0.98	0.06	66,66,66,66	0
3	CL	A	1374	1/1	0.99	0.12	43,43,43,43	0
3	CL	A	1372	1/1	0.99	0.15	48,48,48,48	0
3	CL	A	1373	1/1	0.99	0.23	41,41,41,41	0
4	BR	A	1376	1/1	1.00	0.12	58,58,58,58	0

6.5 Other polymers (i)

There are no such residues in this entry.